

Cayley graphs with forbidden subgraphs

M. Farrokhi D. G.
(Joint work with A. Mohammadian)

Muroran Institute of Technology

Research on Finite Groups and Their Representations, Vertex
Operator Algebras, and Algebraic Combinatorics
Kyoto University, RIMS
January 7, 2016

Definition

A **Cayley graph** $\text{Cay}(G, S)$ of a group G with respect to a subset $S = S^{-1}$ of $G \setminus \{1\}$ is a graph with vertex set G whose edges are $\{g, sg\}$ for all $g \in G$ and $s \in S$.

Definition

A group G is **Cayley \mathcal{P} -group** (resp. **minimal Cayley \mathcal{P} -group**) if all its Cayley graphs (resp. minimal Cayley graphs) satisfy the property \mathcal{P} .

Theorem (Abdollahi and Jazaeri², 2014; Ahmady, Bell and Mohar³, 2014)

A finite group is *Cayley integral* if and only if it is isomorphic to one of the following groups:

- (1) an *abelian group* of exponent 1, 2, 3, 4 or 6;
- (2) S_3 ;
- (3) $C_3 \rtimes C_4 = \langle x, y : x^3 = y^4 = 1, x^y = x^{-1} \rangle$;
- (4) $Q_8 \times C_2^n$ ($n \geq 0$).

¹F. Harary and A. J. Schwenk, Which Graphs Have Integral Spectra? in: Lecture Notes in Mathematics, 406, Springer, 1974, 45–51.

²A. Abdollahi and M. Jazaeri, Groups all of whose undirected Cayley graphs are integral, *European J. Combin.* **38** (2014), 102–109.

³A. Ahmady, J. Bell and B. Mohar, Integral Cayley graphs and groups, *SIAM J. Discrete Math.* **28**(2) (2014), 685–701.

Theorem (Babai and Sós¹; Godsil and Imrich²)

For every finite graph Γ , the order of a Cayley Γ -free group is bounded above by $(2 + \sqrt{3})|\Gamma|^3$.

Theorem

Let Γ be a finite graph. Then there are only finitely many minimal Cayley Γ -free groups if and only if Γ is a union of paths. Moreover, $|G| < |\Gamma|^{|\Gamma|}$ for any minimal Cayley Γ -free group G when Γ is a union of paths.

¹L. Babai and V. T. Sós, Sidon sets in groups and induced subgraphs of Cayley graphs, *European J. Combin.* **6** (1985), 101–114.


²C. D. Godsil and W. Imrich, Embedding graphs in Cayley graphs, *Graphs Combin.* **3** (1987), 39–43.

Lemma

*Let G be a finite group. A connected Cayley graph $\text{Cay}(G, S)$ is **bipartite** if and only if $[G : \langle S^2 \rangle] = 2$ and $S \subseteq G \setminus \langle S^2 \rangle$.*

Lemma

Let G be a finite group. A connected Cayley graph $\text{Cay}(G, S)$ is *bipartite* if and only if $[G : \langle S^2 \rangle] = 2$ and $S \subseteq G \setminus \langle S^2 \rangle$.



$$\langle S^2 \rangle X = X$$

$$\langle S^2 \rangle Y = Y$$

$$[G : \langle S^2 \rangle] = 2$$

Theorem

*A finite group G is a **minimal Cayley bipartite group** if and only if it is a **2-group**.*

Theorem

*A finite group G is a **minimal Cayley bipartite group** if and only if it is a **2-group**.*

G = minimal Cayley bipartite group:

- $\Rightarrow G \setminus \Phi(G) = \{2\text{-elements}\};$
- $\Rightarrow G = 2\text{-group}.$

Theorem

A finite group G is a *minimal Cayley bipartite group* if and only if it is a *2-group*.

G = minimal Cayley bipartite group:

- $\Rightarrow G \setminus \Phi(G) = \{2\text{-elements}\};$
- $\Rightarrow G = 2\text{-group}.$

$G = 2\text{-group}:$

- $\bullet X = \{x_1, \dots, x_n\}$ (a minimal generating set of G);
- $\bullet S = X \cup X^{-1};$
- $\bullet H = \langle \Phi(G), x_1 x_2, \dots, x_1 x_n \rangle;$
- $\Rightarrow \langle S^2 \rangle = H$ and $S \subseteq G \setminus H;$
- $\Rightarrow \text{Cay}(G, S) = \text{bipartite}.$

Definition

- A **proper n -coloring** of a graph Γ is an assignment of n colors to vertices of Γ such that adjacent vertices have distinct colors;
- The **chromatic number** $\chi(\Gamma)$ of Γ is the minimum number of colors to color Γ properly;
- The **clique number** $\omega(\Gamma)$ of Γ is the maximum size of a complete subgraph of Γ ;
- A graph Γ is **perfect** if $\chi(\Gamma') = \omega(\Gamma')$ for every induced subgraph Γ' of Γ .

Theorem (Strong Perfect Graph Theorem¹)

A graph Γ is *perfect* if neither Γ nor Γ^c has an *induced odd cycle* of length ≥ 5 .

¹M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, *Ann. of Math.* (2) **164**(1) (2006), 51–229.

Notation

Suppose G is a finite group, $H \leq G$ and $g \in G$:

- $\overline{G} = G/\Phi(G)$;
- $\overline{H} = H\Phi(G)/\Phi(G)$;
- $\overline{g} = g\Phi(G)$.

Theorem (MFDG¹)

Let G be a finite solvable group and P be a Sylow p -subgroup of G . If either $\overline{P} \trianglelefteq G$ or \overline{P} is cyclic, then $d(P) = d(\overline{P})$.

¹M. Farrokhi D. G., Finite groups with a given Frattini factor group, *In preparation*.

Definition

The **Hughes-Thompson subgroup** of a group G with respect to a prime p is defined as

$$H_p(G) := \langle g \in G : |g| \neq p \rangle.$$

Theorem

Let G be a finite group and $p \in \pi(G)$. Then

- (1) If G is *not* a p -group then either $H_p(G) = 1$, $H_p(G) = G$ or $[G : H_p(G)] = p$ (Hughes and Thompson¹);
- (1) If G is a p -group ($p = 2, 3$) then either $H_p(G) = 1$, $H_p(G) = G$ or $[G : H_p(G)] = p$ (Straus and Szekeres²).

¹D. R. Hughes and J. G. Thompson, The H_p -problem and the structure of H_p -groups, *Pacific J. Math.* **9** (1959), 1097–1101.

²E. G. Straus and G. Szekeres, On a problem of D. R. Hughes, *Proc. Amer. Math. Soc.* **9**(1) (1958), 157–158.

Theorem (Luccini¹)

The *largest size* of a *minimal generating set* of a finite solvable group equals the number of *non-Frattini factors* of a chief series of the group.

¹A. Luccini, The largest size of a minimal generating set of a finite group, *Arch. Math. (Basel)* **101** (2013), 1–8.

Lemma


Let $G = \langle g \rangle$ be a cyclic group. Then

- (1) $\text{Cay}(G, \{g^{\pm 2}, g^{\pm 3}\})$ has an *induced 5-cycle* for $|g| \geq 10$.
- (2) $\text{Cay}(G, \{g^{\pm 1}, g^{\pm 4}\})$ has an *induced 5-cycle* for $|g| \geq 8$.


Lemma

Let $G = \langle g \rangle$ be a cyclic group. Then

- (1) $\text{Cay}(G, \{g^{\pm 2}, g^{\pm 3}\})$ has an *induced 5-cycle* for $|g| \geq 10$.
- (2) $\text{Cay}(G, \{g^{\pm 1}, g^{\pm 4}\})$ has an *induced 5-cycle* for $|g| \geq 8$.



(1)



(2)

Theorem

A finite group G is a *minimal Cayley perfect group* if and only if either G is a *2-group*, or it is isomorphic to one of the groups C_3 , C_6 , S_3 , $C_3 \times C_3$, A_4 or E_3 .

Theorem

A finite group G is a *minimal Cayley perfect group* if and only if either G is a *2-group*, or it is isomorphic to one of the groups C_3 , C_6 , S_3 , $C_3 \times C_3$, A_4 or E_3 .


$G \neq 2$ -group is a minimal Cayley perfect group:

- $\Rightarrow G \setminus \Phi(G) = \{\text{elements of orders } 2^k, 2^k \cdot 3 \text{ and } 3\};$
 - $Q = \text{Sylow } 3\text{-subgroup of } G;$
- $\Rightarrow \exp(Q) = 3 \text{ or } S_3(\Phi(G)) = H_3(Q);$
- $\Rightarrow \exp(Q) = 3.$

(1) $G = 3$ -group:

- $d(G) \geq 3$;
- $X = \{a, b, c, \dots\} =$ a m.g.s. of G ;

$\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle arising from the relation $(abc)^3 = 1$ (*);



$\Rightarrow d(G) \leq 2$;

$\Rightarrow G \cong C_3, C_3 \times C_3$ or E_3 .

(2) G is a $\{2, 3\}$ -group:

- $\mathcal{C} = \{X : X \cong C_6, S_3 \text{ or } A_4\}$;

$\Rightarrow |X| \in \{6, 12\}$, $S = S^{-1} \ni s$ of order 3 and
 $\text{Cay}(X, S) = \text{perfect}$ imply $X \in \mathcal{C}$;

(2) G is a $\{2, 3\}$ -group:

- $\mathcal{C} = \{X : X \cong C_6, S_3 \text{ or } A_4\}$;
- $\Rightarrow |X| \in \{6, 12\}$, $S = S^{-1} \ni s$ of order 3 and
 $\text{Cay}(X, S) = \text{perfect}$ imply $X \in \mathcal{C}$;

Suppose $G \notin \mathcal{C}$ and $|G| = \text{minimum}$:

- $l_f = \text{the number of non-Frattini factors in a chief series of } G$;
 - $l_f = 2$;
- $\Rightarrow \overline{G} \cong C_6, S_3 \text{ or } A_4$;
- (i) $\overline{G} \cong C_6$:
- $\Rightarrow G = \langle x \rangle$ is cyclic and $\text{Cay}(G, \{x^{\pm 2}, x^{\pm 3}\})$ has an induced 5-cycle if $|G| > 6$;
- (ii) $\overline{G} \cong S_3$:
- $\Rightarrow G = \langle x, y : x^3 = y^{2^k} = 1, x^y = x^{-1} \rangle$ and $\text{Cay}(G, \{x^{\pm 1}, y^{\pm 1}\})$ has an induced $(2^k + 3)$ -cycle determine by $y^{2^k - 2} x y x^{-1} y x = 1$ if $k > 1$;

(iii) $\overline{G} \cong A_4$:

- $\Rightarrow \overline{G} = \langle \overline{x}, \overline{y} : \overline{x}^2 = \overline{y}^3 = (\overline{xy})^3 = \overline{1} \rangle$;
- \Rightarrow one can assume that $(xy)^3 = 1$ or $x^{y^{-1}}x^yx = 1$;
 - $|x| = 2^m > 2$;
 - $\langle a, b : a^{2^m} = b^{2^m} = (ab)^{2^m} = 1 \rangle$ is infinite for $m > 1$;
- \Rightarrow there is a relation $w = 1$ (of minimum length) in x, x^y independent of $x^{2^m} = (x^y)^{2^m} = (x^yx)^{2^m} = 1$;
 - $w = x^{a_1y}x^{b_1} \dots x^{a_ky}x^{b_k}$ with $0 < a_i, b_i < 2^m$ and $(a_1, b_1) \neq (1, 1)$;
 - $w' = x^{-y^{-1}}x^{-1}x^{(a_1-1)y}x^{b_1}x^{a_2y}x^{b_2} \dots x^{a_ky}x^{b_k}$;
- \Rightarrow either $w = 1$ or $w' = 1$ determines an induced odd cycle in $\text{Cay}(G, \{x^{\pm 1}, y^{\pm 1}\})$ of length ≥ 7 (*);
- $\Rightarrow |x| = 2$;
- $\Rightarrow G \cong A_4$.

$l_f \geq 3$:

- Let

$$\Phi(G) = G_0 \trianglelefteq G_1 \trianglelefteq \cdots \trianglelefteq G_{l-1} \trianglelefteq G_l = G,$$

be the inverse image of a chief series of $G/\Phi(G)$ and assume $M = G_{l-1}$

- G_{n_i}/G_{n_i-1} ($i = 1, \dots, l_f$) are the non-Frattini factors;
- $\Rightarrow X = \{x_1, \dots, x_{l_f}\}$ ($x_i \in G_{n_i} \setminus G_{n_i-1}$ is a p -element, $i = 1, \dots, l_f$) is a m.g.s. of G ;
- $x_i, x_j \in S_2(G)$ or $S_3(G)$ if $x_i, x_j = p$ -element ($p = 2, 3$);
 - $Y_i = X \setminus \{x_i\}$ for $i = 1, \dots, l_f$;
 - $x_i \in Y_{j_i}$ (containing elements of even and odd order) for all $i = 1, \dots, l_f$;
- $\Rightarrow |x_i|$ is prime for all $i = 1, \dots, l_f$;

Claim: $l_f = l = 3$:

- $Y_i \ni$ elements of even and odd for $i = 1$ or 2 ;
- $\Rightarrow G = G_{n_2} \langle Y_i \rangle$ implies $G/G_{n_2} \cong \langle Y_i \rangle / (G_{n_2} \cap \langle Y_i \rangle) \cong C_2$ or C_3 ;
- $\Rightarrow n_2 = n_3 - 1$ and $n_3 = l$;
- $\Rightarrow \Phi(G/G_1) = G_{l-2}/G_1$ and $G/G_1 = \langle G_1 x_2, G_1 x_3 \rangle$;
- $(|x_2|, |x_3|) = (2, 3)$ or $(3, 2)$;
- $\Rightarrow \langle x_2, x_3 \rangle \in \mathcal{C}$ so that $l = 3$;
- $(|x_2|, |x_3|) = (2, 2)$;
- $\Rightarrow G/G_1 =$ a dihedral group;
- $\{x_1, x_2 x_3, x_3\}$ is a m.g.s. of G ;
- $\Rightarrow (x_2 x_3)^2 = 1$;
- $\Rightarrow G_{l-2} = \langle G_1, (x_2 x_3)^2 \rangle = G_1$ so that $l = 3$;

- $(|x_2|, |x_3|) = (3, 3);$
 - $\Rightarrow G/G_1 \cong C_3 \times C_3$ or $E_3;$
 - $G/G_1 \cong C_3 \times C_3$ implies $l = 3;$
 - $G/G_1 \cong E_3:$
 - $[x_1, x_2] = [x_1, x_3] = 1$ implies $G = \langle x_1 \rangle \times \langle x_2, x_3 \rangle$ so that $[x_2, x_3] \in \Phi(G) (*)$;
 - $\Rightarrow [x_1, x_3] \neq 1$, say;
 - $\Rightarrow \{x_1, x_2x_3, x_3\}$ is a m.g.s. of G if $[x_1, x_2x_3] \neq 1;$
 - $\Rightarrow \langle x_1, x_2x_3 \rangle \cong A_4;$
 - $\Rightarrow (x_1x_2x_3)^3 = 1$ and $\text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle $(*)$;
 - \Rightarrow Replacing x_2 by x_2x_3 we may assume $[x_1, x_2] = 1;$
 - $\Rightarrow [x_1, x_2x_3^{-1}] \neq 1$, $\{x_1, x_2, x_2x_3^{-1}\}$ is a m.g.s. of G , and
- $$x_1x_1^{x_3^{-1}x_2^{-1}} = (x_1x_1^{x_3^{-1}})^{x_2^{-1}} = x_1^{x_3x_2^{-1}} = x_1^{(x_2x_3^{-1})^{-1}} = x_1x_1^{x_2x_3^{-1}} = x_1x_1^{x_2^{-1}x_3^{-1}};$$
- $\Rightarrow [x_2, x_3]$ commutes with $x_1;$
 - $\Rightarrow G = \langle x_1, x_1^{x_3} \rangle \rtimes \langle x_2, x_3 \rangle;$
 - $\Rightarrow [x_2, x_3] \in \Phi(G) (*)$;

Structure of $\Phi(G)$:

- x_u =involution;
- $\Rightarrow gx_u$ =involution for all $g \in \Phi(G)$ so that $g^{x_u} = g^{-1}$;
- $\Rightarrow \Phi(G)$ =abelian;

- $\langle x_u, x_v \rangle \cong A_4$ for some x_v ;
- $\Rightarrow g^{x_u^{x_v^{\pm 1}}} = g^{-1}$ for all $g \in \Phi(G)$ ($x_u \mapsto x_u^{x_v^{\pm 1}}$);
- $\Rightarrow g^{-1} = g^{x_u^{x_v^{-1}}} = g^{x_u x_v} = g$;
- $\Rightarrow \Phi(G)$ =elementary abelian 2-group.

$$H = \langle Y_3 \rangle$$

$$\Rightarrow \text{Cay}(H, Y_3 \cup Y_3^{-1}) = \text{perfect};$$

$$H = \langle Y_3 \rangle$$

$$\Rightarrow \text{Cay}(H, Y_3 \cup Y_3^{-1}) = \text{perfect};$$

$$H = 2\text{-group};$$

$$\Rightarrow [G : M] = 3, M = S_2(G) \text{ and } \Phi(G) = \Phi(M) \text{ so that } \overline{M} = \text{elementary abelian};$$

$$\Rightarrow \langle Y_i \rangle \cong C_6 \text{ or } A_4 \text{ for } i = 1, 2;$$

- $\langle Y_1 \rangle \cong \langle Y_2 \rangle \cong C_6$:

$$\Rightarrow M = H, G/\Phi(G) \cong C_6 \times C_2 \text{ so that } G \text{ is nilpotent};$$

- $M \setminus \Phi(M) \ni \text{an element of order } \geq 4 \text{ implies } G \setminus \Phi(G) \ni \text{an element } x \text{ of order } \geq 12 (*);$

$$\Rightarrow M \setminus \Phi(M) \text{ contains only involutions};$$

$$\Rightarrow M \cong C_2 \times C_2 \text{ so that } G \cong C_6 \times C_2 (*);$$

- $\langle Y_i \rangle \cong A_4$ for some $i = 1, 2$:
 - $\Rightarrow \Phi(G)$ = elementary abelian 2-group;
 - $\langle Y_j \rangle \cong C_6$ for $j \in \{1, 2\} \setminus \{i\}$:
 - $\langle x_2, x_3 \rangle \cong A_4$;
 - $\Rightarrow \{x_1, x_2, x_2x_3\}$ is a m.g.s of G and $|x_2x_3| = 3$;
 - $\Rightarrow [x_1, x_2x_3] \neq 1$ implies $(x_1x_2x_3)^3 = 1$ so that $\text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle (*);
 - $\Rightarrow [x_1, x_2x_3] = 1$;
 - \Rightarrow Replacing x_3 by x_2x_3 one can assume that $[x_1, x_3] = 1$ and $\langle x_1, x_3 \rangle \cong C_6$;
 - $\Rightarrow (gx_i)^{x_3^{-1}} = (gx_i)(gx_i)^{x_3}$ and $(gx_j)^{x_3} = (gx_j)$ for all $g \in \Phi(G)$ ($((x_i, x_j) \mapsto (gx_i, gx_j))$);
 - $\Rightarrow g^{x_3^{-1}} = gg^{x_3} = 1$ so that $\Phi(G) = 1$;
 - $\Rightarrow G \cong A_4 \times C_2$;
 - $\Rightarrow \text{Cay}(G, \{a^{\pm 1}, b^{\pm 1}\})$ has an induced 7-cycle arising from $b^{-1}abab^2a^{-1} = 1$, in which $a := x_3^{x_i}$ and $b := x_jx_3$ (*);

H is a 3-group:

- $\Rightarrow [G : M] = 2$ and $M = S_3(G)$;
- $\Rightarrow \langle Y_i \rangle \cong C_6$ or S_3 for $i = 1, 2$;
- $\Rightarrow x_i^{x_3} = x_i^{\epsilon_i}$ with $\epsilon_i = \pm 1$ for $i = 1, 2$;
- $\Rightarrow |M| = |H| = 9$ or 27 ;
- $\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 7-cycle arising from $x_3 x_2^{\epsilon_2} x_1^{-\epsilon_1} x_3 x_2 x_1 x_2 = 1$ if $[x_1, x_2] = 1$ (*);
- $\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 11-cycle arising from $x_3 x_2^{\epsilon_2} x_1^{-\epsilon_1} x_3 x_2 x_1 x_2 x_1^{-1} x_2^{-1} x_1 x_2 = 1$ if $[x_1, x_2] \neq 1$ (*).

$H = \{2, 3\}$ -group;

$\Rightarrow H \in \mathcal{C}$ so that $H \cong C_6, S_3$ or A_4 ;

- $|x_2| = |x_3| = 2$:

$\Rightarrow [x_2, x_3] = 1$ and $x_1^{x_2}, x_1^{x_3} \in \langle x_1 \rangle$;

$\Rightarrow G \cong C_6 \times C_2$ or $S_3 \times C_2$ (*);

- $|x_2| = |x_3| = 3$:

$\Rightarrow x_1$ commutes with $[x_2, x_3]$ and x_3 (say);

- $[x_1, x_2] = 1$ implies $G = \langle x_1 \rangle \times \langle x_2, x_3 \rangle$;

$\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle arising from $x_1 x_2 x_3 x_2^{-1} x_1 x_2^{-1} x_3 x_2 x_3 = 1$ (*);

- $[x_1, x_2] \neq 1$;

$\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle arising from $(x_1 x_2 x_3)^2 x_1 x_3 x_2 = 1$ if $[x_2, x_3] = 1$ (*);

$\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 13-cycle arising from $(x_1 x_2 x_3)^2 x_2^{-1} x_3^{-1} x_2 x_3 x_1 x_3 x_2 = 1$ if $[x_2, x_3] \neq 1$ (*);

- $|x_2| \neq |x_3|$:
 - $[x_2, x_3] = 1$;
 - \Rightarrow Swapping x_2 and x_3 we get $H = p$ -group ($p \in \{2, 3\}$) (\ast);
 - $\Rightarrow [x_2, x_3] \neq 1$;
 - $[x_1, x_2] = 1$;
 - $\Rightarrow \langle x_2, x_2^{x_3}, x_2^{x_3^{-1}} \rangle \trianglelefteq G$ is elementary abelian;
 - \Rightarrow Swapping x_1 and x_2 we get $H = p$ -group ($p \in \{2, 3\}$) (\ast);
 - $\Rightarrow [x_1, x_2] \neq 1$;
 - $\Rightarrow \{\langle x_1, x_2 \rangle, \langle x_2, x_3 \rangle\} = \{S_3, A_4\}$;
 - $\Rightarrow \Phi(G) = \text{elementary abelian 2-group}$;

- $(|x_1|, |x_2|, |x_3|) = (2, 3, 2)$:
 - $\Rightarrow \langle x_1, x_2 \rangle \cong A_4$ and $\langle x_2, x_3 \rangle \cong S_3$;
 - $\Rightarrow |x_1 x_3| = 2^m$ ($\langle x_1, x_3 \rangle$ =dihedral 2-group);
 - $\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced $(2^{m+1} + 5)$ -cycle arising from $(x_1 x_2)^2 (x_3 x_1)^{2^m - 1} x_2 x_3 x_2^{-1} = 1$ (*);

- $(|x_1|, |x_2|, |x_3|) = (2, 3, 2)$:
 - $\Rightarrow \langle x_1, x_2 \rangle \cong A_4$ and $\langle x_2, x_3 \rangle \cong S_3$;
 - $\Rightarrow |x_1 x_3| = 2^m$ ($\langle x_1, x_3 \rangle$ = dihedral 2-group);
 - $\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced $(2^{m+1} + 5)$ -cycle arising from $(x_1 x_2)^2 (x_3 x_1)^{2^m - 1} x_2 x_3 x_2^{-1} = 1$ (*);
- $(|x_1|, |x_2|, |x_3|) = (3, 2, 3)$:
 - $\Rightarrow \langle x_1, x_2 \rangle \cong S_3$ and $\langle x_2, x_3 \rangle \cong A_4$;
 - $\Rightarrow |x_1 x_3| = 3$;
 - $Q = S_3(G) \ni x_2 x_3$;
 - $y \in \Phi(G) \langle x_2, x_2^{x_3} \rangle = S_2(G)$ such that $x_1^y \in Q$;
 - $\Rightarrow (x_1 x_2 x_3)^3 = 1$ ($x_1 \mapsto x_1^y$);
 - $\Rightarrow \text{Cay}(G, X \cup X^{-1})$ has an induced 9-cycle (*).

Corollary

Let G be a nontrivial finite group. Then G is a **Cayley perfect group** if and only if G is isomorphic to one of the groups C_2 , C_3 , C_4 , $C_2 \times C_2$, S_3 , C_6 , $C_2 \times C_2 \times C_2$, $C_2 \times C_4$, D_8 , Q_8 or $C_3 \times C_3$.

G	S	5-cycle
$\langle a \rangle \times \langle b \rangle, a = b = 4$	$\{a^{\pm 1}, b^{\pm 1}, a^2 b^2\}$	$1, a^{-1}, (ab)^{-1}, ab, a, 1$
$\langle a, b : a^4 = b^4 = 1, [a, b] = a^2 \rangle$	$\{a^{\pm 1}, b^{\pm 1}, a^2 b^2\}$	$1, a^{-1}, (ba)^{-1}, ab, a, 1$
$\langle a \rangle \times \langle b \rangle \times \langle c \rangle, a = 4, b = c = 2$	$\{a^{\pm 1}, b, c, a^2 bc\}$	$1, c, ca^{-1}, ba, b, 1$
$\langle a, b, c : a^4 = b^2 = c^2 = 1, a^c = a^{-1}b \rangle$	$\{a^{\pm 1}, b, c\}$	$1, c, ca^{-1}, ba, b, 1$
$\langle a, b, c : b^2 = c^2 = [a, b] = [a, c] = 1, [b, c] = a^2 \rangle$	$\{a^{\pm 1}, b, c, a^2 bc\}$	$1, a^{-1}, a^{-1}c, ab, a, 1$
$\langle a, b : a^4 = b^2 = 1, a^b = a^{-1} \rangle \times \langle c \rangle, c = 2$	$\{a^{\pm 1}, b, c, a^2 bc\}$	$1, a^{-1}, a^{-1}c, ab, a, 1$
$\langle a, b : a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle \times \langle c \rangle, c = 2$	$\{a^{\pm 1}, b^{\pm 1}, c, (a^2 b^{-1} c)^{\pm 1}\}$	$1, a^{-1}, a^{-1}c, ab, a, 1$
$\langle a \rangle \times \langle b \rangle \times \langle c \rangle \times \langle d \rangle, a = b = c = d = 2$	$\{a, b, c, d, abcd\}$	$1, a, ab, abc, abcd, 1$
$\langle a, b, c : a^3 = b^3 = [a, c] = [b, c] = 1, c = [a, b] \rangle$	$\{a^{\pm 1}, b^{\pm 1}, c^{\pm 1}, (a^{-1}bc)^{\pm 1}\}$	$1, a, ac, bc^{-1}, b, 1$
$\langle a, b : a^2 = b^3 = (ab)^3 = 1 \rangle$	$\{a, b^{\pm 1}, a^b\}$	$1, a^b, b^a, ab, a, 1$

Thank You for Your Attention!