


GEOMETRIC GROUP THEORY

AMENABLE GROUPS

M. FARROKHI D. G.

INSTITUTE FOR ADVANCED STUDIES
IN BASIC SCIENCES

AUGUST 21, 2024



FREE GROUPS

WHAT IS A GROUP?

A group is a **set** of **isometries** of a **mathematical space** closed under **composition** and **inverses** of elements.

WHAT IS A GROUP?

A group is a **set** of **isometries** of a **mathematical space** closed under **composition** and **inverses** of elements.

or

WHAT IS A GROUP?

A group is a **set** of **isometries** of a **mathematical space** closed under **composition** and **inverses** of elements.

or

A group is a **set** equipped with a **binary operation** satisfying the **associativity law**, containing a **neutral** element, and the **inverses** of its elements.

GENERATION

Let G be a group.

- If $a \in G$, then the set of elements

$$\dots, a^{-2}, a^{-1}, 1, a, a^2, \dots$$

form the group $\langle a \rangle$ **generated** by a .

GENERATION

Let G be a group.

- If $a \in G$, then the set of elements

$$\dots, a^{-2}, a^{-1}, 1, a, a^2, \dots$$

form the group $\langle a \rangle$ **generated** by a .

- If $a, b \in G$, then the set of elements

$$\begin{aligned} &1, \\ &a, a^{-1}, b, b^{-1}, \\ &a^2, ab, ab^{-1}, a^{-2}, a^{-1}b, a^{-1}b^{-1}, b^2, ba, ba^{-1}, b^{-2}, b^{-1}a, b^{-1}a^{-1}, \\ &\vdots \end{aligned}$$

form a the group $\langle a, b \rangle$ **generated** by a, b .

PRESENTATION OF GROUPS


$$\blacksquare \mathbb{Z} = \langle 1 \rangle \cong \langle a \rangle,$$

PRESENTATION OF GROUPS

- $\mathbb{Z} = \langle 1 \rangle \cong \langle a \rangle,$
- $\mathbb{Z}_n = \langle [1] \mid [n] = [0] \rangle \cong \langle a \mid a^n = 1 \rangle,$


PRESENTATION OF GROUPS

- $\mathbb{Z} = \langle 1 \rangle \cong \langle a \rangle,$
- $\mathbb{Z}_n = \langle [1] \mid [n] = [0] \rangle \cong \langle a \mid a^n = 1 \rangle,$
- $D_n = \langle a, b \mid a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle,$



PRESENTATION OF GROUPS


- $\mathbb{Z} = \langle 1 \rangle \cong \langle a \rangle,$
- $\mathbb{Z}_n = \langle [1] \mid [n] = [0] \rangle \cong \langle a \mid a^n = 1 \rangle,$
- $D_n = \langle a, b \mid a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle,$



- $S_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i^2 = (\sigma_i \sigma_{i+1})^3 = [\sigma_i, \sigma_j] = 1, |i - j| > 1 \rangle$

PRESENTATION OF GROUPS

- $\mathbb{Z} = \langle 1 \rangle \cong \langle a \rangle$,
- $\mathbb{Z}_n = \langle [1] \mid [n] = [0] \rangle \cong \langle a \mid a^n = 1 \rangle$,
- $D_n = \langle a, b \mid a^n = b^2 = 1, b^{-1}ab = a^{-1} \rangle$,



- $S_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i^2 = (\sigma_i \sigma_{i+1})^3 = [\sigma_i, \sigma_j] = 1, |i - j| > 1 \rangle$
- $F(X) := \langle X \mid \emptyset \rangle$ is a **free group**.

PRESENTATION OF GROUPS

Definition

A **presentation** of a group G with

- generating set X , and
- relation set R

is denoted by $\langle X \mid R \rangle$.

PRESENTATION OF GROUPS

Definition

A **presentation** of a group G with

- generating set X , and
- relation set R

is denoted by $\langle X \mid R \rangle$.

Theorem

If $G = \langle X \mid R \rangle$, then

$$G = F/N,$$

where $F := F(X)$ and

$$N = \langle frf^{-1} \mid r \in R, f \in F \rangle$$

is the **normal closure** of R in F .

CONSTRUCTION OF FREE GROUPS

- Subgroups of free groups,

CONSTRUCTION OF FREE GROUPS

- Subgroups of free groups,
- Ping-pong lemma,

CONSTRUCTION OF FREE GROUPS

- Subgroups of free groups,
- Ping-pong lemma,
- Bijective functions, say $\langle f, g \rangle$ with

$$f(x) := x + 1 \quad \text{and} \quad g(x) := \frac{1}{x},$$

CONSTRUCTION OF FREE GROUPS

- Subgroups of free groups,
- Ping-pong lemma,
- Bijective functions, say $\langle f, g \rangle$ with

$$f(x) := x + 1 \quad \text{and} \quad g(x) := \frac{1}{x},$$

- Invertible matrices, say $\langle A, B \rangle$ with

$$A := \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad B := \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix},$$

CONSTRUCTION OF FREE GROUPS

- Subgroups of free groups,
- Ping-pong lemma,
- Bijective functions, say $\langle f, g \rangle$ with


$$f(x) := x + 1 \quad \text{and} \quad g(x) := \frac{1}{x},$$

- Invertible matrices, say $\langle A, B \rangle$ with


$$A := \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad B := \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix},$$

- Rotations, say $\langle \text{Rot}_x(\alpha), \text{Rot}_y(\beta) \rangle$, where
 - $\text{Rot}_x(\alpha)$ is a rotation around **x-axis** by angle α
 - $\text{Rot}_y(\beta)$ is a rotation around **y-axis** by angle βin \mathbb{R}^n ($n \geq 3$) and α, β are chosen **suitably**.


DIAGRAM OF THE FREE GROUP $F(a, b)$



BANACH-TARSKI PARADOX




BANACH-TARSKI PARADOX



$$F = \{1\} \sqcup R \sqcup L \sqcup U \sqcup D$$


BANACH-TARSKI PARADOX



$$F = \{1\} \sqcup R \sqcup L \sqcup U \sqcup D$$

$$F = a^{-1}R \sqcup L$$

BANACH-TARSKI PARADOX




$$F = \{1\} \sqcup R \sqcup L \sqcup U \sqcup D$$

$$F = a^{-1}R \sqcup L$$

$$F = aL \sqcup R$$

BANACH-TARSKI PARADOX




$$F = \{1\} \sqcup R \sqcup L \sqcup U \sqcup D$$

$$F = a^{-1}R \sqcup L$$

$$F = aL \sqcup R$$

$$F = b^{-1}U \sqcup D$$

BANACH-TARSKI PARADOX



$$F = \{1\} \sqcup R \sqcup L \sqcup U \sqcup D$$

$$F = a^{-1}R \sqcup L$$

$$F = aL \sqcup R$$

$$F = b^{-1}U \sqcup D$$

$$F = bD \sqcup U$$

■ $\theta := \arctan(1/3),$

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3),$
- $a := \text{Rot}_x(\theta), b := \text{Rot}_y(\theta) \text{ in } \mathbb{R}^3,$

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,
- F **acts** on S^2 ,

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,
- F **acts** on S^2 ,
- $Fp := \{f(p) : f \in F\}$ is the **orbit** of $p \in S^2$,

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,
- F **acts** on S^2 ,
- $Fp := \{f(p) : f \in F\}$ is the **orbit** of $p \in S^2$,
- Orbits of points of S^2 **partition** S^2 ,

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,
- F **acts** on S^2 ,
- $Fp := \{f(p) : f \in F\}$ is the **orbit** of $p \in S^2$,
- Orbits of points of S^2 **partition** S^2 ,
- Let M be a set of representatives of **orbits** of F on S^2 ,

BANACH-TARSKI PARADOX

- $\theta := \arctan(1/3)$,
- $a := \text{Rot}_x(\theta)$, $b := \text{Rot}_y(\theta)$ in \mathbb{R}^3 ,
- $F := \langle a, b \rangle$ is **free**,
- F **acts** on S^2 ,
- $Fp := \{f(p) : f \in F\}$ is the **orbit** of $p \in S^2$,
- Orbits of points of S^2 **partition** S^2 ,
- Let M be a set of representatives of **orbits** of F on S^2 ,
- $S^2 = M \cup RM \cup LM \cup UM \cup DM$,

BANACH-TARSKI PARADOX

■ $N := \{a^{-1}, a^{-2}, \dots\}M,$

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$
- Let

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$
- Let
 - ▶ $\Delta_1 := RM \cup M \cup N,$

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$
- Let
 - ▶ $\Delta_1 := RM \cup M \cup N,$
 - ▶ $\Delta_2 := LM \setminus N,$

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$

- Let

- ▶ $\Delta_1 := RM \cup M \cup N,$

- ▶ $\Delta_2 := LM \setminus N,$

- ▶ $\Delta_3 := UM,$

BANACH-TARSKI PARADOX

■ $N := \{a^{-1}, a^{-2}, \dots\}M,$

■ Let

▶ $\Delta_1 := RM \cup M \cup N,$

▶ $\Delta_2 := LM \setminus N,$

▶ $\Delta_3 := UM,$

▶ $\Delta_4 := DM,$

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$
- Let
 - ▶ $\Delta_1 := RM \cup M \cup N,$
 - ▶ $\Delta_2 := LM \setminus N,$
 - ▶ $\Delta_3 := UM,$
 - ▶ $\Delta_4 := DM,$
- $S^2 = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4.$

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$

- Let

- ▶ $\Delta_1 := RM \cup M \cup N,$

- ▶ $\Delta_2 := LM \setminus N,$

- ▶ $\Delta_3 := UM,$

- ▶ $\Delta_4 := DM,$

- $S^2 = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4.$

- Then

BANACH-TARSKI PARADOX

- $N := \{a^{-1}, a^{-2}, \dots\}M,$

- Let

- ▶ $\Delta_1 := RM \cup M \cup N,$

- ▶ $\Delta_2 := LM \setminus N,$

- ▶ $\Delta_3 := UM,$

- ▶ $\Delta_4 := DM,$

- $S^2 = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4.$

- Then

- ▶ $a\Delta_2 = \Delta_2 \cup \Delta_3 \cup \Delta_4,$

BANACH-TARSKI PARADOX

■ $N := \{a^{-1}, a^{-2}, \dots\}M,$

■ Let

▶ $\Delta_1 := RM \cup M \cup N,$

▶ $\Delta_2 := LM \setminus N,$

▶ $\Delta_3 := UM,$

▶ $\Delta_4 := DM,$

■ $S^2 = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4.$

■ Then

▶ $a\Delta_2 = \Delta_2 \cup \Delta_3 \cup \Delta_4,$

▶ $b\Delta_4 = \Delta_1 \cup \Delta_2 \cup \Delta_4.$

BANACH-TARSKI PARADOX

■ $N := \{a^{-1}, a^{-2}, \dots\}M,$

■ Let

▶ $\Delta_1 := RM \cup M \cup N,$

▶ $\Delta_2 := LM \setminus N,$

▶ $\Delta_3 := UM,$

▶ $\Delta_4 := DM,$

■ $S^2 = \Delta_1 \cup \Delta_2 \cup \Delta_3 \cup \Delta_4.$

■ Then

▶ $a\Delta_2 = \Delta_2 \cup \Delta_3 \cup \Delta_4,$

▶ $b\Delta_4 = \Delta_1 \cup \Delta_2 \cup \Delta_4.$

■ Then

$$\begin{array}{c} \Delta_1 \\ \Delta_2 \\ \Delta_3 \\ \Delta_4 \end{array} \implies \begin{array}{c} \Delta_1 \\ a\Delta_2 \\ \Delta_3 \\ b\Delta_4 \end{array} \implies \begin{array}{c} \Delta_1 \\ \Delta_2 \\ \Delta_3 \\ \Delta_4 \end{array} \cup \begin{array}{c} \Delta_1 \\ \Delta_2 \\ \Delta_3 \\ \Delta_4 \end{array}$$

Definition (John von Neumann, 1929)

A finitely presented discrete group G is **amenable** if it admits a **probability measure** μ on 2^G satisfying

- $\mu(G) = 1$,
- $\mu(X \cup Y) = \mu(X) + \mu(Y)$,
- $\mu(gX) = \mu(X)$

for all disjoint $X, Y \subseteq G$ and $g \in G$.

AMENABLE GROUPS

Definition (John von Neumann, 1929)

A finitely presented discrete group G is **amenable** if it admits a **probability measure** μ on 2^G satisfying

- $\mu(G) = 1$,
- $\mu(X \cup Y) = \mu(X) + \mu(Y)$,
- $\mu(gX) = \mu(X)$

for all disjoint $X, Y \subseteq G$ and $g \in G$.

Example

- **Finite groups** are amenable,

AMENABLE GROUPS

Definition (John von Neumann, 1929)

A finitely presented discrete group G is **amenable** if it admits a **probability measure** μ on 2^G satisfying

- $\mu(G) = 1$,
- $\mu(X \cup Y) = \mu(X) + \mu(Y)$,
- $\mu(gX) = \mu(X)$

for all disjoint $X, Y \subseteq G$ and $g \in G$.

Example

- **Finite groups** are amenable,
- **Abelian groups** are amenable (John von Neumann, 1929),

AMENABLE GROUPS

Definition (John von Neumann, 1929)

A finitely presented discrete group G is **amenable** if it admits a **probability measure** μ on 2^G satisfying

- $\mu(G) = 1$,
- $\mu(X \cup Y) = \mu(X) + \mu(Y)$,
- $\mu(gX) = \mu(X)$

for all disjoint $X, Y \subseteq G$ and $g \in G$.

Example

- **Finite groups** are amenable,
- **Abelian groups** are amenable (John von Neumann, 1929),
- **Non-abelian free groups** are not amenable.

Definition

A **class** \mathcal{X} of groups is a collection of groups closed under isomorphism.

Definition

A **class** \mathcal{X} of groups is a collection of groups closed under isomorphism.

- \mathcal{AG} is the class of amenable groups,

Definition

A **class** \mathcal{X} of groups is a collection of groups closed under isomorphism.

- \mathcal{AG} is the class of amenable groups,
- \mathcal{NF} is the class of groups having no non-abelian free subgroups.

Definition

A **class \mathcal{X}** of groups is a collection of groups closed under isomorphism.

- \mathcal{AG} is the class of amenable groups,
- \mathcal{NF} is the class of groups having no non-abelian free subgroups.

Conjecture (John von Neumann, 1929)

$$\mathcal{AG} = \mathcal{NF}.$$

Definition

A **Tarski monster p -group** is a non-cyclic group with all non-trivial subgroups **cyclic** of order p .

Definition

A **Tarski monster p -group** is a non-cyclic group with all non-trivial subgroups **cyclic** of order p .

Example (Alexander Ol'shanskii, 1980)

Definition

A **Tarski monster p -group** is a non-cyclic group with all non-trivial subgroups **cyclic** of order p .

Example (Alexander Ol'shanskii, 1980)

- Tarski monster p -groups **exists** for all prime $p > 10^{75}$.

Definition


A **Tarski monster p -group** is a non-cyclic group with all non-trivial subgroups **cyclic** of order p .

Example (Alexander Ol'shanskii, 1980)


- Tarski monster p -groups **exists** for all prime $p > 10^{75}$.
- Tarski monster p -groups are **not amenable**.

RICHARD THOMPSON'S GROUP F

FIRST CONSTRUCTION: ROOTED BINARY TREES




FIRST CONSTRUCTION: TOOTED BINARY TREES




FIRST CONSTRUCTION: TREE DIAGRAMS

Domain tree




Range tree




(T_1, T_2)

FIRST CONSTRUCTION: REDUCTIONS




(T_1, T_2)

FIRST CONSTRUCTION: REDUCTIONS




(T_1, T_2)

FIRST CONSTRUCTION: REDUCTIONS




(T'_1, T'_2)

FIRST CONSTRUCTION: REDUCTIONS



(T'_1, T'_2)

FIRST CONSTRUCTION: REDUCTIONS




(T''_1, T''_2)

Lemma


- Every *tree diagram* has a *unique reduced* tree diagrams.
- Having *same reduced* tree diagrams is an *equivalence relation* on the set of all tree diagrams.

FIRST CONSTRUCTION: COMBINATION




$$[(T_1, T_2)] \circ [(S_1, S_2)]$$

FIRST CONSTRUCTION: COMBINATION (GRAFTING)



$$[(T_1, T_2)] \circ [(S_1, S_2)]$$

FIRST CONSTRUCTION: COMBINATION (CANCELING)



$$[(T_1, T_2)] \circ [(S_1, S_2)] := [(S'_1, T'_2)]$$

Thompson's group F is the set of equivalence classes of tree diagrams along with the given binary operation.

Thompson's group F is the set of equivalence classes of tree diagrams along with the given binary operation.


$[(T, T)]$ is the identity element

Thompson's group F is the set of equivalence classes of tree diagrams along with the given binary operation.


$[(T, T)]$ is the identity element

$$[(T_1, T_2)]^{-1} = [(T_2, T_1)]$$

FIRST CONSTRUCTION: PRESENTATION




$$a = [(S_1, S_2)]$$



$$b = [(T_1, T_2)]$$

FIRST CONSTRUCTION: PRESENTATION




$$a = [(S_1, S_2)]$$

$$b = [(T_1, T_2)]$$

$$F = \langle a, b \mid [ab^{-1}, aba^{-1}] = [ab^{-1}, a^2ba^{-2}] = 1 \rangle$$

SECOND CONSTRUCTION



SECOND CONSTRUCTION

Thompson's group F is the set of all
bijective functions $f : [0, 1] \rightarrow [0, 1]$
that are **linear** on sub-intervals



$$\left[\frac{a_0}{2^n}, \frac{a_1}{2^n}\right], \left[\frac{a_1}{2^n}, \frac{a_2}{2^n}\right], \dots, \left[\frac{a_{m-1}}{2^n}, \frac{a_m}{2^n}\right] \quad (0 = a_0 < \dots < a_m = 2^n)$$

with **slopes**


$$\frac{f(a_i) - f(a_{i-1})}{a_i - a_{i-1}} = 2^{k_i} \quad (k_i \in \mathbb{Z})$$

for all $i = 1, \dots, m$.


SECOND CONSTRUCTION: PRESENTATION



SECOND CONSTRUCTION: PRESENTATION



f



g

$$F = \langle f, g \mid [fg^{-1}, gfg^{-1}] = [fg^{-1}, g^2fg^{-2}] = 1 \rangle$$

Definition

The class \mathcal{EG} of **elementary amenable** groups is the smallest class of groups that

- contains **finite groups** and **abelian groups**, and it is
- closed under **subgroups**, **quotients**, **extensions**, and **direct limits**.

Proposition

$$\mathcal{EG} \subseteq \mathcal{AG}.$$

AMENABILITY



THANKS!