

Factorization number of finite abelian groups

M. Farrokhi D. G.

Ferdowsi University of Mashhad

Tehran

9-10 March 2011

Definition

Let G be a finite group. The group G is *factorized* into subgroups A and B if $G = AB$ and such an expression is called a *factorization* of G . The number of factorizations of G is denoted by $F_2(G)$.

Definition

Let G be a finite group. The group G is *factorized* into subgroups A and B if $G = AB$ and such an expression is called a *factorization* of G . The number of factorizations of G is denoted by $F_2(G)$.

Definition

If G is a finite group, then the *subgroup commutativity degree* of G , $\text{scd}(G)$, is the probability that two subgroups of G commute.

Introduction

Definition

Let G be a finite group. The group G is *factorized* into subgroups A and B if $G = AB$ and such an expression is called a *factorization* of G . The number of factorizations of G is denoted by $F_2(G)$.

Definition

If G is a finite group, then the *subgroup commutativity degree* of G , $scd(G)$, is the probability that two subgroups of G commute.

Connection with subgroup commutativity degree

If G is a finite group, then

$$scd(G) = \frac{1}{|L(G)|^2} \sum_{H \leq G} F_2(H),$$

where $L(G)$ is the set of all subgroups of G .

Known result

The factorization number is known for the following classes of groups:

Known result

The factorization number is known for the following classes of groups:

- 1 projective special linear groups $PSL(2, p^n)$ (Ito, 1953).

Known result

The factorization number is known for the following classes of groups:

- ① projective special linear groups $PSL(2, p^n)$ (Ito, 1953).
- ② the simple groups $G_2(q)$ (Tchakerian and Gentchev, 1985).

Known result

The factorization number is known for the following classes of groups:

- ① projective special linear groups $PSL(2, p^n)$ (Ito, 1953).
- ② the simple groups $G_2(q)$ (Tchakerian and Gentchev, 1985).
- ③ sporadic simple groups (Gentchev, 1986).

Known result

The factorization number is known for the following classes of groups:

- ① projective special linear groups $PSL(2, p^n)$ (Ito, 1953).
- ② the simple groups $G_2(q)$ (Tchakerian and Gentchev, 1985).
- ③ sporadic simple groups (Gentchev, 1986).
- ④ simple groups of Lie type of Lie rank 1 and 2 (Gentchev, 1986).

Known result

The factorization number is known for the following classes of groups:

- ① projective special linear groups $PSL(2, p^n)$ (Ito, 1953).
- ② the simple groups $G_2(q)$ (Tchakerian and Gentchev, 1985).
- ③ sporadic simple groups (Gentchev, 1986).
- ④ simple groups of Lie type of Lie rank 1 and 2 (Gentchev, 1986).
- ⑤ projective special linear groups $PSL(3, p^n)$ and projective special unitary groups $PSU(3, p^n)$ (Blaum, 1983).

Known result

The factorization number is known for the following classes of groups:

- ① projective special linear groups $PSL(2, p^n)$ (Ito, 1953).
- ② the simple groups $G_2(q)$ (Tchakerian and Gentchev, 1985).
- ③ sporadic simple groups (Gentchev, 1986).
- ④ simple groups of Lie type of Lie rank 1 and 2 (Gentchev, 1986).
- ⑤ projective special linear groups $PSL(3, p^n)$ and projective special unitary groups $PSU(3, p^n)$ (Blaum, 1983).
- ⑥ cyclic groups \mathbb{Z}_n , elementary abelian p -groups \mathbb{Z}_p^n , dihedral groups D_{2n} , quasi-dihedral groups QD_{2n} , generalized quaternion groups Q_{4n} and Modular p -group M_{p^n} (Farrokhi and Saeedi, Submitted).

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

① $\mathcal{X}_i = (x_1, \dots, x_i);$

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

- ① $\mathcal{X}_i = (x_1, \dots, x_i)$;
- ② $G_p(\mathcal{X}) = \mathbb{Z}_{p^{x_1}} \times \cdots \times \mathbb{Z}_{p^{x_n}}$ is an abelian p -group of type \mathcal{X} ;

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

- ① $\mathcal{X}_i = (x_1, \dots, x_i)$;
- ② $G_p(\mathcal{X}) = \mathbb{Z}_{p^{x_1}} \times \cdots \times \mathbb{Z}_{p^{x_n}}$ is an abelian p -group of type \mathcal{X} ;
- ③ $S(\mathcal{X}) = \{(y_1, \dots, y_m) : m \leq n, y_m \leq \cdots \leq y_1 \text{ and } y_i \leq x_i, \text{ for } 1 \leq i \leq m\}$;

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

- ① $\mathcal{X}_i = (x_1, \dots, x_i)$;
- ② $G_p(\mathcal{X}) = \mathbb{Z}_{p^{x_1}} \times \dots \times \mathbb{Z}_{p^{x_n}}$ is an abelian p -group of type \mathcal{X} ;
- ③ $S(\mathcal{X}) = \{(y_1, \dots, y_m) : m \leq n, y_m \leq \dots \leq y_1 \text{ and } y_i \leq x_i, \text{ for } 1 \leq i \leq m\}$;
- ④ if $\mathcal{Y} = (y_1, \dots, y_m) \in S(\mathcal{X})$, then $T_p(\mathcal{X} : \mathcal{Y})$ is the set of all m -tuples $(g_1, \dots, g_m) \in G_p(\mathcal{X})^m$ such that $|g_i| = p^{y_i}$ and $\langle g_1, \dots, g_m \rangle \cong G_p(\mathcal{Y})$, for each $i = 1, \dots, m$, and

Notations

Let $\mathcal{X} = (x_1, \dots, x_n)$ be a non-increasing sequence of natural numbers. Then for each $i = 1, \dots, n$

- ① $\mathcal{X}_i = (x_1, \dots, x_i)$;
- ② $G_p(\mathcal{X}) = \mathbb{Z}_{p^{x_1}} \times \dots \times \mathbb{Z}_{p^{x_n}}$ is an abelian p -group of type \mathcal{X} ;
- ③ $S(\mathcal{X}) = \{(y_1, \dots, y_m) : m \leq n, y_m \leq \dots \leq y_1 \text{ and } y_i \leq x_i, \text{ for } 1 \leq i \leq m\}$;
- ④ if $\mathcal{Y} = (y_1, \dots, y_m) \in S(\mathcal{X})$, then $T_p(\mathcal{X} : \mathcal{Y})$ is the set of all m -tuples $(g_1, \dots, g_m) \in G_p(\mathcal{X})^m$ such that $|g_i| = p^{y_i}$ and $\langle g_1, \dots, g_m \rangle \cong G_p(\mathcal{Y})$, for each $i = 1, \dots, m$, and
- ⑤ $\mu_i(\mathcal{X}) = i \max\{j : x_j \geq i\} + x_{\max\{j : x_j \geq i\}+1} + \dots + x_n$,

First formula for factorization number

Lemma

If $\mathcal{A} = (a_1, \dots, a_n)$ is a non-increasing sequence of natural numbers and $\mathcal{B} = (b_1, \dots, b_m) \in S(\mathcal{A})$, then

$$|T_p(\mathcal{A} : \mathcal{B})| = \prod_{i=1}^m \left(p^{\mu_{b_i}(\mathcal{A})} - \frac{p^{\mu_{b_i-1}(\mathcal{A}) + \mu_{b_i}(\mathcal{B}_{i-1})}}{p^{\mu_{b_i-1}(\mathcal{B}_{i-1})}} \right).$$

First formula for factorization number

Lemma

If $\mathcal{A} = (a_1, \dots, a_n)$ is a non-increasing sequence of natural numbers and $\mathcal{B} = (b_1, \dots, b_m) \in S(\mathcal{A})$, then

$$|T_p(\mathcal{A} : \mathcal{B})| = \prod_{i=1}^m \left(p^{\mu_{b_i}(\mathcal{A})} - \frac{p^{\mu_{b_i-1}(\mathcal{A}) + \mu_{b_i}(\mathcal{B}_{i-1})}}{p^{\mu_{b_i-1}(\mathcal{B}_{i-1})}} \right).$$

Corollary

The number of p -subgroups of type \mathcal{B} of a finite abelian p -group of type \mathcal{A} is

$$\binom{\mathcal{A}}{\mathcal{B}}_p = \frac{T_p(\mathcal{A} : \mathcal{B})}{T_p(\mathcal{B} : \mathcal{B})}.$$

First formula for factorization number

Corollary

The number of subgroups of a finite abelian p -group G of type \mathcal{A} is

$$|L(G)| = \sum_{\mathcal{B} \in \mathcal{S}(\mathcal{A})} \binom{\mathcal{A}}{\mathcal{B}}_p.$$

First formula for factorization number

Corollary

The number of subgroups of a finite abelian p -group G of type \mathcal{A} is

$$|L(G)| = \sum_{\mathcal{B} \in S(\mathcal{A})} \binom{\mathcal{A}}{\mathcal{B}}_p.$$

Theorem

If G is a finite abelian p -group of type \mathcal{A} , then

$$F_2(G) = |L(G)|^2 - \sum_{\mathcal{B} \in S(\mathcal{A}) \setminus \{\mathcal{A}\}} \binom{\mathcal{A}}{\mathcal{B}}_p F_2(G_p(\mathcal{B})).$$

Second formula for factorization number

Lemma

Let G be an elementary abelian p -group and $X \leq G$. Then the number of subgroups Y of G of order p^n ($n \leq d(G) - d(X)$) such that $X \cap Y = 1$ is

$$p^{nd(X)} \binom{d(G) - d(X)}{n}_p.$$

Second formula for factorization number

Lemma

Let G be an elementary abelian p -group and $X \leq G$. Then the number of subgroups Y of G of order p^n ($n \leq d(G) - d(X)$) such that $X \cap Y = 1$ is

$$p^{nd(X)} \binom{d(G) - d(X)}{n}_p.$$

Lemma

Let G be an elementary abelian p -group and $X \leq Y \leq G$. Then the number of subgroups Z of G of order $p^{d(G)-d(Y)+n}$ ($n \leq d(Y) - d(X)$) such that $X \cap Z = 1$ and $YZ = G$ is

$$p^{nd(X) + (d(Y) - n)(d(G) - d(Y))} \binom{d(Y) - d(X)}{n}_p.$$

Second formula for factorization number

Theorem

Let G be a finite abelian p -group. Then

$$F_2(G) = \sum_{G^p=AB} \sum_{\substack{0 \leq i, j \leq n \\ n \leq i+j \leq 2n}} \frac{|\Omega_1(G)|^{d(A)+d(B)} |\Omega_1(G^p)|^{i+j}}{|\Omega_1(A)|^{d(A)} |\Omega_1(B)|^{d(B)}} \cdot \frac{p^{(n-i)(n-j)}}{p^{id(A)+jd(B)}} \cdot \binom{n}{i}_p \binom{i}{n-j}_p,$$

where $n = d(\Omega_1(G)) - d(\Omega_1(G^p))$.

Corollary

Let G be a finite abelian p -group of type \mathcal{A} . Then

Corollary

Let G be a finite abelian p -group of type \mathcal{A} . Then

$$\textcircled{1} \quad |\text{Aut}(G)| = |T_p(\mathcal{A} : \mathcal{A})|.$$

Application: Automorphisms

Corollary

Let G be a finite abelian p -group of type \mathcal{A} . Then

- ① $|\text{Aut}(G)| = |T_p(\mathcal{A} : \mathcal{A})|$.
- ② if $\mathcal{A} = (a_1, \dots, a_1, \dots, a_m, \dots, a_m) = (x_1, \dots, x_n)$, where the number of a_i is b_i and $a_i > a_{i+1}$, then

$$|\text{Aut}(G)| = \prod_{i=1}^m \prod_{j=N_{i-1}+1}^{N_i} \left(p^{x_j N_i + x_{N_i+1} + \dots + x_m} - p^{(x_j-1)N_{i'} + x_{N_{i'}+1} + \dots + x_m + j-1} \right)$$

and in particular

$$|\text{Aut}(G)|_p = \prod_{i=1}^m \prod_{j=N_{i-1}+1}^{N_i} p^{(x_j-1)N_{i'} + x_{N_{i'}+1} + \dots + x_m + j-1},$$

where $i' = i + 1 - \text{Sign}(a_{i+1} - a_i + 1)$ and $N_i = b_1 + \dots + b_i$, for each $i = 1, \dots, m$.

Application: Gaussian binomial coefficients

Definition

Let q be a prime power and n be an integer. Then the numbers $[n]_q$ and $[n]_q!$ are called the q -integer and q -factorial and defined as follows:

$$[n]_q = \frac{q^n - 1}{q - 1} \text{ and } [n]_q! = [n]_q [n-1]_q \cdots [2]_q [1]_q.$$

The Gaussian binomial coefficient can be defined in terms of q -factorials by

$$\binom{n}{i}_q = \frac{[n]_q!}{[i]_q! [n-i]_q!}$$

and as usual the Gaussian polynomial coefficients can be defined by

$$\binom{n}{i_1, \dots, i_k}_q = \frac{[n]_q!}{[i_1]_q! \cdots [i_k]_q! [n - i_1 - \cdots - i_k]_q!},$$

where $0 \leq i_1 + \cdots + i_k \leq n$.

Application: Gaussian binomial coefficients

Remark

The Gaussian binomial coefficient $\binom{n}{i}_q$ is the number of subspaces of dimension i in a vector space of dimension n over the field of order q and also appear in the theory of partition of integers.

Application: Gaussian binomial coefficients

Remark

The Gaussian binomial coefficient $\binom{n}{i}_q$ is the number of subspaces of dimension i in a vector space of dimension n over the field of order q and also appear in the theory of partition of integers.

Corollary






Let q be a prime power and n be a natural number. Then

$$\left(\sum_{i=0}^n \binom{n}{i}_q \right)^2 = \sum_{0 \leq i+j+k \leq n} \binom{n}{i, j, k}_q q^{j(n-i-j-k)}.$$







In particular

$$4^n = \sum_{0 \leq i+j+k \leq n} \binom{n}{i, j, k}.$$






References I

-  G. E. Andrews and K. Eriksson, *Integer Partitions*, Cambridge University Press, 2004.
-  W. N. Bailey, *A note on certain q -identities*, Quart. J. Math., **12** (1941), 173-175.
-  J. N. S. Bidwell and M. J. Curran, *Automorphisms of finite Abelian groups*, Math. Proc. R. Ir. Acad., **110A**(10) (2010), 57-71.
-  M. Blaum, *Factorizations of the simple groups $PSL_3(q)$ and $PSU_3(q^2)$* , Arch. Math., **40** (1983), 8-13.
-  H. Cohn, *Projective geometry over F_1 and the Gaussian binomial coefficients*, Amer. Math. Monthly, **111** (2004), 487-495.






References II






-  I. J. Davies, *Enumeration of certain subgroups of abelian p -groups*, Proc. Edinb. Math. Soc. (2), (), 1-4.
-  S. Delsarte, *Fonctions de Mobius sur les groupes Abeliens finis*, Ann. of Math. (2), **49** (1948), 600-609.
-  M. Farrokhi D. G. and F. Saeedi, *Factorization numbers of some finite groups*, Submitted.
-  T. R. Gentchev, *Factorizations of the sporadic simple groups*, Arch. Math., **47** (1986), 97-102.
-  T. R. Gentchev, *Factorizations of the groups of Lie type of Lie rank 1 or 2*, Arch. Math., **47** (1986), 493-499.
-  M. Golasiński and D. L. Gonçalves, *On automorphisms of finite Abelian p -groups*, Math. Slovaca, **58**(4) (2008), 405-412.

References III

-  C. J. Hillar and D. L. Rhea, *Automorphisms of finite abelian groups*, Amer. Math. Monthly, **11** (2007), 17-23.
-  N. J. S. Hughes, *The structure and order of the group of central automorphisms of a finite group*, Proc. London Math. Soc., **52**(2) (1951), 377-385.
-  N. Ito, *On the factorizations of the linear fractional group $LF(2, p^n)$* , Acta Sci. Math. Szeged, **15** (1953), 79-84.
-  F. H. Jackson, *Certain q -identities*, Quart. J. Math., **12** (1941), 167-172.
-  Y. Kinosita, *On an enumeration of certain subgroups of a p -group*, J. Osaka Inst. Sci. Tech., Part I, **1** (1949), 13-20.

References IV

-  J. Konvalina, *Generalized binomial coefficients and the subset-subspace problem*, Adv. in Appl. Math., **21** (1998), 228-240.
-  G. A. Miller, *Form of the number of the prime power subgroups of an abelian group*, Proc. Natl. Acad. Sci. USA, **12** (1926), 470-472.
-  G. A. Miller, *Number of the subgroups of any given abelian group*, Proc. Natl. Acad. Sci. USA, **25** (1939), 258-262.
-  G. A. Miller, *Independent generators of the subgroups of an abelian group*, Proc. Natl. Acad. Sci. USA, **25** (1939), 364-367.
-  P. R. Sanders, *The central automorphisms of a finite group*, J. London Math. Soc, **44** (1969), 225-228.

-  K. Shoda, *Über die Automorphismen einer endlichen Abelschen Gruppe*, Math. Ann., **100** (1928), 674-686.
-  T. Stehling, *On computing the number of subgroups of a finite Abelian group*, Combinatorica, **12**(4) (1992), 475-479.
-  M. Tărnăuceanu, *Subgroup commutativity degrees of finite groups*, J. Algebra, **321**(9) (2009), 2508-2520.
-  K. B. Tchakerian and T. R. Gentchev, *Factorizations of the groups $G_2(q)$* , Arch. Math., **44** (1985), 230-232.
-  Y. Yeh, *On prime power abelian groups*, Bull. Amer. Math. Soc., **54** (1948), 323-327.

Thank You