

Computational Experiment

To demonstrate that the no: zero eigen values of the (unnormalized) graph Laplacian is equal to the no: disconnected components or disconnected subgraphs (or the clusters)

Toy examples

These are toy examples.

Understand the concept using toy examples !

Demonstrate your understanding using toy examples !

Example 1

2 zero eigen values \Rightarrow 2 components

```
clearvars
rng(100)
N = 4; %no: nodes
```

Make a random adjacency matrix(square symmetric)

```
A = rand(N);
A = A'*A;
Z = zeros(size(A));
```

Build the Laplacian matrix of a graph with 2 disconnected components.

```
W = [A Z ; Z A];
W = W-diag(diag(W));



n = size(W,1);

D = diag(sum(W));

L_u = D - W;
L_n = eye(n) - pinv(D)*W;
L_ns = eye(n) - D^(-0.5)*W*D^(-0.5);

subplot(1,2,1)
spy(W)
title("W")

subplot(1,2,2)
plot(graph(W))
title("2 component Graph")
```



```
[eig(L_u) eig(L_n) eig(L_ns)]
```

```
ans = 8x3
-0.0000         0 -0.0000
-0.0000    1.4338    1.4338
 2.7175    1.3618    1.3618
 2.7175    1.2043    1.2043
 3.7223         0 -0.0000
 3.7223    1.4338    1.4338
 3.8766    1.3618    1.3618
 3.8766    1.2043    1.2043
```

Example 2

3 zero eigen values \Rightarrow 3 components

```
clearvars
rng(100)
N = 4; %no: nodes
```

Make a random adjacency matrix(square symmetric)

```
A = rand(N);
A = A'*A;
Z = zeros(size(A));
```

Build the Laplacian matrix of a graph with 3 disconnected components.

```
W = [A Z Z; Z A Z; Z Z A];
W = W-diag(diag(W));
```



```

n = size(W,1);

D = diag(sum(W));
L_u = D - W;
L_n = eye(n) - pinv(D)*W;
L_ns = eye(n) - D^(-0.5)*W*D^(-0.5);

subplot(1,2,1)
spy(W)
title("W")
subplot(1,2,2)
plot(graph(W))
title("3 component Graph")

```



```
[eig(L_u) eig(L_n) eig(L_ns)]
```

```

ans = 12×3
-0.0000    0   -0.0000
-0.0000   1.4338   1.4338
-0.0000   1.3618   1.3618
 2.7175   1.2043   1.2043
 2.7175    0   -0.0000
 2.7175   1.4338   1.4338
 3.7223   1.3618   1.3618
 3.7223   1.2043   1.2043
 3.7223    0   -0.0000
 3.8766   1.4338   1.4338
 3.8766   1.3618   1.3618
 3.8766   1.2043   1.2043
  ⋮
  ⋮
  ⋮

```

Example 3 Using 'blkdiag' function

3 zero eigen values \Rightarrow 3 components

```
clearvars
rng(100)
```

Make a random adjacency matrix(square symmetric)

```
A = rand(4);
A = A'*A;
```

Build the Laplacian matrix of a graph with 3 disconnected components using blkdiag function.

```
W = blkdiag(A,A,A);
W = W-diag(diag(W));



n = size(W,1);

D = diag(sum(W));

L_u = D - W;
L_n = eye(n) - pinv(D)*W;
L_ns = eye(n) - D^(-0.5)*W*D^(-0.5);

subplot(1,2,1)
spy(W)
title("W")

subplot(1,2,2)
plot(graph(W))
title("3 component Graph")
```



```
[eig(L_u) eig(L_n) eig(L_ns)]
```

```
ans = 12x3
-0.0000         0 -0.0000
-0.0000    1.4338    1.4338
-0.0000    1.3618    1.3618
 2.7175    1.2043    1.2043
 2.7175         0 -0.0000
 2.7175    1.4338    1.4338
 3.7223    1.3618    1.3618
 3.7223    1.2043    1.2043
 3.7223         0 -0.0000
 3.8766    1.4338    1.4338
 3.8766    1.3618    1.3618
 3.8766    1.2043    1.2043
  :
  :
```

```
cd( "/media/user/DATA4LINUX/new1/Repos/MFC4_22MAT230/" )
matlab.internal.liveeditor.openAndConvert( 'U1_EigValues_GraphLaplacian.mlx',
'U1_EigValues_GraphLaplacian.m' );
```