

Tentative Title: Predictive capacity of oceanographic indices of recruitment

Megan L. Feddern Kristin Marshall Nick Tolimieri

2026-03-17

Abstract

To be determined...

1 Does predictive capacity criteria select different variables?

1.1 Methods

Recruitment deviations were extracted from the 2023 sablefish stock assessment ([Johnson et al. 2023](#)), the 2023 petrale sole stock assessment ([Taylor et al. 2023](#)), and the 2025 northern yellowtail stock assessment ([Oken et al. 2025](#)). GLORYS derived ocean time series begin in 1993 but for some groundfish lifestages, observations from the previous year is necessary to appropriately align the oceanographic condition with the recruitment estimates (i.e., lagged by one year). Therefore, to ensure a standardized length of time series were used for all analyses, species, and oceanographic predictors that did not include late recruitment deviations models were fit to the time period 1994 - 2018. The exponentiated recruitment deviations are the multiplicative factor by which the expected recruitment from a Beverton-Holt stock-recruitment relationship is multiplied to obtain the estimate of recruitment from the stock assessment model, and represents the synthesis of age and length composition data and survey indices.

Oceanographic conditions were linked to early life history for each species based on literature-informed conceptual life history for yellowtail rockfish ([Feddern et al. 2026](#)), sablefish ([Tolimieri et al. 2018](#)), and petrale sole ([Haltuch et al. 2020](#)). For petrale sole and sablefish, ROMs based oceanographic indices for the initial publications were converted to GLORYS for the 2023 and 2025 stock assessments, respectively ([Taylor et al. 2023](#), [Wetzel et al. 2025](#)). Physical ocean conditions are often correlated due to shared atmospheric forcing, interrelated

physics driving variability, and a high degree of both spatial and temporal autocorrelation. As such, many conditions identified in the conceptual life history model are described by highly or moderately correlated time series. In order to prevent multicollinearity and overfitting of the index, correlations between each individual time series were evaluated *a priori*. Only time series with an absolute value of Pearson’s correlation coefficient of less than 0.5 (moderate correlation) were considered for inclusion in the same model (?). Up to three oceanographic conditions were included in a single model. **(Question: given the results and challenge of differentiating some of these models, this could be reduced to 0.3, that is what we did for yellowtail)**

Generalized Additive Models (GAMs) were used to fit the relationship between ecological conditions and recruitment deviations from 1994 - 2018 [GAMs implemented in ‘mgcv’; (?)]. Recruitment deviations were modeled as response variables with each possible combination of oceanographic drivers modeled as predictors. Each smoothed term was only allowed to have up to three knots ($k = 3$). As such, the relationship could represent linear, threshold, or dome-shaped relationships, but relationships were not permitted to be more “wiggly” than a parabola.

Predictive capacity was evaluated using a suite of criteria. For each model we considered the root mean square error (RMSE) from a leave-one-out cross validation (LOO), one year ahead leave-future-out RMSE for the most recent 5 years (train: 1994 - 2013; predict: 2014 - 2018) (LFO5), and one year ahead leave-future-out RMSE for the most recent 10 years (train: 1994 - 2008; predict: 2009 - 2018) (LFO10). We compared the best model based on each predictive capacity criteria

1.2 Findings

- 1) Leave-future-out cross validation as a selection criteria does a poor job at identifying the best performing model across a suite of similar models. For petrale sole, Table 1 LFO5 and LFO10 were not able to differentiate between 100 similar models (note: these models do not all have identical RMSE but it appears this way because of rounding). Petrale sole was the most extreme case of this, LFO5 was not able to differentiate between 15 models for sable fish and LFO10 was not able to differentiate between 10 models for yellowtail, only 6 models were not able to be differentiated between LFO5 and LFO10 Table 3. It is not surprising that LFO10 was able to differentiate models slightly better given it has more observations to use. For
- 2) All models, regardless of selection criteria performed relatively poorly for one-year-ahead predictions (Figure 1 A, C, E). **(Question: should the LFO5 and LFO10 models depicted here be an ensemble of the competing models? currently just the most parsimonious)** and

for sablefish and yellowtail were only able to predict whether recruitment would be above or below mean values 50% of the time petrale sole performed slightly better, but was unable to pick up interannual variability effectively, particularly magnitudes of change (Figure 1 A).

- 3) The best models that were identified in previous studies were not consistent with our results likely due to methodological differences. Yellowtail was a notable exception, but employed similar analyses compared to this study. **For sablefish**, Wetzel et al. (2025) reported CSTbjuv, and CSTpjuv as the most supported variables - none of these were included in the top 5 models, although CSTpjuv was highly ranked for marginal mean improvement of RMSE (Figure 2). Other highly supported variables included DDpre, CSTyolk, CSTegg, and CSTyolk, CSTegg were both highly ranked for marginal mean improvement of RMSE. **For petrale sole**, DDpre, MLDegg, CSTlarv, *CSTbjuv* were included in the best model in Taylor et al. (2023). MLDegg and CSTbjuv were both in the best model selected with LOO and were highly ranked based on marginal mean improvement of RMSE. **For yellowtail**, the same best model was selected with LOO compared to Feddern et al. (2026), this is likely because the methods were consistent (GAMs) compared to other analyses that focused on linear models. Every top ranking model across all there selection approaches contained oni_pjuv, which was also consistently highly ranked based on marginal mean improvement RMSE.

Table 1: Top 5 models for petrale sole using each of the three selection criteria. Leave-future-out could not differentiate between 100 competing models

variables	Hit	dev.ex	AIC	RMSE_loo	RMSE5	RMSE10	SelectionCriteria
LOO							
CSTbjuv.a DDlarv MLDegg	0.72	0.65	13.16	0.35	0.37	0.35	LOO
CSTbjuv.a DDpjuv MLDegg	0.72	0.63	14.51	0.37	0.37	0.35	LOO
CSTbjuv.a DDpjuv LSTlarv	0.72	0.72	9.53	0.37	0.37	0.35	LOO
CSTbjuv.a DDlarv LSTlarv	0.68	0.74	8.18	0.37	0.37	0.35	LOO
CSTbjuv.a CSTbjuv.b	0.76	0.40	25.30	0.39	0.37	0.35	LOO
LSTbjuv.b							
LFO5							
CSTbjuv.a	0.76	0.16	27.96	0.42	0.37	0.35	LFO5
CSTbjuv.a DDpre	0.72	0.17	29.65	0.45	0.37	0.35	LFO5
CSTbjuv.a Tpre.a	0.72	0.17	29.80	0.43	0.37	0.35	LFO5
CSTbjuv.a Tpre.b	0.72	0.23	28.26	0.44	0.37	0.35	LFO5
CSTbjuv.a DDegg1	0.72	0.26	28.23	0.48	0.37	0.35	LFO5
LFO10							
CSTbjuv.a	0.76	0.16	27.96	0.42	0.37	0.35	LFO10
CSTbjuv.a DDpre	0.72	0.17	29.65	0.45	0.37	0.35	LFO10
CSTbjuv.a Tpre.a	0.72	0.17	29.80	0.43	0.37	0.35	LFO10
CSTbjuv.a Tpre.b	0.72	0.23	28.26	0.44	0.37	0.35	LFO10
CSTbjuv.a DDegg1	0.72	0.26	28.23	0.48	0.37	0.35	LFO10

Table 2: Top 5 models for sablefish using each of the three selection criteria

variables	Hit	dev.ex	AIC	RMSE_loo	RMSE5	RMSE10	SelectionCriteria
LOO							
CSTegg LSTbjuv	0.60	0.54	80.62	1.20	1.33	1.58	LOO
CSTegg LSTbjuv MLDegg	0.60	0.56	81.54	1.20	1.21	1.24	LOO
CSTbjuv CSTegg LSTbjuv	0.60	0.61	78.14	1.25	NA	NA	LOO
CSTegg DDlarv LSTbjuv	0.60	0.54	82.38	1.26	1.46	1.32	LOO
CSTegg LSTbjuv LSTlarv	0.56	0.55	82.00	1.26	1.43	1.28	LOO
LFO5							
LSTbjuv	0.60	0.30	86.96	1.43	1.05	1.16	LFO5
LSTbjuv Tpre	0.56	0.31	88.59	1.46	1.05	1.16	LFO5
LSTbjuv Tspawn	0.56	0.37	86.58	1.54	1.05	1.16	LFO5
LSTpjuv MLDegg	0.56	0.10	93.77	1.54	1.05	1.16	LFO5
LSTegg LSTpjuv	0.44	0.09	93.93	1.56	1.05	1.16	LFO5
LFO10							
LSTpjuv	0.48	0.09	91.95	1.51	1.20	1.07	LFO10
LSTpjuv Tpre	0.44	0.09	93.85	1.63	1.20	1.07	LFO10
LSTpjuv Tspawn	0.56	0.16	92.07	1.55	1.20	1.07	LFO10
LSTlarv MLDegg	0.40	0.04	95.11	1.54	1.20	1.07	LFO10
CSTbjuv LSTegg	0.44	0.06	94.61	1.64	1.20	1.07	LFO10

Table 3: Top 5 models for yellowtail using each of the three selection criteria

variables	Hit	dev.ex	AIC	RMSE_loo	RMSE5	RMSE10	SelectionCriteria
LOO							
CutiSTIpjuv LSTpjuv oni_pjuv	0.68	0.51	18.30	0.34	0.38	0.42	LOO
CutiSTIpjuv DDegg oni_pjuv	0.60	0.46	22.41	0.38	0.38	0.42	LOO
LSTpjuv MLDpart oni_pjuv	0.56	0.34	24.88	0.38	0.32	0.35	LOO
LSTpjuv MLDlarv oni_pjuv	0.64	0.33	24.94	0.38	0.32	0.35	LOO
LSTpjuv oni_pjuv	0.60	0.28	24.89	0.38	0.32	0.35	LOO
LFO5							
oni_pjuv	0.48	0.14	26.67	0.40	0.29	0.31	LFO5
oni_pjuv Tcop	0.56	0.18	27.56	0.41	0.29	0.31	LFO5
oni_pjuv Tpart	0.48	0.15	28.42	0.42	0.29	0.31	LFO5
oni_pjuv oni_pre	0.52	0.22	27.60	0.43	0.29	0.31	LFO5
oni_pjuv pdo_larv	0.52	0.17	27.84	0.41	0.29	0.31	LFO5
LFO10							
oni_pjuv	0.48	0.14	26.67	0.40	0.29	0.31	LFO10
oni_pjuv Tcop	0.56	0.18	27.56	0.41	0.29	0.31	LFO10
oni_pjuv Tpart	0.48	0.15	28.42	0.42	0.29	0.31	LFO10
oni_pjuv oni_pre	0.52	0.22	27.60	0.43	0.29	0.31	LFO10
oni_pjuv pdo_larv	0.52	0.17	27.84	0.41	0.29	0.31	LFO10

Table 4: Top 5 models for hake using each of the three selection criteria

variables	Hit	dev.ex	AIC	RMSE_loo	RMSE5	RMSE10	SelectionCriteria
LOO							
CSTeggs CSTpjuvs MLDlate	0.68	0.48	97.23	1.72	2.04	1.84	LOO
CSTeggs CSTpjuvs DDspawn	0.68	0.56	94.99	1.72	2.04	1.84	LOO
CSTeggs CSTpjuvs MLDlarv	0.64	0.47	97.90	1.73	2.04	1.84	LOO
CSTeggs CSTpjuvs MLDyolk	0.68	0.46	98.36	1.73	2.04	1.84	LOO
CSTeggs CSTpjuvs Tspawn	0.68	0.54	96.02	1.77	2.04	1.84	LOO
LFO5							
PUTpjuv	0.44	0.04	104.89	1.87	1.96	1.83	LFO5
PUTpjuv SSHjas.pre	0.56	0.10	105.28	1.88	1.96	1.83	LFO5
PUTpjuv SSHmsn.pre	0.40	0.06	106.37	1.94	1.96	1.83	LFO5
PUTpjuv SSHmsc	0.40	0.10	106.18	2.14	1.96	1.83	LFO5
PUTpjuv SSHmss	0.36	0.11	106.09	2.17	1.96	1.83	LFO5
LFO10							
CSTlaten	0.56	0.04	104.97	1.87	2.04	1.76	LFO10
CSTlaten MLDyolk	0.52	0.04	106.90	1.89	2.04	1.76	LFO10
CSTlaten SSHjas.pre	0.44	0.07	106.14	1.91	2.04	1.76	LFO10
CSTlaten SSHmsn.pre	0.52	0.05	106.57	1.94	2.04	1.76	LFO10
CSTlaten PUTpjuv	0.44	0.06	106.33	1.89	2.04	1.76	LFO10

One reason for why these selection criteria produce such different results is that these models are highly sensitive to the training versus prediction period. There are three, interlarded, factors that can contribute to this 1) the relative importance of an individual on recruitment varies through time compared to other variables based on a myriad of interrelated factors and conditions, 2) the time period over which a model is trained may result in differences in the functional relationship, 3) the contrast of the environmental data during the training period relative to the prediction period likely contributes to both 1 and 2. We examined how each of these things contributed to the differences in variable importance and model selection discussed above.

2 How does the relative importance of individual oceanographic conditions vary through time?

2.1 Methods

To do this, used a rolling window approach following the methods described above. Specifically, we fit all model combinations to 15-year training periods and calculated the marginal mean improvement of each variable for each period. As such, we fit a model from starting year 1994 - 2008, then 1995 - 2009 and so on for the entire 1994 - 2018 time frame.

2.2 Findings

All species had variables that were inconsistent through time. This is most dramatic for Petrale sole and CSTegg2 which had >10% marginal mean improvement during the 1997 - 2011 time period, but declined dramatically during time periods that included 2012 and beyond. This same time frame showed a change in relative importance for a number of variables for petrale sole, including DDegg and LST egg declining in importance, while DDbjuv increased in importance. Sablefish and yellowtail also experienced a switching of variable importance that. For example, CSTbjuv not very important (~2.5%) in time periods from 1994 - 2014 but starting in the 2001 - 2015 time period, the importance increased to 7.5%. The importance of Tpre also increased during this time period, while CSTpjuv and CSTyolk decreased. Yellowtail had more consistent variable importance for some variables, specifically ONI being consistently important while DD variables were consistently unimportant, but the variable importance of yellowtail also declined starting in the 2001 - 2015 window.

Question: This could be a factor or heatwave conditions? It will be interesting to see what comes with the hake data...currently the




Figure 1: One year ahead predictions of recruitment deviations (black line) for the last ten years (A., B., C.) and and leave-one-out predictions for all years (B., D., F.) for the best supported models based on RMSE for one year ahead predictions for the last 10 years (LFO10, blue), last 5 years (LFO5, yellow) and RMSE for leave-one-out predictions (red, LOO). Rows depict individual species, Petrale Sole (A.,B.), Sablefish (C. D.) and Yellowtail reockfish (E. F.)




Figure 2: Marginal mean improvement of RMSE for each oceanographic driver and each selection criteria (leave-future-out last 10 years, LFO10; leave-future-out last 5 years, LFO5; leave-one-out all years)

evidence is there but also seems weak and another species will be interesting! I will try to get the hake stuff into this before we meet




Figure 3: Marginal mean improvement of each oceanographic driver using 15-year time windows for each of the three groundfish species of interest

3 Are these differences driven by changes in the functional relationships?

3.1 Methods

To evaluate the stability of the functional relationship between oceanographic conditions and recruitment, we did evaluated the retrospective predictive capacity of the top models selected by LOO (note: I also did this with the best LFO models but not super meaningful since there are so many competing models) and calculated mohn's rho (?) of the estimated smoothed function using a 15-year peel. At its core, this tells us how much the parameter estimate of the oceanographic condition deviates from a linear response by peeling off the last 15 years of data. We repeated this analysis using single predictor models to evaluate the stability of each individual driver for each species.

3.2 Findings

Yellowtail had the most stable functional relationships of any species, with relatively small Mohn's rho estimated for most variables (Figure 5) and consistent functional relationships for most of the 15-year peel (Figure 4) for the best supported model based on LOO. In contrast, the stability of the functional relationship for sablefish began to decline with terminal year 2010, and declined substantially by terminal year 2003 (Figure 4). Petrale sole experienced instability in MLDegg with terminal year 2015, which increased for both MLDegg and DDlarv as years were peeled back (Figure 4).




Figure 4: Mohn's rho (right) and associated functional relationships(left) for each of the 15 "peeled" years for each species. Mohn's rho can be interpreted as deviations from 0 indicate functional relationships that differ from the model fit to the full 1994 - 2018 time period

4 Are these differences driven by contrast in the environmental data?

Question: how might we show this? I am wondering about good ways to show contrast, which shouldn't be hard given the datasets are already standardized. Nick - I remember you might have had an idea on this when we were




Figure 5: Absolute value of Mohn's rho for each oceanographic predictor




Figure 6: Mean Mohns rho by terminal year for single predictor models showing depicting mean values across oceanographic predictors by species with a loess smooth by species

chatting some time but it has escaped my mind! I will try and draft something between now and when we meet and maybe we can workshop it a bit...

5 Investigating the 2012 break point as a potential driver of stability/observed changes

5.1 Motivation

Based on the findings in 1 - 3 there appears to be 1) a decline in model stability that is delineated by a 2012 - 2015 breakepoint Figure 6 and 2) a change in the relative importance of different drivers Figure 3 with a breakpoint for most species in the early 2010s.

5.2 Methods

To dig into this a bit, I divided oceanographic conditions into two groups 1) stable variables that have a Mohn's rho of 0 and 2) unstable variables that have a Mohn's rho of 0.1 (NOTE: I played around with how I divided these quite a bit and the results were not sensitive to it).

I fit a bayesian DFA to the two groups of oceanographic conditions to see if there were differences in the underlying patterns of variability. I tested up to three trends and for both groups of variables the best supported models only had 1.

5.3 Findings

We do not observe differences between the two groups of variables and there are strong loadings across most time series across a similar latent trend. One very notable observation about this trend is that for most of the 1994 - 2018 time period the latent trend is at or below the long-term mean. However, two notable exceptions exist 1) 1997/1998 and 2) 2014 onward. 1997/1998 was the second largest El Nino event on record that had huge consequences for dynamic height, water mass characteristics, and a temporary reversal of net alongshore transport ((?)) which also had substantial impact on the ecosystem ((?)). The 2014 onward period doesn't need much explaining (marine heatwaves, pan basin warming).

6 Preliminary Conclusion

Collectively, it seems like the inclusion(exclusion) of the 2010 - 2018 time period in the training of the model greatly impacts variable importance and model




Figure 7: Bayesian dynamic factor analysis for “stable” and “unstable” variables

stability for recruitment (Figure 6). This in turn, results in inconsistent model selection across selection criteria, as they weigh the 2009 - 2018 time period differently across criteria. We also find that the stability of these models and the differences in prediction capacity does vary by species, with yellowtail showing more stability and consistent variable importance and also greater consistency in model selection across the suite of selection criteria we used.

Question: to dig into this a little deeper we could fit the models from 1994 - 2010 and see if those models are more stable? and generate the same plots of Figure 6 and Figure 5

- Fedder, A., Megan L. and Darby, N. Tolimieri, and K. L. Oken. 2026. Stage-specific indicators of northern yellowtail rockfish (*Sebastes flavidus*) recruitment in the California Current Ecosystem. Fisheries Oceanography.
- Haltuch, M. A., N. Tolimieri, Q. Lee, and M. G. Jacox. 2020. [Oceanographic drivers of petrale sole recruitment in the California Current Ecosystem](#). Fisheries Oceanography 29:122–136.
- Johnson, K. F., C. R. Wetzel, and N. Tolimieri. 2023. Status of sablefish (*Anoplopoma fimbria*) along the U.S. West Coast in 2023. Pacific Fishery Management Council, Portland, Oregon. 145 p.
- Oken, K. L., I. G. Taylor, M. L. Fedder, A. D. Whitman, and F. P. Caltabellotta. 2025. Status of the yellowtail rockfish stock off the US West Coast north of 40°10' in 2025. Pacific Fishery Management Council, Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.
- Taylor, I., V. Gertseva, and N. Tolimieri. 2023. Status of petrel sole (*Eopsetta jordan*) along the US West Coast in 2013. Pacific Fishery Management Council, Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.
- Tolimieri, N., M. Haltuch, Q. Lee, M. Jacox, and S. J. Bograd. 2018. Oceanographic drivers of sablefish recruitment in the California current. Fisheries Oceanography 27:458–474.
- Wetzel, C. R., A. M. Berger, B. C. Z. J. A, T. N, W. E. J, and H. M. 2025. Status of sablefish (*Anoplopoma fimbria*) off the U.S. West Coast in 2025. Pacific Fisheries Management Council, Portland, Oregon.