

Qualify Exam 2015

Mingwei Fei

January 19, 2023

1 Problem 1- MVN and Chi-square

Suppose that $Y \sim N(\mu, \Sigma)$ where Σ is symmetric and full rank, Let A be a symmetric matrix.

1.1 a. Quadratic Form $Y^T A Y$ and Chi-square distribution

Show that the quadratic form $Y^T A Y$ can be represented as

$$Y^T A Y = \sum_{i=1}^k \lambda_i W_i$$

where the W_i 's are independently distributed as noncentral chi-squared variables with d_i degrees of freedom and noncentrality parameter δ_i , that is, $W_i \sim \chi_{d_i}^2(\delta_i), i = 1, 2, \dots, k$. Indicate what λ_i, d_i, δ_i are equal to.

1.1.1 Question

(i) *Quadratic form of normal random variable with a specific variance:*

Assume Y is a normal random variable with mean 0 and a specific variance is

$$\begin{aligned}\Sigma &= [\Sigma^{1/2}]^T [\Sigma^{1/2}], && \text{Cholesky decomposition} \\ Y &= [\Sigma^{1/2}] Z, && Z \sim N(0, 1) \\ Y_i^T Y_i &= \left([\Sigma^{1/2}] Z\right)^T [\Sigma^{1/2}] Z = Z^T \Sigma Z = \sigma^2 \chi^2\end{aligned}$$

(ii) non-centrality chi-square distribution

And if assume Y the mean is not 0, $Y \sim N(\mu, 1)$, then we have

$$\begin{aligned}Y_i^T Y_i &= \sigma^2 \chi^2(\delta) \\ \delta &= \mu_i^2\end{aligned}$$

The mean (mu) of the chi-square distribution is its degrees of freedom, k . Because the chi-square distribution is right-skewed, the mean is greater than the median and mode. The variance of the chi-square distribution is $2k$.

Suppose $Y_{n \times n}$, and Σ is full rank. so Σ is $n \times n$ dimension matrix,

We can transform Y_i into $N(\mu, 1)$ distribution, so that the quadratic form will be a non-central chi-square distribution.

If Z_1, \dots, Z_k are independent, standard normal random variables, then the sum of their squares is chi-square distribution,

$$Q = Z_i^2 \sim \chi^2(k)$$

$$p(k) = \frac{1}{2^{k/2} \Gamma(k/2)} x^{k/2-1} \exp(-\frac{x}{2})$$

Non-central Chi-square:

Here the k is unknown, we need to show that the sum of non-central chi-square distribution is also a non-central chi-square distribution with distribution transformation. The distribution transformation generally use Moment Generating Function.

Lemma:

Let $Q_i \sim \chi_{k_i}^2(\lambda_i)$ for $i = 1, \dots, n$, be independent. Then, $Q = \sum_{i=1}^n Q_i$ is a noncentral $\chi_k^2(\lambda)$, where $k = \sum_{i=1}^n k_i$ and $\lambda = \sum_{i=1}^n \lambda_i$.

Chi-square distribution and non-central chi-square distribution are totally different. I need to understand the components δ_i and d_i in the non-central chi-square distribution.

The non-central chi-square distribution: Let $(X_1, X_2, \dots, X_i, \dots, X_k)$ be k independent, normally distributed random variables with means μ_i and unit variances. Then the random variable

$$Q = \sum_{i=1}^k X_i^2 \sim \chi^2(k, \lambda), \quad \lambda = \sum_{i=1}^k \mu_i^2$$

(iii) Theorem using spectral decomposition

For any $k \times k$ symmetric matrix A the quadratic form defined by A can be written using its spectral decomposition as

$$Q(x) = x'Ax = \sum_{i=1}^k \lambda_i \|q_i^T x\|^2$$

where the eigendecomposition of A is $Q'\Lambda Q$ with Λ diagonal with diagonal elements $\lambda_i, i = 1, \dots, k$, Q is the orthogonal matrix with the eigenvectors, $q_i, i = 1, \dots, k$ as its columns.

The proof: For any $x \in R^k$ let $y = Q^T x = Q^{-1}x$ Then

$$\begin{aligned} Q(x) &= \text{tr}(x'Ax) = \text{tr}(x'Q'\Lambda Qx) = \text{tr}(\Lambda Qx(Qx)^T) \\ &= \text{tr}(\Lambda y'y) = \sum_{i=1}^k y_i^2 = \sum_{i=1}^k \lambda_i \|q_i^T x\|^2 \end{aligned}$$

This proof uses a common trick of expressing the scalar $Q(x)$ as the trace of a 1×1 matrix so we can reverse the order of matrix multiplications.

So in the general case of linear combination of non-central chi-square. We will need to take care of the Σ , assume $Z \sim N(\mu, I)$

$$\begin{aligned} Z &= \Sigma^{-1/2}Y \\ Q(Y) &= Y^T A Y = (\Sigma^{1/2}Z)^T A (\Sigma^{1/2}Z) = Z^T (\Sigma^{1/2})' A (\Sigma^{1/2}) Z \\ &= \sum_{i=1}^k \lambda_i \|q_i^T Z\|^2 \\ W_i &= \|q_i^T Z\|^2 \\ &\sim \chi^2(d_i, \delta_i) \end{aligned}$$

$\Sigma^{1/2}$ is symmetric as well, so $\Sigma^{1/2}A\Sigma^{1/2}$ is also symmetric. Symmetric matrix could be spectral decomposed.

I need to understand the product of orthogonal matrix and other matrix, orthogonal matrix is idempotent, which the eigenvalues are either 0 or 1.

(iv) The quadratic form question often occurs in the multivariate normal distribution.

Quadratic form theorem 1: If $y \sim N(\mu_y, \Sigma_y)$, then

$$z = AY = N(\mu_z = A\mu_y, \Sigma_z = A\Sigma_y A')$$

where A is a matrix of constants.

Proof:

$$\begin{aligned} E[z] &= E[AY] = AE[Y] = A\mu_y \\ \text{Var}[z] &= E[(z - E[z])(z - E[z])'] = E[(AY - A\mu_y)(AY - A\mu_y)'] \\ &= E[A(Y - \mu_y)(Y - \mu_y)'A'] = AE[(Y - \mu_y)(Y - \mu_y)']A' \\ &= A\Sigma_y A' \end{aligned}$$

Quadratic form theorem 2: quadratic form of n standard normal is chi-square with n degrees of freedom. Here we need to pay attention to the n degrees of freedom of chi-square distribution.

Quadratic form theorem 3: If $y \sim N(0, \sigma^2 I)$ and M is a symmetric idempotent matrix of rank m then

$$\frac{y'My}{\sigma^2} = \chi^2(trM)$$

Proof: Since M is symmetric it can be diagonalized with an orthogonal matrix Q .

$$Q'MQ = \Lambda = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \dots & \dots & \lambda_n \end{pmatrix}$$

Furthermore, since M is idempotent all these roots are either zero or 1. Thus we can choose Q so that Λ look like

$$Q'MQ = \Lambda = \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix}$$

The dimension of the identity matrix will be equal to the rank of M , since the number of non-zero roots is the rank of the matrix. Since the sum of the roots is equal to the trace, the dimension is also equal to the trace of M . Now let $v = Q'y$, compute the moments of

$$\begin{aligned} v &= Q'y \\ E[v] &= Q'E[y] = 0 \\ Var[v] &= Q'Var[y]Q = Q'\sigma^2 I Q = \sigma^2 Q'Q = \sigma^2 I, \quad Q \text{ is orthogonal} \\ v &\sim N(0, \sigma^2 I) \end{aligned}$$

Now consider the distribution of $y'My$ using the transformation v . Since Q is orthogonal, its inverse is equal to its transpose. This means that $y = (Q')^{-1}v = Qv$. Now write the quadratic form as follows

$$\begin{aligned} \frac{y'My}{\sigma^2} &= \frac{v'Q'MQv}{\sigma^2} \\ &= \frac{1}{\sigma^2} v' \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} v \\ &= \frac{1}{\sigma^2} \sum_{i=1}^{trM} v_i^2 \\ &= \sum_{i=1}^{trM} \frac{v_i^2}{\sigma^2} \end{aligned}$$

This is the sum of squares of $(tr M)$ standard normal variables and so is a χ^2 variable with trM degrees of freedom.

1.1.2 Proof

$$Y^T AY = \sum_{i=1}^k \lambda_i W_i$$

where W_i are independently distributed as noncentral chi-squared variables with d_i degrees of freedom and noncentrality parameter δ_i , that is, $W_i \sim \chi_{d_i}^2(\delta_i)$, $i = 1, 2, \dots, k$. Indicate what λ_i, d_i, δ_i are equal to.

$$\begin{aligned} \Sigma &= \Sigma^{1/2} \Sigma^{1/2^T}, & \text{by semi-definite matrix} \\ \Sigma^{1/2} &= (\Sigma^{1/2})^T \\ Y^T AY &= Y^T \Sigma^{-1/2} \Sigma^{1/2} A \Sigma^{1/2} \Sigma^{-1/2} Y \\ Z &= \Sigma^{-1/2} Y \sim N(\Sigma^{-1/2} \mu, \Sigma^{-1/2} \Sigma \Sigma^{-1/2}) = N(\Sigma^{-1/2} \mu, I) \\ \Sigma^{1/2} A \Sigma^{1/2} &= P \Lambda P', & \text{semi-definition matrix, symmetric, spectral decomposition} \end{aligned}$$

By now, we have constructed quadratic form into $Z^T B Z$, which we can recognize that we will use the spectral decomposition

$$\begin{aligned} Z &= \Sigma^{-1/2} Y \\ B &= \Sigma^{1/2} A \Sigma^{1/2} = P \Lambda P', & P \text{ is orthogonal matrix} \\ Z^T B Z &= \text{tr}(Z^T B Z) = \text{tr}(Z^T P \Lambda P' Z) = \text{tr}(\Lambda [P' Z][P' Z]^T) \end{aligned}$$

We can see that Λ is the matrix of eigenvalues of $B = \Sigma^{1/2} A \Sigma^{1/2} = A \Sigma$ which we can prove later.

The trace of $[P' Z][P' Z]^T$ is actually $W_i = \|P'_i Z\|^2$, and we can see that it is a non-centrality chi-square distribution with noncentrality parameter δ_i . The degrees of freedom of chi-square distribution is the number of sum of chi-squares. We can see that, in multivariate normal distribution setting, there is only one variable Z , so the degrees of freedom for W_i is 1.

$$\begin{aligned} P'_i Z &= N(P_i^T \Sigma^{-1/2} \mu, P_i^T I P_i) = N(P_i^T \Sigma^{-1/2} \mu, I), & P_i \text{ is orthogonal vector} \\ \|P'_i Z\|^2 &\sim \chi^2(1, \delta_i), & \delta_i = \frac{[P_i^T \Sigma^{-1/2} \mu]^2}{2} \end{aligned}$$

Need to be familiar with non-centrality chi-square, the non-central parameter is half of the mean square.

To prove eigenvalues of $\lambda_1, \lambda_2, \dots, \lambda_n$ of $\Sigma^{1/2} A \Sigma^{1/2}$ are also eigenvalues of $A \Sigma$.

$$\begin{aligned} \left| \Sigma^{1/2} A \Sigma^{1/2} - \lambda I \right| &= 0 \\ \rightarrow \left| \Sigma^{-1/2} \Sigma^{1/2} A \Sigma^{1/2} \Sigma^{1/2} - \Sigma^{-1/2} \lambda I \Sigma^{1/2} \right| &= 0 \\ \rightarrow \left| A \Sigma - \lambda I \right| &= 0 \end{aligned}$$

1.2 b. MGF of $Y^T AY$

Use part (a) to derive the moment generating function of $Y^T AY$. Let $m(t)$ denote the moment generating function. Show that $m(t)$ exists in a small neighborhood of $t = 0$, say, $|t| < t_0$ for some positive constant t_0 . Find the maximal value of t_0 i.i.d chi-square distribution sum MGF.

1.2.1 Preparation

In part (a), we already derive that $Y^T AY = \sum \lambda_i W_i$, while W_i is a one degree of freedom of chi-square distribution with non-centrality parameter. So the MGF could be directly used for the sum of chi-square distribution.

I may need to remember the standard form of MGF of chi-square, or I need to derive that from scratch.

first, get the MGF for non-centrality chi-square, then use the transformation to put λ_i .

Another thing needs to pay attention is that, Y is multivariate normal, not univariate normal. The MGF derivation is different from univariate normal variable.

The MGF for univariate normal:

$$\begin{aligned}
 M(t) &= \prod_{i=1}^k M_i(t) \\
 p(x_i) &= \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(X - \mu_i)^2}{2}\right) \\
 M_i(t) &= E[\exp(x_i^2 t)] = \frac{1}{\sqrt{2\pi}} \int \exp\left[-\frac{(1-2t)x^2 - 2\mu x + \mu_i^2}{2}\right] dx \\
 &= \frac{1}{\sqrt{2\pi}} \int \exp\left[-\frac{x^2 - 2\mu_i/(1-2t)x + \mu_i^2/(1-2t)^2 - \mu_i^2/(1-2t)^2 + \mu_i^2/(1-2t)}{2((1-2t)^{-1})}\right] dx \\
 &= \exp\left[\frac{\mu_i^2/(1-2t)^2 - \mu_i^2/(1-2t)}{2((1-2t)^{-1})}\right] (1-2t)^{-1/2} \int \frac{1}{\sqrt{2\pi(1-2t)^{-1}}} \exp\left[-\frac{[x - \mu_i/(1-2t)]^2}{2((1-2t)^{-1})}\right] dx \\
 &= (1-2t)^{-1/2} \exp\left[\frac{\mu_i^2 t}{(1-2t)}\right] \\
 M(t) &= \prod_{i=1}^k (1-2t)^{-1/2} \frac{\mu_i^2 t}{(1-2t)} = (1-2t)^{-k/2} \exp\left[\frac{\sum_i \mu_i^2 t}{(1-2t)}\right]
 \end{aligned}$$

In which, $(1-2t) > 0$, $t < 1/2$. We can see that the product of non-centrality chi-square distributions is also a non-central chi-square distribution.

$$M_Z(t) = E[\exp(t^T Z)] = e^{\frac{1}{2} t^T t}$$

while for the transformation

$$P'_i Z = N(P_i^T \Sigma^{-1/2} \mu, I)$$

$$\|P'_i Z\|^2 \sim \chi^2(1, \delta_i), \quad \delta_i = \frac{[P_i^T \Sigma^{-1/2} \mu]^2}{2}$$

Then the MGF

$$X_i = Z + P_i^T \Sigma^{-1/2} \mu$$

$$M_{X_i}(t) = E[\exp(t^T X_i)] = \exp(t^T [P_i^T \Sigma^{-1/2} \mu]) e^{\frac{1}{2} t^T t}$$

1.2.2 Proof

However, in part (a), we broke down the vector into sum of each univariate chi-square distribution. In that sense, it is still a univariate chi-square MGF question.

Another method is to let $Z \sim N(0, 1)$, then $(Z + \mu)^2$ has a noncentral chi-square distribution with one degree of freedom, the MGF of $(Z + \mu)^2$

$$E[\exp(t(Z + \mu)^2)] = \frac{1}{\sqrt{2\pi}} \int \exp(t(Z + \mu)^2) \exp(-\frac{Z^2}{2}) dZ$$

$$= \frac{1}{\sqrt{2\pi}} \int \exp[-\frac{(1 - 2t)Z^2 - 2\mu Z + \mu^2}{2}] dZ$$

$$= (1 - 2t)^{-1/2} \exp[\frac{\mu^2 t}{(1 - 2t)}]$$

$$M_{W_i}(t) = (1 - 2t)^{-1/2} \exp[\frac{2\delta_i t}{(1 - 2t)}], \quad \delta_i = \frac{1}{2} \mu_i^2$$

We also note that, chi-square with 1 degree of freedom is a gamma distribution $\chi^2(1, 0) = \text{Gamma}(1/2, 2)$.

As $Y^T A Y = \sum_{i=1}^n \lambda_i W_i$, then

$$M(t) = E\left[\exp\left(\left[\sum_{i=1}^n \lambda_i W_i\right]t\right)\right] = \prod_{i=1}^k M_i(t)$$

$$= \prod_i^k (1 - 2\lambda_i t)^{-1/2} \exp[\frac{2\delta_i \lambda_i t}{(1 - 2\lambda_i t)}]$$

As we see that $1 - 2\lambda_i t > 0$, then we have $t < \frac{1}{2\lambda_i}$.

By definition, a non-central chi-square random variable $\chi_{n,\lambda}^2$ with n df and parameters

$\lambda = \sum_i^n \mu_i^2$ is the sum of n **independent** normal variables $X_i = Z_i + \mu_i, i = 1, 2, \dots, n$.

$$\begin{aligned}\chi_{n,\lambda}^2 &= \sum_i^n X_i^2 = \sum_i^n (Z_i + \mu_i)^2 \\ M(t) &= \prod_{i=1}^k M_i(t) = \prod_i^k (1-2t)^{-1/2} \exp\left[\frac{\mu_i^2 t}{(1-2t)}\right] \\ &= (1-2t)^{-k/2} \exp\left[\frac{\sum_i^n \mu_i^2 t}{(1-2t)}\right] = (1-2t)^{-n/2} \exp\left[\frac{\lambda t}{(1-2t)}\right]\end{aligned}$$

1.3 c. Trace and Rank

Use part (a) to show that $\text{tr}[(A\Sigma)^2] = \text{tr}(A\Sigma) = r$, where r is the rank of A , then $Y^T A Y$ has a chi-squared distribution. Determine its degrees of freedom and noncentrality parameter.

1.3.1 Preparation

From part (a), we see that $Y^T A Y$ is sum of chi-square distributions. In order to show it is a chi-square, then the $\lambda_i = 1, 0$. Well, $\text{tr}[(A\Sigma)^2] = \text{tr}(A\Sigma) = r$ is similar to the idempotent matrix definition, and λ_i is actually the eigenvalue of $A\Sigma$.

But how can we show the idempotency of $A\Sigma$? Rank is the sum of eigenvalues, the link is how to go from the rank to show that the eigenvalues are exactly 0 or 1.

Because Σ is full rank, so $R(A\Sigma) = R(A)$ as the column space won't change when it is the right product of a full rank matrix. So the number of non-zero eigenvalues is r .

$$\begin{aligned}\sum_{i=1}^k \lambda_i &= \sum_{i=1}^k I(\lambda_i \neq 0) \\ I(\lambda_i \neq 0) &= \begin{cases} 1, & \lambda_i \neq 0 \\ 0, & \lambda_i = 0 \end{cases}\end{aligned}$$

So we can see that $\lambda_i = I(\lambda_i \neq 0) = 1$ when $\lambda_i \neq 0$.

Use the spectral decomposition of $A\Sigma = P\Lambda P'$.

$$\begin{aligned}
A\Sigma &= P\Lambda P' \\
(A\Sigma)^2 &= P\Lambda^2 P^T \\
tr((A\Sigma)^2) &= tr(A\Sigma) \\
&\rightarrow tr(P\Lambda P') = tr(P\Lambda^2 P^T) \\
&\rightarrow tr(\Lambda^2) = tr(\Lambda), \\
&\rightarrow \sum_{i=1}^k \lambda_i^2 = \sum_{i=1}^k \lambda_i \\
&\rightarrow \sum_{i=1}^k \lambda_i(\lambda_i - 1) = 0 \\
&\rightarrow \lambda_i = 0, 1
\end{aligned}$$

1.3.2 Proof

From part (a) that

$$\begin{aligned}
tr(A\Sigma) &= tr(\Sigma^{1/2} A \Sigma^{1/2}) \\
tr\left([\Sigma^{1/2} A \Sigma^{1/2}]^2\right) &= tr\left([\Sigma^{1/2} A \Sigma^{1/2}][\Sigma^{1/2} A \Sigma^{1/2}]\right) \\
&= tr\left([\Sigma^{1/2} A \Sigma A \Sigma^{1/2}]\right) \\
&= tr\left([A \Sigma A \Sigma^{1/2} \Sigma^{1/2}]\right) \\
&= tr\left([A \Sigma A \Sigma]\right) \\
&= tr\left([(A\Sigma)^2]\right)
\end{aligned}$$

If $\lambda_i = 0, 1$, then $Y^T A Y$ is the sum of r chi-square $\chi^2(1, \delta_i)$, which is also a chi-square distribution.

$$Y^T A Y = \sum_{i=1}^r W_i \sim \chi^2(r, \delta), \quad \delta = 2 \sum_{i=1}^r \mu_i^2$$

1.4 d. Distribution

Show that $Y^T A Y$ has a noncentral chi-squared distribution if and only if $A\Sigma$ is idempotent.

1.4.1 Preparation

This problem needs to prove "if and only if", that From part(c), we have shown that if $\lambda_i = 0, 1$ which is idempotent will lead to $Y^T AY$ as a noncentral chi-squared distribution.

While if $Y^T AY$ is a noncentral chi-squared distribution, then $\lambda_i = 0, 1$ then it is a idempotent.

Idempotent is $A^2 = A, A^T = A$. So if $A\Sigma = (A\Sigma)^2$, $A\Sigma$ is idempotent.

we have the MGF of linear combination of non-central chi-square distribution Y

$$\begin{aligned} M(t) &= \prod_{i=1}^n M_{W_i}(\lambda_i t) \\ &= \prod_{i=1}^n (1 - 2\lambda_i t)^{-1/2} \exp\left(\frac{2\delta_i \lambda_i t}{1 - 2\lambda_i t}\right) \end{aligned}$$

Then we can see that the shape parameter is $\frac{1}{2\lambda_i}$. If we want to have a non-central chi-square distribution for Y, then all λ_i need to be the same, 1. Otherwise it won't be a chi-square.

1.4.2 Proof

$$\begin{aligned} M_Y(t) &= \prod_{i=1}^n (1 - 2\lambda_i t)^{-1/2} \exp\left(\frac{2\delta_i \lambda_i t}{1 - 2\lambda_i t}\right) \\ &= (1 - 2t)^{-n/2} \exp\left(\frac{\sum_{i=1}^n 2\delta_i t}{1 - 2t}\right) \end{aligned}$$

And for chi-square distribution, the shape parameter has to be 1/2, so the $w_i = 1$. So we prove that if Y is a non-central chi-square distribution, A has to have the eigenvalues either 1 or 0.

When the eigenvalues are only 0, 1, then $A\Sigma$ is a projection matrix, then it is idempotent.

2 Problem 2 - Linear Model

Consider the linear model $Y = X\beta + \epsilon$, where $\epsilon \sim N_n(0, \sigma^2 I)$, X is $n \times p$ of rank p, β is $p \times 1$, and (β, σ^2) are unknown. Define $H = X(X^T X)^{-1} X^T$ and let h_{ii} denote the ith diagonal element of H. Further let $\hat{\sigma}^2$ denote the usual unbiased estimator of σ^2 for the linear regression model and let $\hat{\epsilon}_i$ denote the ordinary residual. Let

$$\begin{aligned} A_i &= \frac{\hat{\epsilon}_i^2}{\sigma^2(1 - h_{ii})} \\ B_i &= \frac{(n - p)\hat{\sigma}^2}{\sigma^2} - \frac{\hat{\epsilon}_i^2}{\sigma^2(1 - h_{ii})} \end{aligned}$$

2.1 a. Show that $B \sim \chi_{n-p-1}^2$

2.1.1 Preparation

In order to prove $B \sim \chi_{n-p-1}^2$, we need to show that it is a sum of n-p-1 normal distribution square (by definition of chi-square distribution).

We can write $\sigma^2, \hat{\sigma}^2$ the quadratic form in the sum of squares of normal distribution. But the fraction between two quadratic forms is what distribution?

Note that the division is not F-distribution, as it didn't divided by degrees of freedom.

$\hat{\sigma}^2$ is the usual unbiased estimator of σ^2 , we need to know that if unbiased, then the degrees of freedom is n-p.

Another thing need to pay attention that, σ^2 is not a distribution, it is fixed and it is the variance of Y_i just unknown.

Variance of Y_i is also the variance of ϵ , but we need to know the difference and how to use them.

$$\begin{aligned} \frac{(n-p)\hat{\sigma}^2}{\sigma^2} &= \frac{Y^T(I-H)Y}{\sigma^2} = \frac{[(I-H)Y]^T[(I-H)Y]}{\sigma^2} \\ &= \sum_{i=1}^n \frac{(y_i - \hat{y}_i)^2}{Var(y_i)} \end{aligned}$$

So is n-p sum of normal distribution square. Then try to link the second term

$$\frac{\hat{\epsilon}_i^2}{\sigma^2(1-h_{ii})} = \frac{\hat{\epsilon}_i^2}{V(e_i)}$$

This question requires understanding the matrix H, or the relationship between o.p.o matrix and scalar form. And the characteristic of H matrix itself $H^2 = H, H^T = H$.

If we write that in scalar form,

$$(h_{i1}, h_{i2} \dots h_{in}) \begin{pmatrix} h_{i1} \\ h_{i2} \\ \dots \\ h_{in} \end{pmatrix} = h_{i1}^2 + h_{i2}^2 + \dots + h_{in}^2 = h_{ii}$$

Variance, Covariance of e

$$\begin{aligned} V(e) &= E[(e - E(e_i))(e - E(e_i))^T] = (I-H)E(\epsilon\epsilon^T)(I-H)^T \\ &= (I-H)I\sigma^2(I-H)^T \\ &= (I-H)(I-H)^T I\sigma^2 = (I-H)\sigma^2 \end{aligned}$$

Need to know that $V(e_i)$ is given by the ith diagonal element $1 - h_{ii}$ and $Cov(e_i, e_j)$ is given by the (i, j)th element of $-h_{ij}$ of the matrix $(I-H)\sigma^2$.

We also need to know that both Y_i and ϵ_i follows the same distribution

$$\begin{aligned} e - E(e) &= (I - H)Y = (Y - X\beta) = (I - H)\epsilon \\ E(\epsilon\epsilon^T) &= V(\epsilon) = I\sigma^2, \quad E(\epsilon) = 0 \end{aligned}$$

We also have correlation

$$\begin{aligned} \rho_{ij} &= \frac{Cov(e_i, e_j)}{\sqrt{V(e_i)V(e_j)}} = -\frac{h_{ij}}{(1 - h_{ii})(1 - h_{jj})} \\ SS(b) &= SS(parameter) = b^T X^T Y = \hat{Y}^T Y = Y^T H^T Y = Y^T H Y = Y^T H^2 Y = \hat{Y}^T \hat{Y} \end{aligned}$$

The average $V(\hat{Y}_i)$ to all data points is

$$\begin{aligned} \sum_{i=1}^n \frac{V(\hat{Y}_i)}{n} &= \frac{trace(H\sigma^2)}{n} = \frac{p\sigma^2}{n} \\ \hat{Y}_i &= h_{ii}Y_i + \sum_{j \neq i} h_{ij}Y_j \end{aligned}$$

Studentized residual

$$V(e_i) = (1 - h_{ii})\sigma^2$$

where σ^2 is estimated by s^2

$$s^2 = \frac{e^T e}{n - p} = \frac{\sum_{i=1}^n e_i^2}{n - p}$$

Sum of Squares attributable to e_i

$$\begin{aligned} e &= (I - H)Y \\ e_i &= -h_{i1}Y_1 - h_{i2}Y_2 - \dots + (1 - h_{ii})Y_i - \dots - h_{in}Y_n = c^T Y \\ Var(e_i) &= c^T Var(Y) c = \sigma^2 c^T c \\ c^T &= (-h_{i1}, -h_{i2}, \dots, (1 - h_{ii}), \dots, -h_{in}), \quad (I - H)I - H = I - H \\ c^T c &= \sum_{j=1}^n h_{ij}^2 + (1 - 2h_{ii}) = (1 - h_{ii}) \\ Var(e_i) &= \sigma^2(1 - h_{ii}) \\ SS(e_i) &= e_i^2 = \left((I - H)y \right)^T (I - H)y = y^T (I - H)y \end{aligned}$$

Need to learn to write in matrix form.

2.1.2 Proof

Write B_i in matrix form

$$\frac{(n-p)\hat{\sigma}^2}{\sigma^2} = \frac{Y^T(I-H)Y}{\sigma^2} = \frac{[(I-H)Y]^T[(I-H)Y]}{\sigma^2}$$

$$\frac{\hat{\epsilon}_i^2}{\sigma^2(1-h_{ii})} = \frac{[e_i^T(I-H)Y]^T[e_i^T(I-H)Y]/(1-h_{ii})}{\sigma^2}$$

where $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ Then

$$\frac{[e_i^T(I-H)Y]^T[e_i^T(I-H)Y]/(1-h_{ii})}{\sigma^2} = \frac{[Y^T(I-H)e_ie_i^T(I-H)Y]/(1-h_{ii})}{\sigma^2}$$

Then

$$B_i = \frac{[Y^T(I-H)Y]}{\sigma^2} - \frac{[Y^T(I-H)e_ie_i^T(I-H)Y]/(1-h_{ii})}{\sigma^2}$$

$$= \frac{Y^T}{\sigma} \left[(I-H) - (I-H) \frac{e_ie_i^T}{1-h_{ii}} (I-H) \right] \frac{Y}{\sigma}$$

$\frac{Y}{\sigma} \sim N_n(0, I)$.

To show that $C = \left[(I-H) - (I-H) \frac{e_ie_i^T}{1-h_{ii}} (I-H) \right]$ is a projection matrix, we need to show $C^2 = C, C^T = C$.

Here $(I-H)(I-H) = I-H, e_i^T(I-H)e_i = 1-h_{ii}$

Therefore, $B_i \sim \chi^2(r)$, where $r = \text{rank}[C]$. For projection matrix, the rank is also the trace.

$$\begin{aligned} \text{rank} \left[(I-H) - (I-H) \frac{e_ie_i^T}{1-h_{ii}} (I-H) \right] &= \text{tr} \left[(I-H) - (I-H) \frac{e_ie_i^T}{1-h_{ii}} (I-H) \right] \\ &= n-p - \frac{1}{1-h_{ii}} \text{tr}[(I-H)e_ie_i^T(I-H)] \\ &= n-p - \frac{1}{1-h_{ii}} \text{tr}[e_i^T(I-H)(I-H)e_i] \\ &= n-p - \frac{1}{1-h_{ii}} \text{tr}[e_i^T(I-H)e_i] \\ &= n-p - \frac{1-h_{ii}}{1-h_{ii}} \\ &= n-p-1 \end{aligned}$$

So $B_i \sim \chi^2(n-p-1)$.

2.2 b. Show A_i and B_i are independent

2.2.1 Preparation

Note that A_i and B_i are quadratic forms, which we need to show that their projection matrix product is 0, which will need to use linear algebra.

Another method is that, there are chi-square distribution, which we just need to show that the two statistics are ancillary.

Traditional method is to derive the distributions, and if we could write the distribution of $A_i + B_i$ as the product of distribution of A_i and B_i , then we can prove the independence.

One more question, are we able to use the moment generating function to do this? It is always easier to prove in MGF.

First, we need to write the MGF of A_i, B_i

$$M_{A_i} = E[\exp[A_i t]] = \int \exp[A_i t] \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right) dx$$

$$M_{B_i} = E[\exp[B_i t]] = \int \exp[B_i t] \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{x^2}{2\sigma^2}\right) dx$$

Then we can use MGF properties to prove

$$\begin{aligned} W_i &= A_i + B_i \\ M_{W_i} &= E[\exp(W_i t)] = E[\exp[(A_i + B_i)t]] \\ &= E[\exp[A_i t] \exp[B_i t]], \quad \text{all based on normal distribution} \\ &= E[\exp[A_i t]] E[\exp[B_i t]] = M_{A_i} M_{B_i} \end{aligned}$$

2.2.2 Proof

$$A_i = \frac{[Y^T (I - H) e_i e_i^T (I - H) Y] / (1 - h_{ii})}{\sigma^2}$$

$$B_i = \frac{[Y^T \left[(I - H) - (I - H) \frac{e_i e_i^T}{(1 - h_{ii})} (I - H) \right] Y]}{\sigma^2}$$

We would like to show the product of two projection matrix are 0

$$M_{A_i} M_{B_i} = (I - H) \frac{e_i e_i^T}{1 - h_{ii}} (I - H) \left[(I - H) - (I - H) \frac{e_i e_i^T}{(1 - h_{ii})} (I - H) \right]$$

2.3 Derive exact distribution

Let

$$r_i = \frac{\hat{\epsilon}_i}{\hat{\sigma} \sqrt{1 - h_{ii}}}$$

Using part(a) and (b), derive the exact distribution of $r_i^2/(n-p)$.

2.3.1 Question

To derive the exact the distribution, we can start from get the distribution of r_i . It is a normal distribution from part(a), then we can transform the distribution.

2.4 Outlier Model 1

Suppose we suspect that the i th case is an outlier and we consider the mean shift outlier model $Y = X\beta + d_i\phi + \epsilon$, where $\epsilon \sim N_n(0, \sigma^2 I)$, ϕ is an unknown scalar, and d_i is an $n \times 1$ vector with a 1 in the i th position and zeros elsewhere. Derive the maximum likelihood estimate of ϕ .

2.4.1 Question

We need to under the outlier model is introducing a new parameter ϕ and we need to estimate the parameter with the existing parameter β .

The way to get MLE is to construct likelihood function first, Y is a normal distribution, and only i th case is an outlier, while all other cases are not. Solve the problem similar as the random model.

2.5 Outlier Model 2

Suppose we wish to test $H_0 : \phi = 0$. Derive the test statistic for this hypothesis and derive its exact distribution under H_0 .

2.5.1 Question

The hypothesis test always comes with the estimate. There are several hypothesis test method we can do, wald test, score test and likelihood ratio test. The score test is generally the way as it is easy to get the estimate, score function and fisher information under H_0 .

2.6 Cook's Distance

Let $I = \{1, ..m\}$ be the subset of the first m cases in the dataset. Let D_I denote the Cook's distance based on simultaneously deleting m cases from the dataset, which is given by

$$D_I = \frac{(\hat{\beta} - \hat{\beta}_I)^T (X^T X)(\hat{\beta} - \hat{\beta}_I)}{p\hat{\sigma}^2}$$

where $\hat{\beta}^I$ denotes the least squares estimate of β with the cases deleted from set I and $\hat{\beta}$ denotes the estimate of β based on the full data. Show that D_I can be written as

$$D_I = \frac{1}{p} \sum_{i=1}^m h_i^2 \left(\frac{\lambda_i}{1 - \lambda_i} \right)$$

where the $\lambda_i, i = 1, ..m$, are the eigenvalues of the matrix $P_I = X_I(X^T X)^{-1} X_I^T$ based on a spectral decomposition of P_I .