




Georgia Tech College of Computing
School of Computational
Science and Engineering

Particle-in-cell in MFEM

Rushan Zhang, Joseph Signorelli, Ketan Mittal, Tzanio Kolev, Qi Tang

MFEM Workshop 2026

Particle tracing in MFEM



Particle Tracking in MFEM

Joseph Signorelli, Ketan Mittal, Tzanio Kolev

MFEM Workshop 2025


Particle tracing in MFEM

Known field $\rightarrow$ particle trajectories

Particles move through a field represented as an MFEM grid function.



Particle tracing provides

- **Particle class** with $\mathbf{x}$, $\mathbf{p}$, m and q etc.
- **Localization** of particles in the mesh
- **Interpolation** of fields at particle positions
- **Boris** push for particle $\mathbf{x}$ and $\mathbf{p}$ updates
- **Redistribution** of particles that cross subdomain boundaries



Tokamak magnetic field generated from: R. Zhang, G. Wimmer, Q. Tang. Structure-preserving transfer of Grad-Shafranov equilibria to magnetohydrodynamic solvers, Journal of Computational Physics, 568:115369, 2027.

Particle tracing in MFEM



$\begin{matrix} y \\ z \\ x \end{matrix}$

X. Garbet, Y. Idomura, L. Villard, T. H. Watanabe. Gyrokinetic simulations of turbulent transport, Nuclear Fusion, 50:043002, 2010.

Tokamak magnetic field generated from: R. Zhang, G. Wimmer, Q. Tang. Structure-preserving transfer of Grad-Shafranov equilibria to magnetohydrodynamic solvers, Journal of Computational Physics, 568:115369, 2027.

The natural next step: particle-in-cell

particles $\xrightarrow{\text{deposit charge}}$ field $\xrightarrow{\text{push}}$ particles

Repeat this feedback loop at every time step.

Governing Equations

Consider the Vlasov-Poisson equation:

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f + \frac{e}{m} \mathbf{E} \cdot \nabla_{\mathbf{v}} f = 0.$$

Distribution function approximated by macro-particles:

$$f \approx \sum_{p=1}^{N_p} \delta(\mathbf{x} - \mathbf{x}_p) \delta(\mathbf{v} - \mathbf{v}_p).$$

Integrate over velocity to get the charge density:

$$\rho(\mathbf{x}) = e \int f(\mathbf{x}, \mathbf{v}) d\mathbf{v} = e \sum_{p=1}^{N_p} \delta(\mathbf{x} - \mathbf{x}_p).$$

Governing Equations

Poisson's equation $-\epsilon_0 \Delta \phi(\mathbf{x}) = \rho(\mathbf{x}) - \rho_0$ yields

$$-\epsilon_0 \Delta \phi(\mathbf{x}) = e \sum_{p=1}^{N_p} \delta(\mathbf{x} - \mathbf{x}_p) - en_0.$$

For a test function $\varphi \in H^1(\Omega)$, the finite element solve is

$$\epsilon_0 \langle \nabla \varphi, \nabla \phi \rangle = e \sum_{p=1}^{N_p} \varphi(\mathbf{x}_p) - en_0 \int_{\Omega} \varphi.$$

OrthoSolver is needed to remove the constant nullspace of the periodic Poisson operator and ensure a unique solution for the potential ϕ .

Governing Equations

The electric field satisfies $\mathbf{E}(\mathbf{x}) = -\nabla\phi(\mathbf{x})$, so for a test function $\mathbf{v} \in H(\text{curl}; \Omega)$, we have

$$\langle \mathbf{v}, \mathbf{E} + \nabla\phi \rangle = 0.$$

The discrete de Rham complex gives a compatible gradient,

$$\begin{array}{ccc} H^1(\Omega) & \xrightarrow{\nabla} & H(\text{curl}; \Omega) \\ \downarrow & & \downarrow \\ CG & \xrightarrow{\nabla_h} & ND \end{array}$$

following which total energy can be shown to be conserved in the discrete PIC system.

One PIC time step in MFEM

- 1. Deposit charge:** For $\varphi \in H^1(\Omega)$, assemble $e \sum_p \varphi(\mathbf{x}_p)$.
- 2. Solve Poisson:** Solve $\epsilon_0 \langle \nabla \varphi, \nabla \phi \rangle = e \sum_p \varphi(\mathbf{x}_p) - en_0 \int_{\Omega} \varphi$.
- 3. Compute field:** $\langle \mathbf{v}, \mathbf{E} + \nabla \phi \rangle = 0$.
- 4. Gather field:** interpolate $\mathbf{E}$ at each $\mathbf{x}_p$.
- 5. Leapfrog push:**

$$\mathbf{p}_p^{t+\frac{1}{2}\Delta t} = \mathbf{p}_p^{t-\frac{1}{2}\Delta t} + e\mathbf{E}^t(\mathbf{x}_p^t)\Delta t,$$

$$\mathbf{x}_p^{t+1} = \mathbf{x}_p^t + \frac{\mathbf{p}_p^{t+\frac{1}{2}\Delta t}}{m_p}\Delta t.$$


- 6. Redistribute:** transfer particles that cross MPI subdomain boundaries.

Linear Landau damping



Field energy decays at a rate close to the reference rate.

256 MPI ranks · 20,000 particles per rank

64 × 64 grid · 400 steps · $\Delta t = 0.05$





Linear Landau damping



Left: kinetic and field energy exchange. Right: total energy. (Both normalized by the initial total energy.)

Parallel scaling



Left: weak scaling, 40,960 particles per rank. Right: strong scaling, 2,621,440 particles total.

Both use 8-128 MPI ranks and a 64×64 grid. Tested on NERSC.

Summary

- Scalable electrostatic PIC implementation in MFEM
- Validated with linear Landau damping

On-going work

- Improving **scalability** on CPU and **GPU** (See Eddy Luo and Elliot Day's poster presentation)
- Implementing a **shape function** beyond Dirac delta (See Rushan Zhang's poster presentation)

Questions?

