

Mobility and flexibility enable resilience of human harvesters to environmental perturbation.

Owen R. Liu¹, Mary Fisher^{2,3}, Blake E. Feist¹, Briana Abrahms⁴, Kate Richerson⁵, Jameal F. Samhouri¹

1 Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanographic and Atmospheric Administration, 2725 Montlake Blvd E, Seattle, Washington 98112 USA

2 School of Environmental and Forest Sciences, University of Washington, Seattle, WA 98195

3 NSF Graduate Research Internship Program, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Seattle, WA 98112

4 Center for Ecosystem Sentinels, Department of Biology, University of Washington, Seattle, WA 98195.

5 Fishery Resource Analysis and Monitoring Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanographic and Atmospheric Administration, Newport, OR, 97365

*** Corresponding author:** Owen R. Liu; Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanographic and Atmospheric Administration, 2725 Montlake Blvd E, Seattle, Washington 98112 USA; 781-622-8343; owen.liu@noaa.gov

Classification: Biological Sciences | Sustainability Science

Keywords: climate change adaptation | environmental perturbation | marine heatwave | fisheries dynamics

Author contributions: O.R.L., M.F., B.E.F., B.A., K.R., and J.F.S. designed research; O.R.L. and J.F.S. performed research and analyzed data; O.R.L., M.F., B.E.F., B.A., K.R., and J.F.S. wrote the paper.

This PDF file includes:

Main text, including all main text figures

Supplemental tables and figures