


# Fast Algorithms for Geometric Consensuses

---


Sariel Har-Peled<sup>1</sup>   Mitchell Jones<sup>1</sup>

SoCG 2020, June 23–26


<sup>1</sup>University of Illinois at Urbana-Champaign



# The yolk: problem setup




# The yolk: problem setup




- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side

# The yolk: problem setup




- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side

# The yolk: problem setup




- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side
- $\ell$  is **extremal** if it passes through 2 points of  $P$

# The yolk: problem setup




- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side
- $\ell$  is **extremal** if it passes through 2 points of  $P$
- **Extremal yolk** is smallest radius ball intersecting all extremal median lines

# The yolk: problem setup



- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side
- $\ell$  is **extremal** if it passes through 2 points of  $P$
- **Extremal yolk** is smallest radius ball intersecting all extremal median lines

# The yolk: problem setup



- $\ell$  is a **median line** if  $\geq \lceil n/2 \rceil$  points of  $P$  lie on either side
- $\ell$  is **extremal** if it passes through 2 points of  $P$
- **Extremal yolk** is smallest radius ball intersecting all extremal median lines




- Yolk is smallest radius ball intersecting all median lines

- Yolk is smallest radius ball intersecting all median lines
- Motivation: voting theory (generalizes plurality point)

- Yolk is smallest radius ball intersecting all median lines
- Motivation: voting theory (generalizes plurality point)

yolk  $\neq$  extremal yolk [Stone and Tovey, 1992]



Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]

Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

$(1 + \varepsilon)$ -apx in  $\mathbb{R}^2$ :

Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

$(1 + \varepsilon)$ -apx in  $\mathbb{R}^2$ :

- $O(n \log^7 n \log^4(1/\varepsilon))$  for **yolk** [Gudmundsson and Wong, 2019]

Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

$(1 + \varepsilon)$ -apx in  $\mathbb{R}^2$ :

- $O(n \log^7 n \log^4(1/\varepsilon))$  for **yolk** [Gudmundsson and Wong, 2019]
- $O((n/\varepsilon^3) \log^3 n)$  for **extremal yolk** [Berg, Chung, et al., 2019]



Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

$(1 + \varepsilon)$ -apx in  $\mathbb{R}^2$ :

- $O(n \log^7 n \log^4(1/\varepsilon))$  for **yolk** [Gudmundsson and Wong, 2019]
- $O((n/\varepsilon^3) \log^3 n)$  for **extremal yolk** [Berg, Chung, et al., 2019]

## Our result

In  $\mathbb{R}^2$ :  $O(n \log n)$  expected time for yolk/extremal yolk

Exact:

- $O_d(n^{(d+1)^2})$  for **yolk** (improved to  $O(n^4)$  in  $\mathbb{R}^2$ ) [Tovey, 1992]
- $O(n^{4/3} \log^{O(1)} n)$  in  $\mathbb{R}^2$  for **yolk and extremal yolk** [Berg, Gudmundsson, et al., 2018]

$(1 + \varepsilon)$ -apx in  $\mathbb{R}^2$ :


- $O(n \log^7 n \log^4(1/\varepsilon))$  for **yolk** [Gudmundsson and Wong, 2019]
- $O((n/\varepsilon^3) \log^3 n)$  for **extremal yolk** [Berg, Chung, et al., 2019]

## Our result


In  $\mathbb{R}^2$ :  $O(n \log n)$  expected time for yolk/extremal yolk, and  $O_d(n^{d-1} \log n)$  for  $\mathbb{R}^d$

- Extremal median lines induced by 2 points of  $P$   
 $\implies \leq \binom{n}{2}$  **implicit** constraints

- Extremal median lines induced by 2 points of  $P$   
 $\implies \leq \binom{n}{2}$  **implicit** constraints




- Extremal median lines induced by 2 points of  $P$   
 $\implies \leq \binom{n}{2}$  **implicit** constraints



- Solve LP in  $O(n^2)$  time

- Extremal median lines induced by 2 points of  $P$   
 $\implies \leq \binom{n}{2}$  **implicit** constraints



- Solve LP in  $O(n^2)$  time
- Faster?

- Uses Chan's framework for implicit LPs [Chan, 2004]

# Solving the implicit LP



- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$



# Solving the implicit LP



- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$

# Solving the implicit LP



- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

(a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ?

- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ?
- (b) Given  $Q$ : construct  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $L(Q) = \cup_{i=1}^r L(Q_i)$

- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ?
- (b) Given  $Q$ : construct  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $L(Q) = \cup_{i=1}^r L(Q_i)$

**Result (using [Chan, 2004])**

Given  $P$ , can compute extremal yolk in  $O(D(n))$  expected time.

- Uses Chan's framework for implicit LPs [Chan, 2004]
- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- **Goal:** compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,


- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ?
- (b) Given  $Q$ : **construct**  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $L(Q) = \cup_{i=1}^r L(Q_i)$

**Result (using [Chan, 2004])**

Given  $P$ , can compute extremal yolk in  $O(D(n))$  expected time.

## Definition: $k$ -levels


Given lines  $L$ ,  $k$ -level = {points lying above or on  $k$  lines of  $L$ }.






## Definition: $k$ -levels

Given lines  $L$ ,  $k$ -level = {points lying above or on  $k$  lines of  $L$ }.




## Definition: $k$ -levels

Given lines  $L$ ,  $k$ -level = {points lying above or on  $k$  lines of  $L$ }.




## Definition: $k$ -levels

Given lines  $L$ ,  $k$ -level = {points lying above or on  $k$  lines of  $L$ }.




## Definition: $k$ -levels

Given lines  $L$ ,  $k$ -level = {points lying above or on  $k$  lines of  $L$ }.



### Definition: Zone of a surface

Given lines  $L$ , curve  $\gamma$ , the zone  $\mathcal{Z}(\gamma, L)$  are cells of  $\mathcal{A}(L)$  intersecting  $\gamma$ .



Lemma [Aronov et al., 1993, Berg, Dobrindt, et al., 1995]

$\mathcal{Z}(\gamma, L)$  can be computed in  $O(n \log n)$  expected time.

- $P$  points,  $L(P)$  extremal median lines

# Computing the extremal yolk





- $P$  points,  $L(P)$  extremal median lines
- In the dual:

# Computing the extremal yolk



- $P$  points,  $L(P)$  extremal median lines
- In the dual:
  - Points  $\iff$  lines







# Computing the extremal yolk





- $P$  points,  $L(P)$  extremal median lines
- In the dual:
  - Points  $\iff$  lines
  - Ball  $\iff$  region  $R$  between hyperbola




# Computing the extremal yolk



- $P$  points,  $L(P)$  extremal median lines
- In the dual:
  - Points  $\iff$  lines
  - Ball  $\iff$  region  $R$  between hyperbola
  - Extremal median line  $\iff$  vertex of the  $n/2$ -level




# The key subproblem



Is there a vertex of the  $n/2$ -level **outside**  $R$ ?

# The key subproblem




Is there a vertex of the  $n/2$ -level **outside**  $R$ ?

Check vertices of  $\mathcal{A}(L(P))$  near **boundary** of  $R$ !

# The key subproblem




**Task:** Is there a vertex of  $n/2$ -level in  $\mathbb{R}^2 \setminus R$ ?



# The key subproblem



**Task:** Is there a vertex of  $n/2$ -level in  $\mathbb{R}^2 \setminus R$ ?




**Idea:** Compute zone of  $\partial R$

# The key subproblem



**Task:** Is there a vertex of  $n/2$ -level in  $\mathbb{R}^2 \setminus R$ ?




**Idea:** Compute **zone** of  $\partial R$ , walk around vertices in **zone**

# The key subproblem



**Task:** Is there a vertex of  $n/2$ -level in  $\mathbb{R}^2 \setminus R$ ?




**Idea:** Compute **zone** of  $\partial R$ , walk around vertices in **zone**, check if any vertex in  $\mathbb{R}^2 \setminus R$  has level  $n/2$ .



# The key subproblem



**Task:** Is there a vertex of  $n/2$ -level in  $\mathbb{R}^2 \setminus R$ ?



**Idea:** Compute **zone** of  $\partial R$ , walk around vertices in **zone**, check if any vertex in  $\mathbb{R}^2 \setminus R$  has level  $n/2$ .

**Result:** Decider takes  $O(n \log n)$  time.

# Solving the implicit LP



- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- Goal: compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ?
- (b) Given  $Q$ : **construct**  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $L(Q) = \cup_{i=1}^r L(Q_i)$

# Solving the implicit LP



- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- Goal: compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ? ✓
- (b) Given  $Q$ : construct  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $L(Q) = \cup_{i=1}^r L(Q_i)$

- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- Goal: compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ? ✓
- (b) Given  $Q$ : construct  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $H(Q) = \cup_{i=1}^r H(Q_i)$  ✓

- $P$ : a set of points;  $n = |P|$
- $L(P)$ : extremal median lines induced by  $P$
- Goal: compute a solution for  $L(P)$

For any  $Q \subseteq P$ ,  $m = |Q|$ , assume in  $D(m)$  time,

- (a) Given  $Q$ , candidate ball  $b$ : does  $b$  violate  $L(Q)$ ? ✓
- (b) Given  $Q$ : construct  $O(1)$  sets  $Q_1, \dots, Q_r$ ,  $|Q_i| \leq \alpha m$ ,  
 $H(Q) = \cup_{i=1}^r H(Q_i)$  ✓

## Our result


$$D(m) = O_d(m^{d-1} \log m) \implies$$

extremal yolk in  $O_d(n^{d-1} \log n)$  expected time.

- Yolk: Smallest ball intersecting all **median** hyperplanes

- Yolk: Smallest ball intersecting all **median** hyperplanes
- Egg: Smallest ball intersecting all **extremal** hyperplanes

- Yolk: Smallest ball intersecting all **median** hyperplanes
- Egg: Smallest ball intersecting all **extremal** hyperplanes
  - Easy modification! Check if any vertex of  $\mathcal{A}(L(P))$  lies **outside**  $R$



$\Rightarrow O_d(n^{d-1} \log n)$  expected time



- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$

- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$
- E.g.,  $\mathcal{T}_1 = \text{conv}(P)$ ,  $\mathcal{T}_{n/(d+1)}$  contains centerpoint of  $P$

- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$
- E.g.,  $\mathcal{T}_1 = \text{conv}(P)$ ,  $\mathcal{T}_{n/(d+1)}$  contains centerpoint of  $P$
- If  $\mathcal{T}_k \neq \emptyset$ , **center ball** = largest ball inside  $\mathcal{T}_k$

- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$
- E.g.,  $\mathcal{T}_1 = \text{conv}(P)$ ,  $\mathcal{T}_{n/(d+1)}$  contains centerpoint of  $P$
- If  $\mathcal{T}_k \neq \emptyset$ , **center ball** = largest ball inside  $\mathcal{T}_k$
- If  $\mathcal{T}_k = \emptyset$ , **Tukey ball** = smallest ball intersecting all halfspaces  $H_k(P)$

- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$
- E.g.,  $\mathcal{T}_1 = \text{conv}(P)$ ,  $\mathcal{T}_{n/(d+1)}$  contains centerpoint of  $P$
- If  $\mathcal{T}_k \neq \emptyset$ , **center ball** = largest ball inside  $\mathcal{T}_k$
- If  $\mathcal{T}_k = \emptyset$ , **Tukey ball** = smallest ball intersecting all halfspaces  $H_k(P)$

### Our result

Center ball and Tukey ball can be computed in  $O_d(n^{d-1} \log n)$  expected time











- $H_k(P) = \{\text{open halfspaces containing } > n - k \text{ points of } P\}$
- $\mathcal{T}_k = \bigcap \{h \mid h \in H_k(P)\}$
- E.g.,  $\mathcal{T}_1 = \text{conv}(P)$ ,  $\mathcal{T}_{n/(d+1)}$  contains centerpoint of  $P$
- If  $\mathcal{T}_k \neq \emptyset$ , **center ball** = largest ball inside  $\mathcal{T}_k$
- If  $\mathcal{T}_k = \emptyset$ , **Tukey ball** = smallest ball intersecting all halfspaces  $H_k(P)$

### Our result

Center ball and Tukey ball can be computed in  $O_d(n^{d-1} \log n)$  expected time

Thank you!

-  Richard E. Stone and Craig A. Tovey. *Limiting median lines do not suffice to determine the yolk*. *Social Choice and Welfare*, 9(1): 33–35, 1992.
-  Craig A. Tovey. *A polynomial-time algorithm for computing the yolk in fixed dimension*. *Math. Program.*, 57: 259–277, 1992.
-  Mark de Berg, Joachim Gudmundsson, and Mehran Mehr. *Faster algorithms for computing plurality points*. *ACM Trans. Algorithms*, 14(3): 36:1–36:23, 2018.
-  Joachim Gudmundsson and Sampson Wong. *Computing the yolk in spatial voting games without computing median lines*. *33th Conf. Artificial Intell. (AAAI)*,

-  Mark de Berg, Jonathan Chung, and Joachim Gudmundsson. *Computing the Yolk in Spatial Voting Games*. 2019.
-  Timothy M. Chan. *An optimal randomized algorithm for maximum Tukey depth*. *Proc. 15th ACM-SIAM Sympos. Discrete Algs. (SODA)*, 430–436, 2004.
-  Boris Aronov, Marco Pellegrini, and Micha Sharir. *On the zone of a surface in a hyperplane arrangement*. *Discrete Comput. Geom.*, 9: 177–186, 1993.
-  Mark de Berg, Katrin Dobrindt, and Otfried Schwarzkopf. *On lazy randomized incremental construction*. *Discrete Comput. Geom.*, 14(3): 261–286, 1995.