

Commercial Fish Harvest ~ Water Quality

Shallow Natural Lakes of Northwest Iowa

Matthew Foli & Martin Simonson

Overall Objective

Our aim is to quantify relationships between commercial harvest of “rough fish” (common carp and bigmouth buffalo) against a suite of water quality variables. The term *water quality* itself can be subjective, but here we have identified multiple potential response variables that are often associated with “clean” or “impaired” water body designations. They fall into three broad categories:

- **Nutrients:** Trophic State Index (TSI, Carlson 1977); Total Phosphorous ($\mu\text{g/L}$); Total Nitrogen (mg/L)
- **Physical:** Secchi depth (m , *meters*); Total Suspended Solids ($\mu\text{g/L}$); Dissolved Oxygen (mg/L)
- **Biological:** Total Zooplankton Biomass; Total Phytoplankton Biomass; Phytoplankton Biomass of *Cyanophyta*; Phytoplankton Biomass of *Chlorophyta*

Study Area

Marty’s dissertation research is focused on shallow, natural lakes of Iowa. Commercial Harvest of bigmouth buffalo occurs in some 20 lakes and rivers in Iowa, and over half of them fall into the shallow natural lake category. Since 1) Common carp are strongly suspected to disturb ecosystems and force shallow lakes into a turbid, algae-dominated state and 2) Bigmouth Buffalo may regulate zooplankton that graze on algae, the Iowa DNR is interested to see if managing these populations of fish can restore elusive water quality in Iowa and shift these lakes into a clear-water macrophyte and piscivore dominated lake systems.

Commercial harvest of these species is complex and variable with weather conditions, equipment, staffing, invasive species control, permitting, supply chain management, and market fluctuations affecting how much effort commercial anglers can put into catching fish, and where they focus that effort. Fishing is restricted to times of year when water temperatures are below 70°F , or basically between Memorial Day and October-ish. That being said, *there might be some overlap between harvest dates and ambient lake monitoring samples (described below)*.

One final note on the study area: We can incorporate other lakes that fit our selection criteria but do not have commercial harvest of rough fish species. Not that they are a perfect control, but it could help explain any year-to-year variation in response variables across our study region.

Introducing the Data

Harvest Data

We do not have abundance data for carp and buffalo for more than a couple of years in a subset of lakes. However, based on available data from the Spirit Lake District Fisheries Office, we have total pounds removed at multiple time points spanning 25 years in about a dozen lakes. We have identified a subset of these lakes based on similarity in geographic location, lake morphometry, fish communities, and regular harvest periods. Data is provided in total pounds removed, however due to differences in lake size and capacity to hold fish, we will standardize data by kilograms per hectare. The lakes are summarized here:

Lake	Center Lake	Five Island Lake	Ingham Lake	Lost Island Lake	Silver Lake (Dickinson County)
Area (hectare)	89	390	150	470	418
Area (m^2)	890,308	3,901,170	1,497,337	4,702,447	4,176,356
Watershed Area (hectare)	296	3516	129	1838	6890
Watershed:Lake Ratio	3.3:1	9.2:1	0.9:1	3.9:1	16.5:1
Mean Depth (m)	2.9	1.5	1.9	3.1	2
Max Depth (m)	4.7	6.1	3.7	4.3	3
Row Crop Usage (hectare)		2657	55		5876
Percent Row Crop		75.6	42.2		85.3
Shoreline (m)	4,023	22,853	6,828	11,748	15,463
Shoreline Complexity	1.20	3.26	1.57	1.53	2.13

- *Note that Center Lake TMDL reports are scheduled to begin in the next couple of years, and not all information is readily available*

Water Quality Data

Water quality data (summarized below) is available from ISU and Iowa DNR through an online database and the Limnology Laboratory at ISU and provided by Dr. Grace Wilkonson, EEOB. This is part of the Ambient Lake Monitoring Program that collects a suite of variables at 130 lakes in Iowa that are designated as “Principal Recreation Lakes” and data is available from 2000 to 2015, except in 2008. Further data (2016-2019) may also be available for inclusion toward the end of this project’s timeframe. These lakes are visited 2-3 (almost always 3) times per year from spring to fall and in-situ measurements are taken as well as samples removed for laboratory analysis of biological and chemical components.

A Trophic State Index (TSI) can also be calculated from Chlorophyll a , Secchi Depth and Total Phosphorus. The index formulas were designed so that a difference of 10 units corresponds to a doubling of algal biomass. We chose to pick chlorophyll a as a initial response variable, because the the developer of the index (Carlson 1977) said later in 1983 that averaging the three values was not appropriate, and chlorophyll a was the best measurement to use out of the three. We do not expect the TSI to decrease out of the eutrophic range, but a directional shift in TSI may be indicative of the desired change in water quality from biomanipulation of carp and buffalo.

Lakes fluctuate over time in cycles of stratification, turnover, zooplankton and algal blooms, etc. so these snapshots in time can be problematic. Still, we have identified a suite of variables that we expect to respond to the biomanipulation (harvest) of carp and buffalo. Also, these variables are common metrics used to designate water bodies as “degraded” or “healthy” and could provide insight into food web processes in the lake. The variables are summarized here:


Variable	Units	Hypothesized short-term response
Secchi Depth	meters	Increase from carp removal, no difference from buffalo
Total Suspended Solids	$\mu g/L$	Decrease from carp removal, no difference from buffalo
Dissolved Oxygen	mg/L	Decrease if large buffalo removal caused enough increase in zooplankton to reduce O ₂ production
Total Phosphorous	$\mu g/L$	Decrease with carp removal (resuspension)
Total Nitrogen	mg/L	Decrease with carp removal (resuspension)

Variable	Units	Hypothesized short-term response
Zooplankton Biomass	$\mu g/L$	Increase with buffalo removal (trophic release)
Chlorophyll <i>a</i>	$\mu g/L$	Decrease with carp and buffalo removal
Phytoplankton Biomass	mg/L	Decrease with both carp and buffalo removal (bottom up and top down pathways)
<i>Cyanophyta</i>	mg/L	Decrease with carp and buffalo removal (nutrient shift and zoop shift)
<i>Chlorophyta</i>	mg/L	Decrease with both carp and buffalo removal
Trophic State Index	Calculated from Chl <i>a</i>	Decrease with both carp and buffalo removal; but not out of hyper-eutrophic state.

Data Analysis

- Predictor variable data comes from commercial harvest reports from 1997 to 2019 out of 5 Iowa Lakes
- For background, we plot the distribution of harvest in total pounds removed (left) and pounds per acre removed (right). The number of bars across the box plots indicate the number of harvest dates in that year due to “stacking” multiple harvests within a year. The colors represent the different species.
- Clearly, effort expended and harvest success are highly variable through time.
- Note the incentivized harvest in Lost Island Lake in 2010 and 2011.
- Also note non-incentivized contracts at Ingham 1999 & 2019 as well as Center 2016 approached the same biomass density removals as Lost Island Lake (357 kg/hectare).

Harvest through time



Update 2/24/2020

After meeting with Audrey and Phil, we have a new format for how to arrange the data. The general model format is:

$$TSI \approx \beta_0 + L_i + (\beta_c * C_{li} * H_i) + (\beta_b * (1 - C_{li}) * H_i)$$

where i is indexed on year, lowercase l is indexed on lake, uppercase L is a fixed lake effect, C is the proportion of harvest made up by carp, $1 - C$ is the proportion of harvest made up by buffalo, H is the total harvest. *In this example, TSI is used as response variable but any of the other 10 water quality variables may be used.*

Mesurements of TSI and other variables are taken from the June Ambient Lake Monitoring measurements. When a measurement from June is not available, we took the next closest measurement to within 365 days after the previous year's measurement date.

Completed Action Items:

- Calculate TSI from Chl a formula.
- From those TSI values, calculate the *change* in TSI from year A to year B.
- Apply same change in June measurement for other water quality variables between year A and year B.
- Harvest data above began in 1999 to match water quality data; our data set starts in year “2000” which represents the commercial fish removal from September 1999 to May 2000. This is coupled with the water quality variable for June 2000, and so on.

New data frame:


- *Lake*: Name of each lake.
- *Year*: Year B for water quality response variables (represents the change in TSI, etc. between year A and B). For harvest data, this value is the total and proportional harvests amounts between years A and B (September A to April/May B).
- *Carp Harvest*: The amount (kilograms per hectare) of carp harvested between years A and B.
- *Buffalo Harvest*: Same as immediately above, for buffalo.
- *Total Removed*: Per the model above, variable H_i (kilograms per hectare).
- *Proportion of Carp*: Per the model above, the proportion (0 to 1) of the harvest made up of common carp, variable C_{li} .
- *Proportion of Buffalo*: Per the model above, and complement to previous bullet point, the proportion (0 to 1) of the harvest made up of bigmouth buffalo, variable $1 - C_{li}$.
- *Delta TSI Chl. a*: Change in Trophic State Index calculated from Chlorophyll a . Each 10 unit increase represents a doubling of algal biomass. Lakes over 50 TSI have characteristics of eutrophic lakes. Represents June measurement in year B minus June measurement in Year A.
- *Delta Chla, TSS, TN, TP, SD (Secchi Depth), Dissolved Oxygen, Turbidity, Phytoplankton biomass, Zooplankton biomass, Chlorophyta density, Cyanophyta density*: Change in June (or closest) measurement of the variable from year A to B. For Year 2000, this represents the measurement in year 2000 minus the measurement in 1999.

Loading in Yearly Change Dataset


Distribution of response variables

These represent the count (frequency distribution) for each response variable along its range. The function ‘geom_density’ from ggplot2 is a smoothed version of the histogram, useful for continuous random variables (shaded red area on figures, not always visible). Note that these are centered at around zero since we are examining the change in these variables from year to year


Distribution of Physical Response Variables



Distribution of TSI and Nutrient Response Variables



Distribution of Biotic Response Variables



Pairs Plots - correlation of variables

From the package “ggAlly” this displays both the paired scatterplots and correlation coefficients of all pairs of data. Instead of displaying a 12x12 grid of 144 plots, we’ll split them into our three categories: Physical, Nutrient, and Biotic.

