SWITCHING RECONSTRUCTION OF ORIENTED GRAPHS

Milana Golich
with Muse Spark and Watermelon via meta.ai–TBD

Orchestrators
Charles Arnal, Vivien Cabannes

Reviewed by

Abstract. We prove the existence of a switching, non-reconstructible digraph on $n \geq 12$ vertices. We remark on the use of AI in this research, see Statement of AI Use.

1. Introduction

Human

A natural question to ask is to what extent a mathematical object can be recovered by its properties. If a mathematical property uniquely determines the object (up to categorical equivalence), then identifying the object can be reduced to simply computing an invariant. Here we are concerned with this question in the graph theoretic setting, and we refer to such deterministic objects as reconstructible. The graph property that we explore is its "deck", a multi-set of objects formed from the original graph. There are many examples of decks that have been studied. For instance, Kelly and Ulam ([Kel57], [Ula60]) investigated the deck consisting of the multi-set of subgraphs each obtained from deleting a single vertex from the original graph. This led to the following conjecture which is referred to as the graph reconstruction conjecture:

Conjecture 1.1 (Kelly, Ulam)**.** *Every finite simple graph on at least 3 vertices is reconstructible from its multiset of 1-vertex-deleted subgraphs.*

Similar reconstruction questions to that of Kelly and Ulam were asked for various other kinds of decks. For example, Stanley [Sta85] asked an analagous question for graphs with decks that are obtained from a 'switching'. Here Stanley defined the switching of a graph about a vertex to be the switching of edges incident with a vertex into non-edges, and vice-versa. Bondy and Mercier [BM11] continued to ask switching reconstruction problems in the setting of directed graphs, or digraphs. The switching in this context refers to switching the orientation of the directed edges incident with a vertex. In [BM11], the authors proved the existence of large sets of digraphs on 4 and 8 vertices that were switching, non-reconstructible. In fact, it was shown that any switching, non-reconstructible digraph must have order divisible by 4, but there were no known examples with $n \geq 12$ vertices. In this exposition, we present an example for $n = 12$. We also refer the reader to [MS14] for additional background on this topic.

Statement of AI Use

This paper was developed through collaboration between researchers and Watermelon and Muse Spark (AI models) via the meta.ai chat interface. The model helped explore ideas, develop candidate arguments, and draft material. Researchers guided the work, checked and corrected the mathematics, and take ownership for the final manuscript. Following [Sch25], margin markers identify human-drafted (**Human**) , joint(Human + AI), and AI-assisted material (**AI**) separately.

2. Problem Statement

Human + AI

For a digraph D and a vertex v, we define switch(D, v) to be the graph obtained by the **switching in** D **about** v where every edge incident with v is reversed and all other edges remain unchanged. A pair of digraphs D and H have the same switching deck if there exists a bijection $\pi : V(D) \to V(H)$ such that switch$(D, v) \cong$ switch$(H, \pi(v))$ for every v. We say that a digraph D is **non-reconstructible** if there

2020 *Mathematics Subject Classification.* 00A05.

Human + AI

exists a digraph H, not isomorphic to D, that has the same switching deck as D.

The following question was posed in [BM11]:

QUESTION. Does there exist a switching, non-reconstructible digraph on twelve or more vertices?

3. Solution

We exhibit a switching, non-reconstructible oriented graph on $n = 12$ vertices. That is, we present a pair of non-isomorphic oriented graphs D and H on 12 vertices that have the same switching deck.

We recall that an **oriented graph** is a digraph with no loops and no symmetric pairs of edges. That is, for an n-vertex oriented graph, its associated $n \times n$ signed adjacency matrix A is skew-symmetric $A^T = -A$ with entries $A_{ij} \in \{-1, 0, 1\}$ and $A_{ii} = 0$. For $i \neq j$, $A_{ij} = 0$ if there exists no edge between vertices v_i and v_j.

ORIENTED GRAPHS D and H. We define D to be the oriented graph of order 12 with vertices $V = \{v_0, v_1, v_2, ..., v_{11}\}$ and skew-symmetric signed adjacency matrix A:

$$A = [A_{ij}] = \begin{pmatrix} 0 & -1 & -1 & 0 & -1 & 1 & -1 & 1 & 1 & -1 & 1 & -1 \\ 1 & 0 & 1 & -1 & 0 & -1 & 1 & -1 & -1 & 1 & -1 & 1 \\ 1 & -1 & 0 & -1 & -1 & 0 & 1 & -1 & -1 & 1 & -1 & 1 \\ 0 & 1 & 1 & 0 & 1 & -1 & -1 & 1 & 1 & -1 & 1 & -1 \\ 1 & 0 & 1 & -1 & 0 & -1 & -1 & 1 & 1 & -1 & 1 & -1 \\ -1 & 1 & 0 & 1 & 1 & 0 & 1 & -1 & -1 & 1 & -1 & 1 \\ 1 & -1 & -1 & 1 & 1 & -1 & 0 & -1 & 1 & 0 & -1 & -1 \\ -1 & 1 & 1 & -1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 & -1 \\ -1 & 1 & 1 & -1 & -1 & 1 & -1 & -1 & 0 & -1 & -1 & 0 \\ 1 & -1 & -1 & 1 & 1 & -1 & 0 & -1 & 1 & 0 & -1 & -1 \\ -1 & 1 & 1 & -1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 & -1 \\ 1 & -1 & -1 & 1 & 1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 \end{pmatrix}$$

We can see from the above matrix that the only non-adjacent, distinct vertices v_i and v_j in D are given by the following six pairs:

$$(i, j) = \{(0, 3), (1, 4), (2, 5), (6, 9), (7, 10), (8, 11)\}.$$

Equivalently, we can derive the directed edges of D from A as the following:

$$\begin{aligned} &0 \to \{5, 7, 8, 10\}, && 1 \to \{0, 2, 6, 9, 11\}, && 2 \to \{0, 6, 9, 11\}, \\ &3 \to \{1, 2, 4, 7, 8, 10\}, && 4 \to \{0, 2, 7, 8, 10\}, && 5 \to \{1, 3, 4, 6, 9, 11\}, \\ &6 \to \{0, 3, 4, 8\}, && 7 \to \{1, 2, 5, 6, 8, 9\}, && 8 \to \{1, 2, 5\}, \\ &9 \to \{0, 3, 4, 8\}, && 10 \to \{1, 2, 5, 6, 8, 9\}, && 11 \to \{0, 3, 4, 6, 7, 9, 10\}. \end{aligned}$$

Next, we obtain an oriented graph H by switching the oriented graph D at the vertices v_0 and v_1. By definition, this means that H is represented by the skew-symmetric signed adjacency matrix C given by

$$C = [C_{ij}], \quad C_{ij} = \begin{cases} -A_{ij} & \text{if exactly one of } i, j \in \{0, 1\}, \\ A_{ij} & \text{otherwise.} \end{cases}$$

This gives the following matrix representation of H:

Human + AI

$$C = [C_{ij}] = \begin{pmatrix} 0 & -1 & 1 & 0 & 1 & -1 & 1 & -1 & -1 & 1 & -1 & 1 \\ 1 & 0 & -1 & 1 & 0 & 1 & -1 & 1 & 1 & -1 & 1 & -1 \\ -1 & 1 & 0 & -1 & -1 & 0 & 1 & -1 & -1 & 1 & -1 & 1 \\ 0 & -1 & 1 & 0 & 1 & -1 & -1 & 1 & 1 & -1 & 1 & -1 \\ -1 & 0 & 1 & -1 & 0 & -1 & -1 & 1 & 1 & -1 & 1 & -1 \\ 1 & -1 & 0 & 1 & 1 & 0 & 1 & -1 & -1 & 1 & -1 & 1 \\ -1 & 1 & -1 & 1 & 1 & -1 & 0 & -1 & 1 & 0 & -1 & -1 \\ 1 & -1 & 1 & -1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 & -1 \\ 1 & -1 & 1 & -1 & -1 & 1 & -1 & -1 & 0 & -1 & -1 & 0 \\ -1 & 1 & -1 & 1 & 1 & -1 & 0 & -1 & 1 & 0 & -1 & -1 \\ 1 & -1 & 1 & -1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 & -1 \\ -1 & 1 & -1 & 1 & 1 & -1 & 1 & 1 & 0 & 1 & 1 & 0 \end{pmatrix}$$

SAME SWITCHING DECK. For simplicity, we refer to the vertice v_i, or merely v, as i. Based on the derivation of H, we naturally define the bijection $\pi : V(D) \to V(H)$ to be the permutation $\pi = (0\,1)(2\,5)(3\,4)(6\,7)(8\,11)(9\,10)$. This gives

$$\pi(V) = \{\pi(0), \pi(1), \pi(2), ..., \pi(11)\} = \{1, 0, 5, 4, 3, 2, 7, 6, 11, 10, 9, 8\}.$$

Under π, we consider the signed adjacency matrices of switch(D, v) and switch$(H, \pi(v))$ for each $v \in V$, which we will denote by D_v and $H_{\pi(v)}$ respectively. First, we note that the switching in D about v results in multiplying the signed adjacency matrix A by -1 in the row and column corresponding to v. Equivalently, this corresponds to the following matrix multiplication:

$$D_v = S_v A S_v \quad \text{where} \quad S_v = \text{diag}(s_0, \ldots, s_{11}),$$

$$s_i = \begin{cases} -1 & i = v \\ 1 & i \neq v \end{cases} = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & \underset{\text{position } v}{-1} & & & \\ & & & \ddots & & \\ & & & & 1 & \end{pmatrix}_{12\times 12}$$

Thus given that H was obtained from switching D about the vertices v_0 and v_1, we can realize the signed adjacency matrix C as the multiplication of A by two switching matrices S_0 and S_1 as follows:

$$C = S_{01} A S_{01}, \quad S_{01} = S_0 S_1$$

Furthermore, we can realize D_v and $H_{\pi(v)}$ at $v = 0$ as the following:

$$D_0 = S_0 A S_0 = S_1 S_{01} A S_{01} S_1 = S_1 C S_1 = H_1 = H_{\pi(0)}.$$

Hence for the switching at $v = 0$, we act on D_0 by conjugation of the identity permutation matrix P_0 which represents the trivial action on the vertices V:

$$P_0^T D_0 P_0 = H_1$$

In this way, we naturally define the bijection $\psi_v : V(\text{switch}(D, v)) \to V(\text{switch}(H, \pi(v)))$ to be the permutation which induces the conjugation action on D_v by the respective permutation matrices for each v:

$$\Psi_v = P_{\psi(v)}^T D_v P_{\psi(v)} = H_{\pi(v)}$$

In other words, P_{ψ_v} is the matrix representation of ψ_v. For convenience, we will write the permutation ψ_v of V as the list $[\psi_v(0), \ldots, \psi_v(11)]$. For each switching of D and H under π, we may continue this computational exercise to derive ψ_v for each v:

Human + AI

v	$\pi(v)$	ψ_v
0	1	$[0,1,2,3,4,5,6,7,8,9,10,11]$
1	0	$[3,4,5,0,1,2,6,7,8,9,10,11]$
2	5	$[1,2,3,4,5,0,6,7,8,9,10,11]$
3	4	$[3,4,5,0,1,2,6,7,8,9,10,11]$
4	3	$[4,5,0,1,2,3,6,7,8,9,10,11]$
5	2	$[4,5,0,1,2,3,6,7,8,9,10,11]$
6	7	$[2,3,4,5,0,1,11,6,7,8,9,10]$
7	6	$[2,3,4,5,0,1,7,8,9,10,11,6]$
8	11	$[2,3,4,5,0,1,11,6,7,8,9,10]$
9	10	$[2,3,4,5,0,1,8,9,10,11,6,7]$
10	9	$[2,3,4,5,0,1,10,11,6,7,8,9]$
11	8	$[2,3,4,5,0,1,10,11,6,7,8,9]$

Claim 3.1. In terms of the corresponding adjacency matrices, we have for every v and all i,j that

$$[D_v]_{ij} = [H_{\pi(v)}]_{\psi_v(i),\psi_v(j)}.$$

By definition, this implies that ψ_v is an isomorphism between the oriented graphs $\text{switch}(D,v)$ and $\text{switch}(H,\pi(v))$ for each v. Equivalently, $\Psi_v : D_v \to H_{\pi(v)}$ is an isomorphism for each v.

D IS NOT ISOMORPHIC TO H. For $v \in V$, the out-degree of v is given by

$$\text{outdeg}(v) = |\{w : v \to w\}|,$$

where w is a vertice of V. By definition, this equals the number of $+1$'s in row v of A. Alternatively, the in-degree of v is given by

$$\text{indeg}(v) = |\{u : u \to v\}|$$

and equals the number of -1's in row v of A. It follows that if there exists a graph isomorphism $\varphi : D \to H$, then $\text{outdeg}_D(v) = \text{outdeg}_H(\varphi(v))$ and $\text{indeg}_D(v) = \text{indeg}_H(\varphi(v))$ for every $v \in V$.

For an oriented graph G and $v \in V$, we define $P_G(v)$ to be

$$P_G(v) = (z_1, z_2, z_3), \quad \begin{cases} z_1 = \text{outdeg}(v), \\ z_2 = \text{sorted list of outdeg}(y) \text{ over } y \in N^+(v), \\ z_3 = \text{sorted list of outdeg}(z) \text{ over } z \in N^-(v). \end{cases}$$

Given that an isomorphism preserves outdegrees of vertices and of their neighbours, the multiset $\{P_G(v) : v \in V\}$ would be invariant. We thus continue by computing $P_D(v)$ explicitly for $v = 0$.

A straightforward computation on the first row of A corresponding to vertex v_0 gives $\text{outdeg}_D(0) = 4$ with directed edges $0 \to \{5,7,8,10\}$. The out-neighbours $\{5,7,8,10\}$ then have out-degrees $(6,6,3,6)$ and the in-neighbours $\{1,2,4,6,9,11\}$ have outdegrees $(5,4,5,4,4,7)$. This yields

$$P_D(0) = (4,\ (3,6,6,6),\ (4,4,4,5,5,7)).$$

Next, by a direct check from the matrix C, we have that the vertices of H with outdegree 4 are exactly $2,4,6,9$ with

$$P_H(2) = (4,\ (4,4,6,7),\ (3,4,5,5,6,6)),$$

$$P_H(4) = (4,\ (3,4,6,6),\ (4,4,5,5,6,7)),$$

$$P_H(6) = (4,\ (3,4,5,6),\ (4,5,6,6,6,7)),$$

$$P_H(9) = (4,\ (3,4,5,6),\ (4,5,6,6,6,7)).$$

For example, we derive the $P_H(2)$ above since vertex 2 in H has out-neighbours $\{1,6,9,11\}$ with respective out-degrees $(6,4,4,7)$, and in-neighbours $\{0,3,4,7,8,10\}$ with respective out-degrees $(5,5,4,6,3,6)$. However, this means that $P_D(0) \neq P_H(v)$ for $v = 2,4,6,9$. Therefore, we conclude that D and H are not

Human + AI

isomorphic.

CONCLUSION. We have shown the existence of a pair of oriented graphs D and H with $n = 12$ vertices that have the same switching deck, but are not isomorphic. This proves the existence of switching, non-reconstructible oriented graphs on twelve vertices.

FURTHER QUESTIONS

Human

To motivate further study on this topic, we conclude with the following questions of interest:

Q1: Under what conditions determine whether a digraph is switching, non-reconstructible?

In [MS14], the authors performed an exhaustive search on all 12-vertex tournaments, all orientations of graphs with maximum degree at most 3, and all orientations of quartic graphs, and found them all to be determined by their switching decks. Additionally, Figure 12 in [MS14] provides an explicit example of a pair of digraphs on 12 vertices that are determined by their switching deck, but the digraphs contain 2-cycles and are not oriented graphs. This leads to the following question:

Q2: Are switching, non-reconstructible digraphs restricted to n-vertex oriented graphs for $n \geq 12$? If so, what kinds of oriented graphs can admit this property and for what values of n?

References

[BM11] J. A. Bondy and F. Mercier. Switching reconstruction of digraphs. *J. Graph Theory*, 67:332–348, 2011.

[Kel57] Paul J. Kelly. A congruence theorem for trees. *Pacific J. Math.*, 7:961–968, 1957.

[MS14] Brendan D. McKay and Pascal Schweitzer. Switching reconstruction of digraphs. *J. Graph Theory*, 76(4):279–296, 2014.

[Sch25] Johannes Schmitt. Extremal descendant integrals on moduli spaces of curves: An inequality discovered and proved in collaboration with ai. *arXiv preprint arXiv:2512.14575*, 2025.

[Sta85] R. P. Stanley. Reconstruction from vertex-switching. *J. Combin. Theory Ser. B*, 38:132–138, 1985.

[Ula60] Stanislaw M. Ulam. *A Collection of Mathematical Problems*. Number 8 in Interscience Tracts in Pure and Applied Mathematics. Interscience Publishers, New York, 1960.