

# Sampling networks of ecological interactions

Pedro Jordano<sup>\*a</sup>

<sup>a</sup>Integrative Ecology Group, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (EBD-CSIC), Avenida Americo Vespucio s/n, E-41092 Sevilla, Spain

In 2nd review, Functional Ecology **FE-2015-00758.R2**

August 30, 2016

## Summary

1. Sampling ecological interactions presents similar challenges, problems, potential biases, and constraints as sampling individuals and species in biodiversity inventories. Robust estimates of the actual number of interactions (links) within diversified ecological networks require adequate sampling effort that needs to be explicitly gauged. Yet we still lack a sampling theory explicitly focusing on ecological interactions.
2. While the complete inventory of interactions is likely impossible, a robust characterization of its main patterns and metrics is probably realistic. We

---

<sup>\*</sup>jordano@ebd.csic.es

must acknowledge that a sizeable fraction of the maximum number of interactions  $I_{max}$  among, say,  $A$  animal species and  $P$  plant species (i.e.,  $I_{max} = AP$ ) is impossible to record due to forbidden links, i.e., life-history restrictions. Thus, the number of observed interactions  $I$  in robustly sampled networks is typically  $I \ll I_{max}$ , resulting in sparse interaction matrices with low connectance.

3. Here I provide a review and outline a conceptual framework for interaction sampling by building an explicit analogue to individuals and species sampling, thus extending diversity-monitoring approaches to the characterization of complex networks of ecological interactions. Contrary to species inventories, a sizable fraction of non-observed pairwise interactions cannot be sampled, due to biological constraints that forbid their occurrence.
4. Reasons for forbidden links are multiple but mainly stem from spatial and temporal uncoupling, size mismatches, and intrinsically low probabilities of interspecific encounter for most potential interactions of partner species. Adequately assessing the completeness of a network of ecological interactions thus needs knowledge of the natural history details embedded, so that forbidden links can be accounted for as a portion of the unobserved links when addressing sampling effort.
5. Recent implementations of inference methods for unobserved species or for individual-based data can be combined with the assessment of forbidden links. This can help in estimating their relative importance, simply by the difference between the asymptotic estimate of interaction richness *in a robustly-sampled* assemblage and the maximum richness  $I_{max}$  of interactions.

This is crucial to assess the rapid and devastating effects of defaunation-driven loss of key ecological interactions and the services they provide and the analogous losses related to interaction gains due to invasive species and biotic homogenization.

## Keywords

complex networks, food webs, frugivory, mutualism, plant-animal interactions, pollination, seed dispersal

## Introduction

Biodiversity sampling is a labour-intensive activity, and sampling is often not sufficient to detect all or even most of the species present in an assemblage. Gotelli & Colwell (2011).

1 Biodiversity species assessment aims at sampling individuals in collections and  
2 determining the number of species represented. Given that, by definition, samples  
3 are incomplete, these collections do not enumerate the species actually present.  
4 The ecological literature dealing with robust estimators of species richness and di-  
5 versity in collections of individuals is immense, and a number of useful approaches  
6 have been used to obtain such estimates (Magurran, 1988; Gotelli & Colwell, 2001;  
7 Colwell *et al.*, 2004; Hortal *et al.*, 2006; Colwell, 2009; Gotelli & Colwell, 2011; Chao  
8 *et al.*, 2014). Recent effort has been also focused at defining essential biodiversity  
9 variables (EBV) (Pereira *et al.*, 2013) that can be sampled and measured repeat-

edly to complement biodiversity estimates. Yet sampling species or taxa-specific EBVs is just probing a single component of biodiversity; interactions among species are another fundamental component, one that supports the existence, but in some cases also the extinction, of species. For example, the extinction of interactions represents a dramatic loss of biodiversity because it entails the loss of fundamental ecological functions (Valiente-Banuet *et al.*, 2014). This missed component of biodiversity loss, the extinction of ecological interactions, very often accompanies, or even precedes, species disappearance. Interactions among species are thus a key component of biodiversity and here I aim to show that most problems associated with sampling interactions in natural communities relate to, and are even worse than, problems associated with sampling species diversity. I consider pairwise interactions among species at the habitat level, in the context of alpha diversity and the estimation of local interaction richness from sampling data (Chao *et al.*, 2014). In the first part I provide a succinct overview of previous work addressing sampling issues for ecological interaction networks. In the second part, I discuss specific rationales for sampling the biodiversity of ecological interactions. Finally, I provide a short overview of asymptotic diversity estimates (Gotelli & Colwell, 2001), and a discussion of its application to interaction sampling. Most of the examples come from the analysis of plant-animal interaction networks, yet are applicable to other types of interspecific interactions.

Interactions can be a much better indicator of the richness and diversity of ecosystem functions than a simple list of taxa and their abundances and/or related biodiversity indicator variables (EBVs) (Memmott *et al.*, 2006; Valiente-Banuet *et al.*, 2014). Thus, sampling interactions should be a central issue when identifying and diagnosing ecosystem services (e.g., pollination, seeding by frugivores,

etc.). Fortunately, the whole battery of biodiversity-related tools used by ecologists to sample biodiversity (species, *sensu stricto*) can be extended and applied to the sampling of interactions (see Table 2 in Colwell *et al.*, 2004). Monitoring interactions is a type of biodiversity sampling and is subject to similar methodological shortcomings, especially under-sampling (Jordano, 1987; Jordano *et al.*, 2009; Vázquez *et al.*, 2009; Dorado *et al.*, 2011; Rivera-Hutinel *et al.*, 2012). We are interested in having a complete list of all the pairwise interactions among species (e.g., all the distinct, species-species interactions, or links, among the pollinators and flowering plants) that do actually exist in a given community. Sampling these interactions thus entails exactly the same problems, limitations, constraints, and potential biases as sampling individual organisms and species diversity. As Mao & Colwell (2005) put it, these are the workings of Preston’s demon, the moving “veil line” (Preston, 1948) between the detected and the undetected interactions as sample size increases.

Early efforts to recognize and solve sampling problems in analyses of interactions stem from research on food webs and to determine how undersampling biases food web metrics (Martinez, 1991; Cohen *et al.*, 1993; Wells & O’Hara, 2012, among others). In addition, the myriad of classic natural history studies documenting animal diets, host-pathogen infection records, plant herbivory records, etc., represent efforts to document interactions occurring in nature. All of them share the problem of sampling incompleteness influencing the patterns and metrics reported. Yet, despite the early recognition that incomplete sampling may seriously bias the analysis of ecological networks (Jordano, 1987), only recent studies have explicitly acknowledged it and attempted to determine its influence (Ollerton & Cranmer, 2002; Nielsen & Bascompte, 2007; Vázquez *et al.*, 2009; Gibson *et al.*,

2011; Olesen *et al.*, 2011; Chacoff *et al.*, 2012; Rivera-Hutinel *et al.*, 2012; Olito &  
 Fox, 2014; Bascompte & Jordano, 2014; Vizentin-Bugoni *et al.*, 2014, 2016; Frund  
*et al.*, 2016). The sampling approaches have been extended to predict patterns of  
 coextinctions in interaction assemblages (e.g., hosts-parasites) (Colwell *et al.*, 2012).  
 Most empirical studies provide no indication of sampling effort, implicitly assuming  
 that the reported network patterns and metrics are robust. Yet recent evidences  
 point out that number of partner species detected, number of actual links, and  
 some aggregate statistics describing network patterns, are prone to sampling bias  
 (Nielsen & Bascompte, 2007; Dorado *et al.*, 2011; Olesen *et al.*, 2011; Chacoff *et al.*,  
 2012; Rivera-Hutinel *et al.*, 2012; Olito & Fox, 2014; Frund *et al.*, 2016). Most of  
 these evidences, however, come either from simulation studies (Frund *et al.*, 2016)  
 or from relatively species-poor assemblages. Most certainly, sampling limitations  
 pervade biodiversity inventories and we might rightly expect that frequent inter-  
 actions may be over-represented and rare interactions may be missed entirely in  
 studies of mega-diverse assemblages (Bascompte & Jordano, 2014); but, to what  
 extent?

## Sampling interactions: methods

When we sample interactions in the field we record the presence of two species that  
 interact in some way. For example, Snow and Snow(1988) recorded an interaction  
 whenever they saw a bird “touching” a fruit on a plant. We observe and record  
 feeding observations, visitation, occupancy, presence in pollen loads or in fecal  
 samples, etc., of *individual* animals or plants and accumulate pairwise interactions,  
 i.e., lists of species partners and the frequencies with which we observe them. We

83 assume that the matrix (species numbers) is predefined (i.e., all species interacting  
84 are well documented).

85 Most types of ecological interactions can be illustrated with bipartite graphs,  
86 with two or more distinct groups of interacting partners (Bascompte & Jordano,  
87 2014); for illustration purposes I'll focus more specifically on plant-animal inter-  
88 actions. Sampling interactions requires filling the cells of an interaction matrix  
89 with data. The matrix,  $\Delta = AP$  (the adjacency matrix for the graph representa-  
90 tion of the network), is a 2D inventory of the interactions among, say,  $A$  animal  
91 species (rows) and  $P$  plant species (columns) (Jordano, 1987; Bascompte & Jor-  
92 dano, 2014). The matrix entries illustrate the values of the pairwise interactions  
93 visualized in the  $\Delta$  matrix, and can be 0 or 1, for presence-absence of a given  
94 pairwise interaction, or take a quantitative weight  $w_{ji}$  to represent the interaction  
95 intensity or unidirectional effect of species  $j$  on species  $i$  (Bascompte & Jordano,  
96 2014; Vázquez *et al.*, 2015). The outcomes of most ecological interactions are  
97 dependent on frequency of encounters (e.g., visit rate of pollinators, number of  
98 records of ant defenders, frequency of seeds in fecal samples). Thus, a frequently  
99 used proxy for interaction intensities  $w_{ji}$  is just how frequent new interspecific  
100 encounters are, whether or not appropriately weighted to estimate interaction ef-  
101 fectiveness (Vázquez *et al.*, 2005).

102 We need to define two basic steps in the sampling of interactions: 1) which  
103 type of interactions we sample; and 2) which type of record we get to document  
104 the existence of an interaction. In step #1 we need to take into account whether  
105 we are sampling the whole community of interactor species (all the animal and  
106 plant species) or just a subset of them, i.e., a sub matrix  $\Delta_{m,n}$  of  $m < A$  animal  
107 species and  $n < P$  plant species of the adjacency matrix  $\Delta_{AP}$ . Subsets can be:

108 a) all the potential plants interacting with a subset of the animals (Fig. 1a); b)  
 109 all the potential animal species interacting with a subset of the plant species (Fig.  
 110 1b); c) a subset of all the potential animal species interacting with a subset of all  
 111 the plant species (Fig. 1c). While some discussion has considered how to establish  
 112 the limits of what represents a network (Strogatz, 2001) (in analogy to discussion  
 113 on food-web limits; Cohen, 1978), it must be noted that situations a-c in Fig. 1  
 114 do not represent complete interaction networks. Subnet sampling is generalized  
 115 in studies of biological networks (e.g., protein interactions, gene regulation), yet  
 116 it is important to recognize that most properties of subnetworks (even random  
 117 subsamples) do not represent properties of whole networks (Stumpf *et al.*, 2005).

118 In step #2 above we face the problem of the type of record we take to sample  
 119 interactions. This is important because it defines whether we approach the problem  
 120 of filling up the interaction matrix in a “zoo-centric” way or in a “phyto-centric”  
 121 way. Zoo-centric studies directly sample animal activity and document the plants  
 122 ‘touched’ by the animal. For example, analysis of pollen samples recovered from the  
 123 body of pollinators, analysis of fecal samples of frugivores, radio-tracking data, etc.  
 124 Phyto-centric studies take samples of focal individual plant species and document  
 125 which animals ‘arrive’ or ‘touch’ the plants. Examples include focal watches of  
 126 fruiting or flowering plants to record visitation by animals, raising insect herbivores  
 127 from seed samples, identifying herbivory marks in samples of leaves, etc.

128 Most recent analyses of plant-animal interaction networks are phyto-centric;  
 129 just 3.5% of available plant-pollinator ( $N = 58$ ) or 36.6% plant-frugivore ( $N = 22$ )  
 130 interaction datasets are zoo-centric (see Schleuning *et al.*, 2012). Moreover, most  
 131 available datasets on host-parasite (parasitoid) or plant-herbivore interactions are  
 132 “host-centric” or phyto-centric (e.g., Thébault & Fontaine, 2010; Morris *et al.*,



2013; Eklöf *et al.*, 2013). This may be related to a variety of causes, like preferred methodologies by researchers working with a particular group or system, logistic limitations, or inherent taxonomic focus of the research questions. A likely result of phyto-centric sampling would be adjacency matrices with large  $A : P$  ratios. In contrast, zoo-centric samplings might be prone to detect plants from outside the habitat, complicating the definition of network boundaries. In any case we don't have a clear view of the potential biases that taxa-focused sampling may generate in observed network patterns, for example by generating consistently asymmetric interaction matrices (Dormann *et al.*, 2009).

Reasonably complete analyses of interaction networks can be obtained when combining both phyto-centric and zoo-centric sampling. For example, Bosch *et al.* (2009) showed that the addition of pollen load data on top of focal-plant sampling of pollinators unveiled a significant number of interactions, resulting in important network structural changes. Olesen *et al.* (2011) identified pollen loads on sampled insects and added the new links to an observation-based visitation matrix, with an extra 5% of links representing the estimated number of missing links in the pollination network. The overlap between observational and pollen-load recorded links was only 33%, underscoring the value of combining methodological approaches. Zoo-centric sampling has recently been extended with the use of DNA-barcoding, for example with plant-herbivore (Jurado-Rivera *et al.*, 2009), host-parasitoid (Wirta *et al.*, 2014; Evans *et al.*, 2016), and plant-frugivore interactions (González-Varo *et al.*, 2014). For mutualistic networks we would expect that zoo-centric sampling could help unveiling interactions of the animals with rare plant species or for relatively common plants species which are difficult to sample by direct observation. Future methodological work may provide signifi-

cant advances showing how mixing different sampling strategies strengthens the completeness of network data. These mixed strategies may combine, for instance, timed watches at focal plants, spot censuses along walked transects, pollen load or seed contents analyses, monitoring with camera traps, and DNA barcoding records. However, there are no tested protocols and/or sampling designs for ecological interaction studies to suggest an optimum combination of approaches. Ideally, pilot studies would provide adequate information for each specific study setting.

## 165 Sampling interactions: rationale

166 The number of distinct pairwise interactions that we can record in a landscape (an area of relatively homogeneous vegetation) is equivalent to the number of distinct classes in which we can classify the recorded encounters among *individuals* of two different species. Yet, individual-based interaction networks have been only recently studied (Dupont *et al.*, 2011; Wells & O’Hara, 2012). The most usual approach has been to pool individual-based interaction data into species-based summaries, an approach that ignores the fact that only a fraction of individuals may actually interact given a per capita interaction effect (Wells & O’Hara, 2012). Wells & O’Hara (2012) illustrate the pros and cons of the approach. We walk in the forest and see a blackbird  $Tm$  picking an ivy  $Hh$  fruit and ingesting it: we have a record for  $Tm - Hh$  interaction. We keep advancing and record again a blackbird feeding on hawthorn  $Cm$  fruits so we record a  $Tm - Cm$  interaction; as we advance we encounter another ivy plant and record a blackcap swallowing a fruit so we now have a new  $Sa - Hh$  interaction, and so on. At the end we have a series of classes (e.g.,  $Sa - Hh$ ,  $Tm - Hh$ ,  $Tm - Cm$ , etc.), along with their

observed frequencies.

We get a vector  $c = [c_1 \dots c_n]'$  where  $c_j$  is the number of classes represented  $j$  times in our sampling:  $c_1$  is the number of singletons (interactions recorded once),  $c_2$  is the number of twin pairs (interactions with just two records),  $c_3$  the number of triplets, etc. The problem thus turns to be estimating the number of distinct classes  $C$  from the vector of  $c_j$  values and the frequency of unobserved interactions (see “The real missing links” below).

More specifically, we usually obtain a type of reference sample (Chao *et al.*, 2014) for interactions: a series of repeated samples (e.g., observation days, 1h watches, etc.) with quantitative information, i.e., recording the number of instances of each interaction type on each day. This replicated abundance data, can be treated in three ways: 1) Abundance data within replicates: the counts of interactions, separately for each day; 2) Pooled abundance data: the counts of interactions, summed over all days (the most usual approach); and 3) Replicated incidence data: the number of days on which we recorded each interaction. Assuming a reasonable number of replicates, replicated incidence data is considered to be the most robust statistically, as it takes account of heterogeneity among days (Colwell *et al.*, 2004, 2012; Chao *et al.*, 2014). Thus, both presence-absence and weighted information on interactions can be accommodated for this purpose.

## The species assemblage

When we consider an observed and recorded sample of interactions on a particular assemblage of  $A_{obs}$  and  $P_{obs}$  species (or a set of replicated samples) as a reference sample (Chao *et al.*, 2014) we may have three sources of undersampling error.

204 These sources are ignored if we treat the reference sample as a true representation  
 205 of the interactions in a well-defined assemblage: 1) some animal species are actually  
 206 present but not observed (zero abundance or incidence in the interactions in the  
 207 reference sample),  $A_0$ ; 2) some plant species are actually present but not observed  
 208 (zero abundance or incidence in the interactions in the reference sample),  $P_0$ ; 3)  
 209 some unobserved links (the zeroes in the adjacency matrix,  $UL$ ) may actually  
 210 occur but not recorded. Thus a first problem is determining if  $A_{obs}$  and  $P_{obs}$  truly  
 211 represent the actual species richness interacting in the assemblage. To this end  
 212 we might use the replicated reference samples to estimate the true number of  
 213 interacting animal  $A_{est}$  and plant  $P_{est}$  species as in traditional diversity estimation  
 214 analysis (Chao *et al.*, 2014). If there are no uniques (species seen on only one day),  
 215 then  $A_0$  and  $P_0$  will be zero (based on the Chao2 formula), and we have  $A_{obs}$  and  
 216  $P_{obs}$  as robust estimates of the actual species richness of the assemblage. If  $A_0$   
 217 and  $P_0$  are not zero they estimate the minimum number of undetected animal and  
 218 plant species that can be expected with a sufficiently large number of replicates,  
 219 taken from the same assemblage/locality by the same methods in the same time  
 220 period. We can use extrapolation methods (Colwell *et al.*, 2012) to estimate how  
 221 many additional replicate surveys it would take to reach a specified proportion  $g$   
 222 of  $A_{est}$  and  $P_{est}$ .

## 223 The interactions

224 We are then faced with assessing the sampling of interactions  $I$ . Table 1 summa-  
 225 rizes the main components and targets for estimation of interaction richness. In  
 226 contrast with traditional species diversity estimates, sampling networks has the

paradox that despite the potentially interacting species being present in the sampled assemblage (i.e., included in the  $A_{obs}$  and  $P_{obs}$  species lists), some of their pairwise interactions are impossible to record. The reason is forbidden links. Independently of whether we sample full communities or subset communities we face a problem: some of the interactions that we can visualize in the empty adjacency matrix  $\Delta$  will simply not occur. With a total of  $A_{obs}P_{obs}$  “potential” interactions (eventually augmented to  $A_{est}P_{est}$  in case we have undetected species), a fraction of them are impossible to record, because they are forbidden (Jordano *et al.*, 2003; Olesen *et al.*, 2011).

Our goal is to estimate the true number of non-null  $AP$  interactions, including interactions that actually occur but have not been observed ( $I_0$ ) from the replicated incidence frequencies of interaction types:  $I_{est} = I_{obs} + I_0$ . Note that  $I_0$  estimates the minimum number of undetected plant-animal interactions that can be expected with a sufficiently large number of replicates, taken from the same assemblage/locality by the same methods in the same time period. Therefore we have two types of non-observed links:  $UL^*$  and  $UL$ , corresponding to the real assemblage species richness and to the observed assemblage species richness, respectively (Table 1).

Forbidden links are non-occurrences of pairwise interactions that can be accounted for by biological constraints, such as spatio-temporal uncoupling (Jordano, 1987), size or reward mismatching, foraging constraints (e.g., accessibility) (Moré *et al.*, 2012), and physiological-biochemical constraints (Jordano, 1987). We still have very little information about the frequency of forbidden links in natural communities (Jordano *et al.*, 2003; Stang *et al.*, 2009; Vázquez *et al.*, 2009; Olesen *et al.*, 2011; Ibanez, 2012; Maruyama *et al.*, 2014; Vizentin-Bugoni *et al.*, 2014)

(Table 1). Forbidden links are thus represented as structural zeroes in the interaction matrix, i.e., matrix cells that cannot get a non-zero value. Therefore, we need to account for the frequency of these structural zeros in our matrix before proceeding.

Our main problem then turns to estimate the number of true missed links, i.e., those that can't be accounted for by biological constraints and that might suggest undersampling. Thus, the sampling of interactions in nature, as the sampling of species, is a cumulative process. In our analysis, we are not re-sampling individuals, but interactions, so we built interaction-based accumulation curves. We add new, distinct, interactions recorded as we increase sampling effort (Fig. 2, and Supplementary Online Material). We can obtain an Interaction Accumulation Curve (*IAC*) analogous to a Species Curve (*SAC*) (see Supporting Information in the online data availability repository): the observed number of distinct pairwise interactions in a survey or collection as a function of the accumulated number of observations or samples (Colwell, 2009).

## Empirical data on Forbidden Links

Adjacency matrices are frequently sparse, i.e., they are densely populated with zeroes, with a fraction of them being structural (unobservable interactions) (Bascompte & Jordano, 2014). Thus, it would be a serious interpretation error to attribute the sparseness of adjacency matrices for bipartite networks to just the result of undersampling. The actual typology of link types in ecological interaction networks is thus more complex than just the two categories of observed and unobserved interactions (Table 1). Unobserved interactions are represented by

zeroes and belong to two categories. Missing interactions may actually exist but  
 require additional sampling or a variety of methods to be observed. Forbidden  
 links, on the other hand, arise due to biological constraints limiting interactions  
 and remain unobservable in nature, irrespectively of sampling effort (Table 1).  
 Forbidden links *FL* may actually account for a relatively large fraction of unob-  
 served interactions *UL* when sampling taxonomically-restricted subnetworks (e.g.,  
 plant-hummingbird pollination networks) (Table 1). Phenological uncoupling is  
 also prevalent in most networks, and may add up to explain ca. 25-40% of the for-  
 bidden links, especially in highly seasonal habitats, and up to 20% when estimated  
 relative to the total number of unobserved interactions (Table 2). In any case, we  
 might expect that a fraction of the missing links *ML* would be eventually explained  
 by further biological reasons, depending on the knowledge of natural details of the  
 particular systems. Our goal as naturalists would be to reduce the fraction of *UL*  
 which remain as missing links; to this end we might search for additional biological  
 constraints or increase sampling effort. For instance, habitat use patterns by hum-  
 mingbirds in the Arima Valley network (Table 2; Snow & Snow, 1972) impose a  
 marked pattern of microhabitat mismatches causing up to 44.5% of the forbidden  
 links. A myriad of biological causes beyond those included as *FL* in Table 1 may  
 contribute explanations for *UL*: limits of color perception, presence of secondary  
 metabolites in fruit pulp and leaves, toxins and combinations of monosaccharides  
 in nectar, etc. For example, aside from *FL*, some pairwise interactions may sim-  
 ply have an asymptotically-zero probability of interspecific encounter between the  
 partner species, if they are very rare. However, it is surprising that just the limited  
 set of forbidden link types considered in Table 1 explain between 24.6-77.2% of  
 the unobserved links. Notably, the Arima Valley, Santa Virgínia, and Hato Ratón

networks have  $> 60\%$  of the unobserved links explained, which might be related to the fact that they are subnetworks (Arima Valley, Santa Virgínia) or relatively small networks (Hato Ratón). All this means that empirical networks may have sizable fractions of structural zeroes. Ignoring this biological fact may contribute to wrongly inferring undersampling of interactions in real-world assemblages.

To sum up, two elements of inference are required in the analysis of unobserved interactions in ecological interaction networks: first, detailed natural history information on the participant species that allows the inference of biological constraints imposing forbidden links, so that structural zeroes can be identified in the adjacency matrix. Second, a critical analysis of sampling robustness and a robust estimate of the actual fraction of missing links,  $M$ , resulting in a robust estimate of  $I$ . In the next sections we explore these elements of inference, using *IACs* as analogs to *SACs* to assess the robustness of interaction sampling.

## Assessing sampling effort when recording interactions: asymptotic diversity estimates

A plot of the cumulative number of species recorded,  $S_n$ , as a function of some measure of sampling effort (say,  $n$  samples taken) yields the species accumulation curve (SAC) or collector's curve (Colwell & Coddington, 1994). Similarly, interaction accumulation curves (IAC), analogous to SACs (Gotelli & Colwell, 2001; Hortal *et al.*, 2006; Chao, 2005; Colwell, 2013), can be used to assess the robustness of interactions sampling for plant-animal community datasets (Jordano, 1987; Jordano *et al.*, 2009; Olesen *et al.*, 2011; Chacoff *et al.*, 2012).



322 The basic method to estimate sampling effort and explicitly show the analogues  
 323 with rarefaction analysis in biodiversity research is to vectorize the interaction ma-  
 324 trix  $AP$  so that we get a vector of all the potential pairwise interactions ( $I_{max}$ ,  
 325 Table 1) that can occur in the observed assemblage with  $A_{obs}$  animal species and  
 326  $P_{obs}$  plant species. The new “species” we aim to sample are the pairwise interactions  
 327 (Table 3), as previously discussed. In general, if we have  $A = 1...i$ , animal species  
 328 and  $P = 1...j$  plant species (assuming a complete list of species in the assemblage),  
 329 we’ll have a vector of “new” species to sample:  $A_1P_1, A_1P_2, ...A_2P_1, A_2P_2, ...A_iP_j$ .  
 330 We can represent the successive samples where we can potentially get records of  
 331 these interactions in a matrix with the vectorized interaction matrix and columns  
 332 representing the successive samples we take (Table 3). This is simply a vectorized  
 333 version of the interaction matrix  $\Delta$ . This is analogous to a biodiversity sam-  
 334 pling matrix with species as rows and sampling units (e.g., quadrats) as columns  
 335 (Jordano *et al.*, 2009). The package *EstimateS* (Colwell, 2013) includes a com-  
 336 plete set of functions for estimating the mean IAC and its unconditional standard  
 337 deviation from random permutations of the data, or subsampling without replace-  
 338 ment (Gotelli & Colwell, 2001); it further reports asymptotic estimators for the  
 339 expected number of distinct pairwise interactions included in a given reference  
 340 sample of interaction records (see also the `specaccum` function in library `vegan`  
 341 of the R Package)(R Development Core Team, 2010; Jordano *et al.*, 2009; Olesen  
 342 *et al.*, 2011). In particular, we may take advantage of replicated incidence data,  
 343 as it takes account of heterogeneity among samples (days, censuses, etc.; R.K Col-  
 344 well, pers. comm.) (see also Colwell *et al.*, 2004, 2012; Chao *et al.*, 2014). Future  
 345 theoretical work will be needed to formally assess the similarities and differences  
 346 between the species vs. interactions sampling approaches and developing biologi-

cally meaningful null models of expected interaction richness with added sampling effort.

Diversity-accumulation analysis (Magurran, 1988; Hortal *et al.*, 2006) comes up immediately with this type of dataset. This procedure plots the accumulation curve for the expected number of distinct pairwise interactions recorded with increasing sampling effort (Jordano *et al.*, 2009; Olesen *et al.*, 2011). Asymptotic estimates of interaction richness and its associated standard errors and confidence intervals can thus be obtained (Hortal *et al.*, 2006) (see Table 4 and Supplementary Online Material). The characteristic feature of interaction datasets is that, due to forbidden links, a number of pairwise interactions among the  $I_{max}$  number specified in the  $\Delta$  adjacency matrix cannot be recorded, irrespective of sampling effort.

We may expect undersampling specially in moderate to large sized networks with multiple modules (i.e., species subsets requiring different sampling strategies) (Jordano, 1987; Olesen *et al.*, 2011; Chacoff *et al.*, 2012); adequate sampling may be feasible when interaction subwebs are studied (Olesen *et al.*, 2011; Vizentin-Bugoni *et al.*, 2014), typically with more homogeneous subsets of species (e.g., bumblebee-pollinated flowers).

Mixture models incorporating detectabilities have been proposed to effectively account for rare species (Mao & Colwell, 2005). In an analogous line, mixture models could be extended to samples of pairwise interactions, also with specific detectability values. These detection rate/odds could be variable among groups of interactions, depending on their specific detectability. For example, detectability of flower-pollinator interactions involving bumblebees could have a higher detectability than flower-pollinator pairwise interactions involving, say, nitidulid beetles.

372 These more homogeneous groupings of pairwise interactions within a network de-  
 373 fine modules (Bascompte & Jordano, 2014), so we might expect that interactions  
 374 of a given module (e.g., plants and their hummingbird pollinators; Fig. 1a) may  
 375 share similar detectability values, in an analogous way to species groups receiving  
 376 homogeneous detectability values in mixture models (Mao & Colwell, 2005). In  
 377 its simplest form, this would result in a sample with multiple pairwise interactions  
 378 detected, in which the number of interaction events recorded for each distinct  
 379 interaction found in the sample is recorded (i.e., a column vector in Table 3, cor-  
 380 responding to, say, a sampling day). The number of interactions recorded for the  
 381  $i_{th}$  pairwise interaction (i.e.,  $A_i P_j$  in Table 3),  $Y_i$  could be treated as a Poisson  
 382 random variable with a mean parameter  $\lambda_i$ , its detection rate. Mixture models  
 383 (Mao & Colwell, 2005) include estimates for abundance-based data (their analogs  
 384 in interaction sampling would be weighted data), where  $Y_i$  is a Poisson random  
 385 variable with detection rate  $\lambda_i$ . This is combined with the incidence-based model,  
 386 where  $Y_i$  is a binomial random variable (their analogous in interaction sampling  
 387 would be presence/absence records of interactions) with detection odds  $\lambda_i$ . Let  
 388  $T$  be the number of samples in an incidence-based data set. A Poisson/binomial  
 389 density can be written as (Mao & Colwell, 2005):

$$g(y; \lambda) = \begin{cases} \frac{\lambda^y}{y! e^\lambda} & [1] \\ \binom{T}{y} \frac{\lambda^y}{(1+\lambda)^T} & [2] \end{cases}$$

390 where [1] corresponds to a weighted network, and [2] to a qualitative network.

391 The detection rates  $\lambda_i$  depend on the relative abundances  $\phi_i$  of the interactions,  
 392 the probability of a pairwise interaction being detected when it is present, and the

sample size (the number of interactions recorded), which, in turn, is a function of the sampling effort. Unfortunately, no specific sampling model has been developed along these lines for species interactions and their characteristic features. For example, a complication factor might be that interaction abundances,  $\phi_i$ , in real assemblages are a function of the abundances of interacting species that determine interspecific encounter rates; yet they also depend on biological factors that ultimately determine if the interaction occurs when the partner species are present. For example,  $\lambda_i$  should be set to zero for all  $FL$ . In its simplest form,  $\phi_i$  could be estimated from just the product of partner species abundances, an approach recently used as a null model to assess the role of biological constraints in generating forbidden links and explaining interaction patterns (Vizentin-Bugoni *et al.*, 2014). Yet more complex models (e.g., Wells & O'hara 2012; 2016) should incorporate not only interspecific encounter probabilities, but also interaction detectabilities, phenotypic matching and incidence of forbidden links. Mixture models are certainly complex and for most situations of evaluating sampling effort better alternatives include the simpler incidence-based rarefaction and extrapolation (Colwell *et al.*, 2012; Chao *et al.*, 2014).

## The *real* missing links

Given that a fraction of unobserved interactions can be accounted for by forbidden links, what about the remaining missing interactions? We have already discussed that some of these could still be related to unaccounted constraints, and still others would be certainly attributable to insufficient sampling. Would this always be the case? A crucial ecological aspect limiting interactions within multispecific assem-

416 blages of distinct taxonomic relatedness (Fig. 2) is the probability of interspecific  
 417 encounter, i.e., the probability that two individuals of the partner species actually  
 418 encounter each other in nature.

419 Given log-normally distributed abundances of the two species groups, the ex-  
 420 pected probabilities of interspecific encounter ( $PIE$ ) would be simply the product  
 421 of the two lognormal distributions. Thus, we might expect that for very low  $PIE$   
 422 values, pairwise interactions would be either extremely difficult to sample, or sim-  
 423 ply do not occur in nature. Consider the Nava de las Correhuelas interaction  
 424 web (NCH, Table 2, 4), with  $A = 36$ ,  $P = 25$ ,  $I = 181$ , and almost half of the  
 425 unobserved interactions not accounted for by forbidden links, thus  $M = 53.1\%$   
 426 (Jordano *et al.*, 2009). A sizable fraction of these possible but missing links would  
 427 be simply not occurring in nature, most likely due to extremely low  $PIE$ , in fact  
 428 asymptotically zero. Given the vectorized list of pairwise interactions for NCH, I  
 429 computed the  $PIE$  values for each one by multiplying element-wise the two species  
 430 abundance distributions. The  $PIE_{max} = 0.0597$ , being a neutral estimate, based  
 431 on the assumption that interactions occur in proportion to the species-specific local  
 432 abundances. With  $PIE_{median} < 1.4 \cdot 10^{-4}$  we may safely expect (note the quantile  
 433 estimate  $Q_{75\%} = 3.27 \cdot 10^{-4}$ ) that a sizable fraction of these missing interactions  
 434 may not occur according to this neutral expectation (Jordano, 1987; Olesen *et al.*,  
 435 2011) (neutral forbidden links, *sensu* Canard *et al.*, 2012).

436 When we consider the vectorized interaction matrix, enumerating all pairwise  
 437 interactions for the  $AP$  combinations, the expected probabilities of finding a given  
 438 interaction can be estimated with a Good-Turing approximation (Good, 1953).  
 439 The technique, developed by Alan Turing and I.J. Good with applications to lin-  
 440 guistics and word analysis (Gale & Sampson, 1995) has been recently extended in

novel ways for ecological analyses (Chao *et al.*, 2015). In our present context it estimates the probability of recording an interaction of a hitherto unseen pair of partners, given a set of past records of interactions between other species pairs. Let a sample of  $N$  interactions so that  $n_r$  distinct pairwise interactions have exactly  $r$  records. All Good-Turing estimators obtain the underlying frequencies of events as:

$$P(X) = \frac{(N_X + 1)}{T} \left(1 - \frac{E(1)}{T}\right) \quad (1)$$

where  $X$  is the pairwise interaction,  $N_X$  is the number of times interaction  $X$  is recorded,  $T$  is the sample size (number of distinct interactions recorded) and  $E(1)$  is an estimate of how many different interactions were recorded exactly once. Strictly speaking Equation (1) gives the probability that the next interaction type recorded will be  $X$ , after sampling a given assemblage of interacting species. In other words, we scale down the maximum-likelihood estimator  $\frac{n}{T}$  by a factor of  $\frac{1-E(1)}{T}$ . This reduces all the probabilities for interactions we have recorded, and makes room for interactions we haven't seen. If we sum over the interactions we have seen, then the sum of  $P(X)$  is  $1 - \frac{1-E(1)}{T}$ . Because probabilities sum to one, we have the left-over probability of  $P_{new} = \frac{E(1)}{T}$  of seeing something new, where new means that we sample a new pairwise interaction.

## Discussion

Recent work has inferred that most data available for interaction networks are incomplete due to undersampling, resulting in a variety of biased parameters and

461 network patterns (Chacoff *et al.*, 2012). It is important to note, however, that in  
 462 practice, most surveyed networks to date have been subnets of much larger net-  
 463 works. This is also true for protein interaction, gene regulation, and metabolic  
 464 networks, where only a subset of the molecular entities in a cell have been sam-  
 465 pled (Stumpf *et al.*, 2005). Despite recent attempts to document whole ecosystem  
 466 meta-networks (Pocock *et al.*, 2012), it is likely that most ecological interaction  
 467 networks will illustrate just major ecosystem compartments. Due to their high  
 468 generalization, high temporal and spatial turnover, and high complexity of associ-  
 469 ation patterns, adequate sampling of ecological interaction networks is challenging  
 470 and requires extremely large sampling effort. Undersampling of ecological net-  
 471 works may originate from the analysis of assemblage subsets (e.g., taxonomically  
 472 or functionally defined), and/or from logistically-limited sampling effort. It is ex-  
 473 tremely hard to robustly sample the set of biotic interactions even for relatively  
 474 simple, species-poor assemblages; thus, we need to assess how robust is the char-  
 475 acterization of the adjacency matrix  $\Delta$ . Concluding that an ecological network  
 476 dataset is undersampled just by its sparseness would be unrealistic. The reason  
 477 stems from a biological fact: a sizeable fraction of the maximum, potential links  
 478 that can be recorded among two distinct sets of species is simply unobservable,  
 479 irrespective of sampling effort (Jordano, 1987). In addition, sampling effort needs  
 480 to be explicitly gauged because of its potential influence on parameter estimates  
 481 for the network.

482 Missing links are a characteristic feature of all plant-animal interaction net-  
 483 works, and likely pervade other ecological interactions. Important natural history  
 484 details explain a fraction of them, resulting in unrealizable interactions (i.e., for-  
 485 bidden interactions) that define structural zeroes in the interaction matrices and

contribute to their extreme sparseness. Sampling interactions is a way to monitor biodiversity beyond the simple enumeration of component species and to develop efficient and robust inventories of functional interactions. Yet no sampling theory for interactions is available. Focusing just on the realized interactions or treating missing interactions as the expected unique result of sampling bias would miss important components to understand how all sorts of interactions coevolve within complex webs of interdependence among species.

Contrary to species inventories, a sizable fraction of non-observed pairwise interactions cannot be sampled, due to biological constraints that forbid their occurrence. Moreover, recent implementations of inference methods for unobserved species (Chao *et al.*, 2015) or for individual-based data (Wells & O’Hara, 2012) can be combined with the forbidden link approach. They do not account either for the existence of these ecological constraints, but can help in estimating their relative importance, simply by the difference between the asymptotic estimate of interaction richness *in a robustly-sampled* assemblage and the maximum richness  $I_{max}$  of interactions.

Ecological interactions provide the wireframe supporting the lives of species, and they also embed crucial ecosystem functions which are fundamental for supporting the Earth system. We still have a limited knowledge of the biodiversity of ecological interactions, and they are being lost (extinct) at a very fast pace, frequently preceding species extinctions (Valiente-Banuet *et al.*, 2014). We urgently need robust techniques to assess the completeness of ecological interactions networks because this knowledge will allow the identification of the minimal components of their ecological complexity that need to be restored to rebuild functional ecosystems after perturbations.



## Acknowledgements

I am indebted to Jens M. Olesen, Alfredo Valido, Jordi Bascompte, Thomas Lewinshon, John N. Thompson, Nick Gotelli, Carsten Dormann, and Paulo R. Guimaraes Jr. for useful and thoughtful discussion at different stages of this manuscript. Jeferson Vizentin-Bugoni kindly helped with the Sta Virginia data. Jens M. Olesen kindly made available the Grundvad dataset; together with Robert K. Colwell, Néstor Pérez-Méndez, JuanPe González-Varo, and Paco Rodríguez provided most useful comments to a final version of the ms. Robert Colwell shared a number of crucial suggestions that clarified my vision of sampling ecological interactions, and the final manuscript was greatly improved with comments from three anonymous reviewers. The study was supported by a Junta de Andalucía Excellence Grant (RNM-5731), as well as a Severo Ochoa Excellence Award from the Ministerio de Economía y Competitividad (SEV-2012-0262). The Agencia de Medio Ambiente, Junta de Andalucía, provided generous facilities that made possible my long-term field work in different natural parks.

## Data accessibility

This review does not use new raw data, but includes some re-analyses of previously published material. All the original data supporting the paper, R code, supplementary figures, and summaries of analytical protocols is available at the author's GitHub repository ([https://github.com/pedroj/MS\\_Network-Sampling](https://github.com/pedroj/MS_Network-Sampling)), with DOI: 10.5281/zenodo.29437.

## References

- Bartomeus, I., Gravel, D., Tylianakis, J., Aizen, M., Dickie, I. & Bernard-Verdier, M. (2016) A common framework for identifying linkage rules across different types of interactions. *Functional Ecology* **00**, 0000–0000.
- Bascompte, J. & Jordano, P. (2014) *Mutualistic networks*. Monographs in Population Biology, No. 53, Princeton University Press, Princeton, NJ.
- Bosch, J., Martín González, A.M., Rodrigo, A. & Navarro, D. (2009) Plant-pollinator networks: adding the pollinator’s perspective. *Ecology Letters* **12**, 409–419.
- Canard, E., Mouquet, N., Marescot, L., Gaston, K.J., Gravel, D. & Mouillot, D. (2012) Emergence of structural patterns in neutral trophic networks. *PLoS ONE* **7**, e38295.
- Chacoff, N.P., Vázquez, D.P., Lomáscolo, S.B., Stevani, E.L., Dorado, J. & Padrón, B. (2012) Evaluating sampling completeness in a desert plant-pollinator network. *Journal of Animal Ecology* **81**, 190–200.
- Chao, A. (2005) Species richness estimation. *Encyclopedia of Statistical Sciences*, pp. 7909–7916, Oxford University Press, New York, USA.
- Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K. & Ellison, A.M. (2014) Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecological Monographs* **84**, 45–67.

- 553 Chao, A., Hsieh, T.C., Chazdon, R.L., Colwell, R.K. & Gotelli, N.J. (2015) Un-  
 554 veiling the species-rank abundance distribution by generalizing the Good-Turing  
 555 sample coverage theory. *Ecology* **96**, 1189–1201.
- 556 Cohen, J.E. (1978) *Food webs and niche space*. Princeton University Press, Prince-  
 557 ton, New Jersey, US.
- 558 Cohen, J.E., Beaver, R.A., Cousins, S.H., DeAngelis, D.L., Goldwasser, L., Heong,  
 559 K.L., Holt, R.D., Kohn, A.J., Lawton, J.H., Martinez, N., O'Malley, R., Page,  
 560 L.M., Patten, B.C., Pimm, S.L., Polis, G., Rejmanek, M., Schoener, T.W.,  
 561 Schoenly, K., Sprules, W.G., Teal, J.M., Ulanowicz, R., Warren, P.H., Wilbur,  
 562 H.M. & Yodzis, P. (1993) Improving Food Webs. *Ecology* **74**, 252–258.
- 563 Colwell, R.K. (2009) Biodiversity: concepts, patterns, and measurement. *The*  
 564 *Princeton Guide to Ecology* (ed. S.A. Levin), pp. 257–263, Princeton Univer-  
 565 sity Press, Princeton.
- 566 Colwell, R.K. (2013) EstimateS: statistical estimation of species richness and  
 567 shared species from samples. Version 9.1.0. User's guide and application.  
 568 [<http://viceroy.eeb.uconn.edu/estimates/>] pp. 1–33.
- 569 Colwell, R.K. & Coddington, J. (1994) Estimating terrestrial biodiversity through  
 570 extrapolation. *Philosophical Transactions of the Royal Society of London Series*  
 571 *B-Biological Sciences* **345**, 101–118.
- 572 Colwell, R.K., Dunn, R.R. & Harris, N.C. (2012) Coextinction and persistence of  
 573 dependent species in a changing world. *Annual Review of Ecology Evolution and*  
 574 *Systematics* **43**, 183–203.

- 575 Colwell, R.K., Mao, C.X. & Chang, J. (2004) Interpolating, extrapolating, and  
576 comparing incidence-based species accumulation curves. *Ecology* **85**, 2717–2727.
- 577 Dorado, J., Vázquez, D.P., Stevani, E.L. & Chacoff, N.P. (2011) Rareness and  
578 specialization in plant-pollinator networks. *Ecology* **92**, 19–25.
- 579 Dormann, C.F., Frund, J., Bluthgen, N. & Gruber, B. (2009) Indices, graphs and  
580 null models: Analyzing bipartite ecological networks. *Open Ecology Journal* **2**,  
581 7–24.
- 582 Dupont, Y.L., Trøjelsgaard, K., Hagen, M., Henriksen, M.V., Olesen, J.M., Ped-  
583 ersen, N.M.E. & Kissling, W.D. (2014) Spatial structure of an individual-based  
584 plant-pollinator network. *Oikos* **123**, 1301–1310.
- 585 Dupont, Y.L., Trøjelsgaard, K. & Olesen, J.M. (2011) Scaling down from species  
586 to individuals: a flower-visitation network between individual honeybees and  
587 thistle plants. *Oikos* **120**, 170–177.
- 588 Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N.P.,  
589 Dalsgaard, B., de Sassi, C., Galetti, M., Guimaraes, P.R., Lomáscolo, S.B.,  
590 Martín González, A.M., Pizo, M.A., Rader, R., Rodrigo, A., Tylianakis, J.M.,  
591 Vázquez, D.P. & Allesina, S. (2013) The dimensionality of ecological networks.  
592 *Ecology Letters* **16**, 577–583.
- 593 Evans, D., Kitson, J., Lunt, D., Straw, N. & Pocock, M. (2016) Merging DNA  
594 metabarcoding and ecological network analysis to understand and build resilient  
595 terrestrial ecosystems. *Functional Ecology* **00**, 0000–0000.
- 596 Frund, J., McCann, K.S. & Williams, N.M. (2016) Sampling bias is a challenge

- for quantifying specialization and network structure: lessons from a quantitative niche model. *Oikos* **125**, 502–513.
- Gale, W.A. & Sampson, G. (1995) Good-Turing frequency estimation without tears. *Journal of Quantitative Linguistics* **2**, 217–237.
- Gibson, R.H., Knott, B., Eberlein, T. & Memmott, J. (2011) Sampling method influences the structure of plant–pollinator networks. *Oikos* **120**, 822–831.
- González-Varo, J.P., Arroyo, J.M. & Jordano, P. (2014) Who dispersed the seeds? The use of DNA barcoding in frugivory and seed dispersal studies. *Methods in Ecology and Evolution* **5**, 806–814.
- Good, I.J. (1953) The population frequencies of species and the estimation of population parameters. *Biometrika* **40**, 237–264.
- Gotelli, N. & Colwell, R. (2001) Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* **4**, 379–391.
- Gotelli, N.J. & Colwell, R.K. (2011) Estimating species richness. *Biological Diversity Frontiers in Measurement and Assessment* (eds. A.E. Magurran & B.J. McGill), pp. 39–54, Oxford University Press, Oxford, UK.
- Hortal, J., Borges, P. & Gaspar, C. (2006) Evaluating the performance of species richness estimators: sensitivity to sample grain size. *Journal of Animal Ecology* **75**, 274–287.
- Ibanez, S. (2012) Optimizing size thresholds in a plant–pollinator interaction web:

- 618 towards a mechanistic understanding of ecological networks. *Oecologia* **170**, 233–  
619 242.
- 620 Jordano, P. (1987) Patterns of mutualistic interactions in pollination and seed dis-  
621 persal: connectance, dependence asymmetries, and coevolution. *The American*  
622 *Naturalist* **129**, 657–677.
- 623 Jordano, P., Bascompte, J. & Olesen, J. (2003) Invariant properties in coevolu-  
624 tionary networks of plant-animal interactions. *Ecology Letters* **6**, 69–81.
- 625 Jordano, P., Vázquez, D. & Bascompte, J. (2009) Redes complejas de interac-  
626 ciones planta—animal. *Ecología y evolución de interacciones planta-animal* (eds.  
627 R. Medel, R. Dirzo & R. Zamora), pp. 17–41, Editorial Universitaria, Santiago,  
628 Chile.
- 629 Jurado-Rivera, J.A., Vogler, A.P., Reid, C.A.M., Petitpierre, E. & Gomez-Zurita,  
630 J. (2009) DNA barcoding insect-host plant associations. *Proceedings Of The*  
631 *Royal Society B-Biological Sciences* **276**, 639–648.
- 632 Magurran, A. (1988) *Ecological diversity and its measurement*. Princeton Univer-  
633 sity Press, Princeton, US.
- 634 Mao, C. & Colwell, R.K. (2005) Estimation of species richness: mixture models,  
635 the role of rare species, and inferential challenges. *Ecology* **86**, 1143–1153.
- 636 Martinez, N. (1991) Artifacts or attributes? Effects of resolution on food-web  
637 patterns in Little Rock Lake food web. *Ecological Monographs* **61**, 367–392.
- 638 Maruyama, P.K., Vizentin-Bugoni, J., Oliveira, G.M., Oliveira, P.E. & Dalsgaard,

- 639 B. (2014) Morphological and spatio-temporal mismatches shape a neotropical  
640 savanna plant-hummingbird network. *Biotropica* **46**, 740–747.
- 641 Memmott, J., Alonso, D., Berlow, E., Dobson, A., Dunne, J.A., Solé, R.V. &  
642 Weitz, J.S. (2006) Biodiversity loss and ecological network structure. *Ecological*  
643 *Networks Linking Structure to Dynamics in Food Webs* (eds. M. Pascual & J.A.  
644 Dunne), pp. 325–347, Oxford University Press, Oxford, UK.
- 645 Moré, M., Amorim, F.W., Benitez-Vieyra, S., Medina, A.M., Sazima, M. &  
646 Cocucci, A.A. (2012) Armament imbalances: match and mismatch in plant-  
647 pollinator traits of highly specialized long-spurred orchids. *PLoS ONE* **7**, e41878.
- 648 Morris, R.J., Gripenberg, S., Lewis, O.T. & Roslin, T. (2013) Antagonistic inter-  
649 action networks are structured independently of latitude and host guild. *Ecology*  
650 *Letters* **17**, 340–349.
- 651 Nielsen, A. & Bascompte, J. (2007) Ecological networks, nestedness and sampling  
652 effort. *Journal of Ecology* **95**, 1134–1141–1141.
- 653 Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H. & Jordano, P. (2011)  
654 Missing and forbidden links in mutualistic networks. *Proceedings of the Royal*  
655 *Society B-Biological Sciences* **278**, 725–732.
- 656 Olito, C. & Fox, J.W. (2014) Species traits and abundances predict metrics of  
657 plant-pollinator network structure, but not pairwise interactions. *Oikos* **124**,  
658 428–436.
- 659 Ollerton, J. & Cranmer, L. (2002) Latitudinal trends in plant-pollinator interac-  
660 tions: are tropical plants more specialised? *Oikos* **98**, 340–350.

- 661 Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes,  
662 R.J., Bruford, M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops,  
663 N., Dulloo, E., Faith, D., Freyhof, J., Gregory, R.D., Heip, C., Hoft, R., Hurtt,  
664 G., Jetz, W., Karp, D.S., Mcgeoch, M., Obura, D., Onoda, Y., Pettorelli, N.,  
665 Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S., Turak, E., Walpole, M.  
666 & Wegmann, M. (2013) Essential biodiversity variables. *Science* **339**, 277–278.
- 667 Pocock, M.J.O., Evans, D.M. & Memmott, J. (2012) The robustness and restora-  
668 tion of a network of ecological networks. *Science* **335**, 973–977.
- 669 Preston, F. (1948) The commonness, and rarity, of species. *Ecology* **29**, 254–283.
- 670 R Development Core Team (2010) *R: A language and environment for statis-*  
671 *tical computing*. R Foundation for Statistical Computing. Vienna, Austria.  
672 <http://www.R-project.org>, Vienna, Austria.
- 673 Rivera-Hutinel, A., Bustamante, R.O., Marín, V.H. & Medel, R. (2012) Effects  
674 of sampling completeness on the structure of plant-pollinator networks. *Ecology*  
675 **93**, 1593–1603.
- 676 Schleuning, M., Frund, J., Klein, A.M., Abrahamczyk, S., Alarcón, R., Albrecht,  
677 M., Andersson, G.K.S., Bazarian, S., Böhning-Gaese, K., Bommarco, R., Dals-  
678 gaard, B., Dehling, D.M., Gotlieb, A., Hagen, M., Hickler, T., Holzschuh, A.,  
679 Kaiser-Bunbury, C.N., Kreft, H., Morris, R.J., Sandel, B., Sutherland, W.J.,  
680 Svenning, J.C., Tschardtke, T., Watts, S., Weiner, C.N., Werner, M., Williams,  
681 N.M., Winqvist, C., Dormann, C.F. & Blüthgen, N. (2012) Specialization of  
682 mutualistic interaction networks decreases toward tropical latitudes. *Current*  
683 *Biology* **22**, 1925–1931.



- 684 Snow, B. & Snow, D. (1972) Feeding niches of hummingbirds in a Trinidad valley.  
685 *Journal of Animal Ecology* **41**, 471–485.
- 686 Snow, B. & Snow, D. (1988) *Birds and berries*. Poyser, Calton, UK.
- 687 Stang, M., Klinkhamer, P., Waser, N.M., Stang, I. & van der Meijden, E. (2009)  
688 Size-specific interaction patterns and size matching in a plant-pollinator inter-  
689 action web. *Annals of Botany* **103**, 1459–1469.
- 690 Strogatz, S. (2001) Exploring complex networks. *Nature* **410**, 268–276.
- 691 Stumpf, M.P.H., Wiuf, C. & May, R.M. (2005) Subnets of scale-free networks  
692 are not scale-free: Sampling properties of networks. *Proceedings of the National*  
693 *Academy of Sciences USA* **102**, 4221–4224.
- 694 Thébault, E. & Fontaine, C. (2010) Stability of ecological communities and the  
695 architecture of mutualistic and trophic networks. *Science* **329**, 853–856.
- 696 Valiente-Banuet, A., Aizen, M.A., Alcántara, J.M., Arroyo, J., Cocucci, A.,  
697 Galetti, M., García, M.B., García, D., Gomez, J.M., Jordano, P., Medel, R.,  
698 Navarro, L., Obeso, J.R., Oviedo, R., Ramírez, N., Rey, P.J., Traveset, A.,  
699 Verdú, M. & Zamora, R. (2014) Beyond species loss: the extinction of ecological  
700 interactions in a changing world. *Functional Ecology* **29**, 299–307.
- 701 Vázquez, D., Morris, W. & Jordano, P. (2005) Interaction frequency as a surrogate  
702 for the total effect of animal mutualists on plants. *Ecology Letters* **8**, 1088–1094.
- 703 Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009) Evaluating multiple determi-  
704 nants of the structure of plant-animal mutualistic networks. *Ecology* **90**, 2039–  
705 2046.

- 706 Vázquez, D.P., Ramos-Jiliberto, R., Urbani, P. & Valdovinos, F.S. (2015) A con-  
 707 ceptual framework for studying the strength of plant-animal mutualistic inter-  
 708 actions. *Ecology Letters* **18**, 385–400.
- 709 Vizentin-Bugoni, J., Maruyama, P.K., Debastiani, V.J., Duarte, L.d.S., Dalsgaard,  
 710 B. & Sazima, M. (2016) Influences of sampling effort on detected patterns and  
 711 structuring processes of a neotropical plant–hummingbird network. *Journal of*  
 712 *Animal Ecology* **85**, 262–272.
- 713 Vizentin-Bugoni, J., Maruyama, P.K. & Sazima, M. (2014) Processes entangling  
 714 interactions in communities: forbidden links are more important than abundance  
 715 in a hummingbird-plant network. *Proceedings of the Royal Society B-Biological*  
 716 *Sciences* **281**, 20132397–20132397.
- 717 Wells, K. & O’Hara, R.B. (2012) Species interactions: estimating per-individual  
 718 interaction strength and covariates before simplifying data into per-species eco-  
 719 logical networks. *Methods in Ecology and Evolution* **4**, 1–8.
- 720 Wirta, H.K., Hebert, P.D.N., Kaartinen, R., Prosser, S.W., Várkonyi, G. & Roslin,  
 721 T. (2014) Complementary molecular information changes our perception of food  
 722 web structure. *Proceedings of the National Academy of Sciences USA* **111**, 1885–  
 723 1890.

## Figure captions

**Figure 1.** Sampling ecological interaction networks (e.g., plant-animal interactions) usually focus on different types of subsampling the full network, yielding submatrices  $\Delta[m, n]$  of the full interaction matrix  $\Delta$  with  $A$  and  $P$  animal and plant species. a) all the potential plants interacting with a subset of the animals (e.g., studying just the hummingbird-pollinated flower species in a community); b) all the potential animal species interacting with a subset of the plant species (e.g., studying the frugivore species feeding on figs *Ficus* in a community); and c) sampling a subset of all the potential animal species interacting with a subset of all the plant species (e.g., studying the plant-frugivore interactions of the rainforest understory).

**Figure 2.** Sampling species interactions in natural communities. Suppose an assemblage with  $A = 3$  animal species (red, species 1–3 with three, two, and 1 individuals, respectively) and  $P = 3$  plant species (green, species a-c with three individuals each) (colored balls), sampled with increasing effort in steps 1 to 6 (panels). In Step 1 we record a total of three interactions (black lines) for animal species 1 and plant species a and b, represented as two distinct pairwise interactions:  $1 - a$  and  $1 - b$ . As we advance our sampling (panels 1 to 6, illustrating e.g., additional sampling days) we record new distinct interactions. Note that we actually sample and record interactions among individuals, yet we pool the data across species to get a species by species interaction matrix. Few network analyses have been carried out on individual data (e.g., Dupont *et al.*, 2014). Above and below each panel are the cumulative number of distinct species and interactions

748 sampled, so that panel 6 illustrates the final network.

749

## 750 Figures

Figure 1:





Figure 2:



Jordano - Figure 1

## Table captions

**Table 1.** A taxonomy of link types for ecological interactions (Olesen *et al.* 2011).  $A_{obs}$ , number of animal species;  $P_{obs}$ , number of plant species;  $I_{obs}$ , number of observed links;  $FL$ , number of forbidden links; and  $ML$ , number of missing links. As natural scientists, our ultimate goal is to eliminate  $ML$  from the equation  $FL = AP - I - ML$ , which probably is not feasible given logistic sampling limitations. When we, during our study, estimate  $ML$  to be negligible, we cease observing and estimate  $I$  and  $FL$ .

**Table 2.** Frequencies of different type of forbidden links in natural plant-animal interaction assemblages.  $AP$ , maximum potential links,  $I_{max}$ ;  $I$ , number of observed links;  $UL$ , number of unobserved links;  $FL$ , number of forbidden links;  $FL_P$ , phenology constraints;  $FL_S$ , size restrictions;  $FL_A$ , accessibility constraints;  $FL_O$ , other types of restrictions;  $ML$ , unknown causes (missing links). Relative frequencies (in parentheses) calculated over  $I_{max} = AP$  for  $I$  and  $UL$ ; those for  $FL$ , and  $ML$ , over  $UL$ ; for all forbidden links types, calculated over  $FL$ . Data for Hato Ratón and Nava Correhuelas include interactions with bird and mammal frugivores. References, from left to right: Olesen *et al.* 2008; Olesen & Myrthue unpubl.; Snow & Snow 1972 and Jordano *et al.* 2006; Vizentin-Bugoni *et al.* 2014; Jordano *et al.* 2009; Olesen *et al.* 2011.

**Table 3.** A vectorized interaction matrix.

**Table 4.** Sampling statistics for three plant-animal interaction networks (Olesen

775 *et al.* 2011). Symbols as in Table 1;  $N$ , number of records;  $Chao1$  and  $ACE$  are  
 776 asymptotic estimators for the number of distinct pairwise interactions  $I$  (Hortal  
 777 *et al.* 2006), and their standard errors;  $C$ , sample coverage for rare interactions  
 778 (Chao & Jost 2012). Scaled asymptotic estimators and their confidence intervals  
 779 ( $CI$ ) were calculated by weighting  $Chao1$  and  $ACE$  with the observed frequencies  
 780 of forbidden links. Data for Hato Ratón and Nava Correhuelas include only plant-  
 781 bird interactions.

782

## 783 Tables



Table 1:

Link type	Formulation	Definition
Potential links	$I_{max} = A_{obs}P_{obs}$	Size of observed network matrix, i.e. maximum number of potentially observable interactions; $A_{obs}$ and $P_{obs}$ , numbers of interacting animal and plant species, respectively. These might be below the real numbers of animal and plant species, $A_{est}$ and $P_{est}$ .
Observed links	$I_{obs}$	Total number of observed links in the network given a sufficient sampling effort. Number of ones in the adjacency matrix.
True links	$I_{est}$	Total number of links in the network given a sufficient sampling effort; expected for the augmented $A_{est}P_{est}$ matrix.
Unobserved links	$UL = I_{max} - I_{obs}$	Number of zeroes in the adjacency matrix.
True unobserved links	$UL* = I_{max} - I_{obs}$	Number of zeroes in the augmented adjacency matrix that, eventually, includes unobserved species.
Forbidden links	$FL$	Number of links, which remain unobserved because of linkage constraints, irrespectively of sufficient sampling effort.
Observed Missing links	$ML = A_{obs}P_{obs} - I_{obs} - FL$	Number of links, which may exist in nature but need more sampling effort and/or additional sampling methods to be observed.
True Missing links	$ML* = A_{est}P_{est} - I_{est} - FL$	Number of links, which may exist in nature but need more sampling effort and/or additional sampling methods to be observed. Augments $ML$ for the $A_{est}P_{est}$ matrix.

Table 2:

Link type	Pollination			Seed dispersal		
	Zackenber	Grundvad	Arima Valley	Sta. Virginia	Hato Ratón	Nava Correhuelas
$I_{max}$	1891	646	522	423	320	900
$I$	384 (0.2031)	212 (0.3282)	185 (0.3544)	86 (0.2033)	151 (0.4719)	181 (0.2011)
$UL$	1507 (0.7969)	434 (0.6718)	337 (0.6456)	337 (0.7967)	169 (0.5281)	719 (0.7989)
$FL$	530 (0.3517)	107 (0.2465)	218 (0.6469)	260 (0.7715)	118 (0.6982)	320 (0.4451)
$FL_P$	530 (1.0000)	94 (0.8785)	... (...)	120 (0.4615)	67 (0.5678)	195 (0.6094)
$FL_S$	... (...)	8 (0.0748)	30 (0.1376)	140 (0.5385)	31 (0.2627)	46 (0.1438)
$FL_A$	... (...)	5 (0.0467)	150 (0.6881) <sup>a</sup>	... (...)	20 (0.1695)	79 (0.2469)
$FL_O$	... (...)	... (...)	38 (0.1743) <sup>b</sup>	... (...)	... (...)	... (...)
$ML$	977 (0.6483)	327 (0.7535)	119 (0.3531)	77 (0.2285)	51 (0.3018)	399 (0.5549)

<sup>a</sup>, Lack of accessibility due to habitat uncoupling, i.e., canopy-foraging species vs. understory species.

<sup>b</sup>, Colour restrictions, and reward per flower too small relative to the size of the bird.

Dots indicate no data available for the  $FL$  type.

Table 3:

Interaction	Sample 1	Sample 2	Sample 3	...	Sample $i$
A1 - P1	12	2	0	...	6
A1 - P2	0	0	0	...	1
...	...	...	...	...	...
A5 - P3	5	0	1	...	18
A5 - P4	1	0	1	...	3
...	...	...	...	...	...
A <sub>i</sub> - P <sub>i</sub>	1	0	1	...	2

Table 4:

	Hato Ratón	Nava Correhuelas	Zackenberg
$A$	17	33	65
$P$	16	25	31
$I_{max}$	272	825	1891
$N$	3340	8378	1245
$I$	151	181	268
$Chao1$	$263.1 \pm 70.9$	$231.4 \pm 14.2$	$509.6 \pm 54.7$
$ACE$	$240.3 \pm 8.9$	$241.3 \pm 7.9$	$566.1 \pm 14.8$
$\% \text{ unobserved}^a$	8.33	15.38	47.80

<sup>a</sup>, estimated with library Jade (R Core Development Team 2010, Chao *et al.* 2015)