

Sampling networks of ecological interactions

Pedro Jordano^{*a}

^aIntegrative Ecology Group, Estación Biológica de Doñana, Consejo
Superior de Investigaciones Científicas (EBD-CSIC), Avenida
Americo Vespucio s/n, E-41092 Sevilla, Spain

In review, Functional Ecology, February 4, 2016

Summary

1. Sampling ecological interactions presents similar challenges, problems, potential biases, and constraints as sampling individuals and species in biodiversity inventories. Interactions are just pairwise relationships among individuals of two different species, such as those among plants and their seed dispersers in frugivory interactions or those among plants and their pollinators. Sampling interactions is a fundamental step to build robustly estimated interaction networks, yet few analyses have attempted a formal approach to their sampling protocols.

^{*}jordano@ebd.csic.es

2. Robust estimates of the actual number of interactions (links) within diversified ecological networks require adequate sampling effort that needs to be explicitly gauged. Yet we still lack a sampling theory explicitly focusing on ecological interactions.
3. While the complete inventory of interactions is likely impossible, a robust characterization of its main patterns and metrics is probably realistic. We must acknowledge that a sizable fraction of the maximum number of interactions I_{max} among, say, A animal species and P plant species (i.e., $I_{max} = AP$) is impossible to record due to forbidden links, i.e., life-history restrictions. Thus, the number of observed interactions I in robustly sampled networks is typically $I \ll I_{max}$, resulting in sparse interaction matrices with low connectance.
4. Reasons for forbidden links are multiple but mainly stem from spatial and temporal uncoupling, size mismatches, and intrinsically low probabilities of interspecific encounter for most potential interactions of partner species. Adequately assessing the completeness of a network of ecological interactions thus needs knowledge of the natural history details embedded, so that forbidden links can be accounted for when addressing sampling effort.
5. Here I provide a review and outline a conceptual framework for interaction sampling by building an explicit analogue to individuals and species sampling, thus extending diversity-monitoring approaches to the characterization of complex networks of ecological interactions. Contrary to species inventories, a sizable fraction of non-observed pairwise interactions cannot be sampled, due to biological constraints that forbid their occurrence.

6. Recent implementations of inference methods for unobserved species or for individual-based data can be combined with the assessment of forbidden links. This can help in estimating their relative importance, simply by the difference between the asymptotic estimate of interaction richness *in a robustly-sampled* assemblage and the maximum richness I_{max} of interactions. This is crucial to assess the rapid and devastating effects of defaunation-driven loss of key ecological interactions and the services they provide and the analogous losses related to interaction gains due to invasive species and biotic homogenization.

Keywords

complex networks, food webs, frugivory, mutualism, plant-animal interactions, pollination, seed dispersal

Introduction

Biodiversity sampling is a labour-intensive activity, and sampling is often not sufficient to detect all or even most of the species present in an assemblage. Gotelli & Colwell (2011).

- 1 Biodiversity species assessment aims at sampling individuals in collections and
- 2 determining the number of species represented. Given that, by definition, samples
- 3 are incomplete, these collections do not enumerate the species actually present.
- 4 The ecological literature dealing with robust estimators of species richness and di-

5 versity in collections of individuals is immense, and a number of useful approaches
6 have been used to obtain such estimates (Magurran, 1988; Gotelli & Colwell, 2001;
7 Colwell, Mao & Chang, 2004; Hortal, Borges & Gaspar, 2006; Colwell, 2009; Gotelli
8 & Colwell, 2011; Chao *et al.*, 2014). Recent effort has been also focused at defining
9 essential biodiversity variables (EBV) (Pereira *et al.*, 2013) that can be sampled
10 and measured repeatedly to complement biodiversity estimates. Yet sampling
11 species or taxa-specific EBVs is just probing a single component of biodiversity;
12 interactions among species are another fundamental component, one that supports
13 the existence, but in some cases also the extinction, of species. For example, the
14 extinction of interactions represents a dramatic loss of biodiversity because it en-
15 tails the loss of fundamental ecological functions (Valiente-Banuet *et al.*, 2014).
16 This missed component of biodiversity loss, the extinction of ecological interac-
17 tions, very often accompanies, or even precedes, species disappearance. Interac-
18 tions among species are a key component of biodiversity and here we aim to show
19 that most problems associated with sampling interactions in natural communities
20 relate to, and are even worse than, problems associated with sampling species di-
21 versity. We consider pairwise interactions among species at the habitat level, in
22 the context of alpha diversity and the estimation of local interaction richness from
23 sampling data (Chao *et al.*, 2014). In the first part we provide a succinct overview
24 of previous work addressing sampling issues for ecological interaction networks. In
25 the second part, we discuss specific rationales for sampling the biodiversity of eco-
26 logical interactions. Finally, we provide a short overview of asymptotic diversity
27 estimates (Gotelli & Colwell, 2001), and a discussion of its application to inter-
28 action sampling. Most of the examples come from the analysis of plant-animal
29 interaction networks, yet are applicable to other types of interspecific interactions.

Interactions can be a much better indicator of the richness and diversity of ecosystem functions than a simple list of taxa and their abundances and/or related biodiversity indicator variables (EBVs) (Memmott *et al.*, 2006; Valiente-Banuet *et al.*, 2014). Thus, sampling interactions should be a central issue when identifying and diagnosing ecosystem services (e.g., pollination, seeding by frugivores, etc.). Fortunately, the whole battery of biodiversity-related tools used by ecologists to sample biodiversity (species, *sensu stricto*) can be extended and applied to the sampling of interactions. Analogs are evident between these approaches (see Table 2 in Colwell, Mao & Chang, 2004). Monitoring interactions is a type of biodiversity sampling and is subject to similar methodological shortcomings, especially undersampling (Jordano, 1987; Jordano, Vázquez & Bascompte, 2009; Coddington *et al.*, 2009; Vázquez, Chacoff & Cagnolo, 2009; Dorado *et al.*, 2011; Rivera-Hutinel *et al.*, 2012). For example, when we study mutualistic networks, our goal is to make an inventory of the distinct pairwise interactions that made up the network. We are interested in having a complete list of all the pairwise interactions among species (e.g., all the distinct, species-species interactions, or links, among the pollinators and flowering plants) that do actually exist in a given community. Sampling these interactions thus entails exactly the same problems, limitations, constraints, and potential biases as sampling individual organisms and species diversity. As Mao & Colwell (2005) put it, these are the workings of Preston’s demon, the moving “veil line” (Preston, 1948) between the detected and the undetected interactions as sample size increases.

Early efforts to recognize and solve sampling problems in analyses of interactions stem from research on food webs and to determine how undersampling biases food web metrics (Martinez, 1991; Cohen *et al.*, 1993; Martinez, 1993; Bersier,

55 Banasek-Richter & Cattin, 2002; Brose, Martinez & Williams, 2003; Banasek-
 56 Richter, Cattin & Bersier, 2004; Wells & O'Hara, 2012). In addition, the myriad
 57 of classic natural history studies documenting animal diets, host-pathogen infection
 58 records, plant herbivory records, etc., represent efforts to document interactions
 59 occurring in nature. All of them share the problem of sampling incompleteness in-
 60 fluencing the patterns and metrics reported. Yet, despite the early recognition that
 61 incomplete sampling may seriously bias the analysis of ecological networks (Jor-
 62 dano, 1987), only recent studies have explicitly acknowledged it and attempted to
 63 determine its influence (Ollerton & Cranmer, 2002; Nielsen & Bascompte, 2007;
 64 Vázquez, Chacoff & Cagnolo, 2009; Gibson *et al.*, 2011; Olesen *et al.*, 2011; Chacoff
 65 *et al.*, 2012; Rivera-Hutinel *et al.*, 2012; Olito & Fox, 2014; Bascompte & Jordano,
 66 2014; Vizentin-Bugoni, Maruyama & Sazima, 2014; Vizentin-Bugoni *et al.*, 2016;
 67 Frund, McCann & Williams, 2015). The sampling approaches have been extended
 68 to predict patterns of coextinctions in interaction assemblages (e.g., hosts-parasites)
 69 (Colwell, Dunn & Harris, 2012). Most empirical studies provide no indication of
 70 sampling effort, implicitly assuming that the reported network patterns and met-
 71 rics are robust. Yet recent evidences point out that number of partner species
 72 detected, number of actual links, and some aggregate statistics describing network
 73 patterns, are prone to sampling bias (Nielsen & Bascompte, 2007; Dorado *et al.*,
 74 2011; Olesen *et al.*, 2011; Chacoff *et al.*, 2012; Rivera-Hutinel *et al.*, 2012; Olito &
 75 Fox, 2014; Frund, McCann & Williams, 2015). Most of these evidences, however,
 76 come either from simulation studies (Frund, McCann & Williams, 2015) or from
 77 relatively species-poor assemblages. Most certainly, sampling limitations pervade
 78 biodiversity inventories in tropical areas (Coddington *et al.*, 2009) and we might
 79 rightly expect that frequent interactions may be over-represented and rare inter-

actions may be missed entirely in studies of mega-diverse assemblages (Bascompte & Jordano, 2014); but, to what extent?

Sampling interactions: methods

When we sample interactions in the field we record the presence of two species that interact in some way. For example, Snow and Snow(1988) recorded an interaction whenever they saw a bird “touching” a fruit on a plant. We observe and record feeding observations, visitation, occupancy, presence in pollen loads or in fecal samples, etc., of *individual* animals or plants and accumulate pairwise interactions, i.e., lists of species partners and the frequencies with which we observe them. Therefore, estimating the sampling completeness of pairwise interactions for a whole network, requires some gauging of how the number (richness) of distinct pairwise interactions accumulates as sampling effort is increased) and/or estimating the uncertainty around the missed links (Wells & O’Hara, 2012).

Most types of ecological interactions can be illustrated with bipartite graphs, with two or more distinct groups of interacting partners (Bascompte & Jordano, 2014); for illustration purposes I’ll focus more specifically on plant-animal interactions. Sampling interactions requires filling the cells of an interaction matrix with data. The matrix, $\Delta = AP$ (the adjacency matrix for the graph representation of the network), is a 2D inventory of the interactions among, say, A animal species (rows) and P plant species (columns) (Jordano, 1987; Bascompte & Jordano, 2014). The matrix entries illustrate the values of the pairwise interactions visualized in the Δ matrix, and can be 0 or 1, for presence-absence of a given pairwise interaction, or take a quantitative weight w_{ji} to represent the interaction

intensity or unidirectional effect of species j on species i (Bascompte & Jordano, 2014; Vazquez *et al.*, 2015). The outcomes of most ecological interactions are dependent on frequency of encounters (e.g., visit rate of pollinators, number of records of ant defenders, frequency of seeds in fecal samples). Thus, a frequently used proxy for interaction intensities w_{ji} is just how frequent new interspecific encounters are, whether or not appropriately weighted to estimate interaction effectiveness (Vazquez, Morris & Jordano, 2005).

We need to define two basic steps in the sampling of interactions: 1) which type of interactions we sample; and 2) which type of record we get to document the existence of an interaction. In step #1 we need to take into account whether we are sampling the whole community of interactor species (all the animals, all the plants) or just a subset of them, i.e., a sub matrix $\Delta_{m,n}$ of $m < A$ animal species and $n < P$ plant species of the adjacency matrix Δ_{AP} (i.e., the matrix representation of interactions among the partner species). Subsets can be: a) all the potential plants interacting with a subset of the animals (Fig. 1a); b) all the potential animal species interacting with a subset of the plant species (Fig. 1b); c) a subset of all the potential animal species interacting with a subset of all the plant species (Fig. 1c). While some discussion has considered how to establish the limits of what represents a network (Strogatz, 2001) (in analogy to discussion on food-web limits; Cohen, 1978), it must be noted that situations a-c in Fig. 1 do not represent complete interaction networks. As vividly stated by Cohen *et al.* (1993): “*As more comprehensive, more detailed, more explicit webs become available, smaller, highly aggregated, incompletely described webs may progressively be dropped from analyses of web structure (though such webs may remain useful for other purposes, such as pedagogy)*”. Subnet sampling is generalized in studies of

biological networks (e.g., protein interactions, gene regulation), yet it is important to recognize that most properties of subnetworks (even random subsamples) do not represent properties of whole networks (Stumpf, Wiuf & May, 2005).

In step #2 above we face the problem of the type of record we take to sample interactions. This is important because it defines whether we approach the problem of filling up the interaction matrix in a “zoo-centric” way or in a “phyto-centric” way. Zoo-centric studies directly sample animal activity and document the plants ‘touched’ by the animal. For example, analysis of pollen samples recovered from the body of pollinators, analysis of fecal samples of frugivores, radio-tracking data, etc. Phyto-centric studies take samples of focal individual plant species and document which animals ‘arrive’ or ‘touch’ the plants. Examples include focal watches of fruiting or flowering plants to record visitation by animals, raising insect herbivores from seed samples, identifying herbivory marks in samples of leaves, etc.

Most recent analyses of plant-animal interaction networks are phyto-centric; just 3.5% of available plant-pollinator ($N= 58$) or 36.6% plant-frugivore ($N= 22$) interaction datasets are zoo-centric (see Schleuning *et al.*, 2012). Moreover, most available datasets on host-parasite (parasitoid) or plant-herbivore interactions are “host-centric” or phyto-centric (e.g., Thébault & Fontaine, 2010; Morris *et al.*, 2013; Eklöf *et al.*, 2013). This may be related to a variety of causes, like preferred methodologies by researchers working with a particular group or system, logistic limitations, or inherent taxonomic focus of the research questions. A likely result of phyto-centric sampling would be adjacency matrices with large $A : P$ ratios. In any case we don’t have a clear view of the potential biases that taxa-focused sampling may generate in observed network patterns, for example by generating consistently asymmetric interaction matrices (Dormann *et al.*, 2009). System sym-

metry has been suggested to influence estimations of generalization levels in plants and animals when measured as I_A and I_P (Elberling & Olesen, 1999); thus, differences in I_A and I_P between networks may arise from different $A : P$ ratios rather than other ecological factors (Olesen & Jordano, 2002).

Reasonably complete analyses of interaction networks can be obtained when combining both phyto-centric and zoo-centric sampling. For example, Bosch *et al.* (2009) showed that the addition of pollen load data on top of focal-plant sampling of pollinators unveiled a significant number of interactions, resulting in important network structural changes. Connectance increased 1.43-fold, mean plant connectivity went from 18.5 to 26.4, and mean pollinator connectivity from 2.9 to 4.1; moreover, extreme specialist pollinator species (singletons in the adjacency matrix) decreased 0.6-fold. Olesen *et al.* (2011) identified pollen loads on sampled insects and added the new links to an observation-based visitation matrix, with an extra 5% of links representing the estimated number of missing links in the pollination network. The overlap between observational and pollen-load recorded links was only 33%, underscoring the value of combining methodological approaches. Zoo-centric sampling has recently been extended with the use of DNA-barcoding, for example with plant-herbivore (Jurado-Rivera *et al.*, 2009), host-parasitoid (Wirta *et al.*, 2014), and plant-frugivore interactions (González-Varo, Arroyo & Jordano, 2014). For mutualistic networks we would expect that zoo-centric sampling could help unveiling interactions of the animals with rare plant species or for relatively common plants species which are difficult to sample by direct observation. Future methodological work may provide significant advances showing how mixing different sampling strategies strengthens the completeness of network data. These mixed strategies may combine, for instance, timed watches at focal plants, spot

censuses along walked transects, pollen load or seed contents analyses, monitoring with camera traps, and DNA barcoding records. We might expect increased power of these mixed sampling approaches when combining different methods from both phyto- and zoo-centric perspectives (Bosch *et al.*, 2009; Blüthgen, 2010). Note also that the different methods could be applied in different combinations to the two distinct sets of species. However, there are no tested protocols and/or sampling designs for ecological interaction studies to suggest an optimum combination of approaches. Ideally, pilot studies would provide adequate information for each specific study setting.

187 Sampling interactions: rationale

The number of distinct pairwise interactions that we can record in a landscape (an area of relatively homogeneous vegetation, analogous to the one we would use to monitor species diversity) is equivalent to the number of distinct classes in which we can classify the recorded encounters among *individuals* of two different species. Yet, individual-based interaction networks have been only recently studied (Dupont, Trøjelsgaard & Olesen, 2011; Wells & O'Hara, 2012). The most usual approach has been to pool individual-based interaction data into species-based summaries, an approach that ignores the fact that only a fraction of individuals may actually interact given a per capita interaction effect (Wells & O'Hara, 2012). Wells & O'Hara (2012) illustrate the pros and cons of the approach. We walk in the forest and see a blackbird *Tm* picking an ivy *Hh* fruit and ingesting it: we have a record for *Tm* – *Hh* interaction. We keep advancing and record again a blackbird feeding on hawthorn *Cm* fruits so we record a *Tm* – *Cm* interaction;

as we advance we encounter another ivy plant and record a blackcap swallowing a fruit so we now have a new $Sa - Hh$ interaction, and so on. At the end we have a series of classes (e.g., $Sa - Hh$, $Tm - Hh$, $Tm - Cm$, etc.), along with their observed frequencies. Bunge & Fitzpatrick (1993) provide an early review of the main aspects and approaches to estimate the number of distinct classes C in a sample of observations.

Our sampling above would have resulted in a vector $n = [n_1 \dots n_C]'$ where n_i is the number of records in the i^{th} class. As stressed by Bunge & Fitzpatrick (1993), however, the i^{th} class would appear in the sample if and only if $n_i > 0$, and we don't know *a priori* which n_i are zero. So, n is not observable. Rather, what we get is a vector $c = [c_1 \dots c_n]'$ where c_j is the number of classes represented j times in our sampling: c_1 is the number of singletons (interactions recorded once), c_2 is the number of twin pairs (interactions with just two records), c_3 the number of triplets, etc. The problem thus turns to be estimating the number of distinct classes C from the vector of c_j values and the frequency of unobserved interactions (see "The real missing links" below).

More specifically, we usually obtain a type of reference sample (Chao *et al.*, 2014) for interactions: a series of replicated samples (e.g., observation days, 1h watches, etc.) with quantitative information, i.e., recording the number of instances of each interaction type on each day. This replicated abundance data, can be treated in three ways: 1) Abundance data within replicates: the counts of interactions, separately for each day; 2) Pooled abundance data: the counts of interactions, summed over all days (the most usual approach); and 3) Replicated incidence data: the number of days on which we recorded each interaction. Assuming a reasonable number of replicates, replicated incidence data is considered the

most robust statistically, as it takes account of heterogeneity among days (Colwell, Mao & Chang, 2004; Colwell, Dunn & Harris, 2012; Chao *et al.*, 2014). Thus, both presence-absence and weighted information on interactions can be accommodated for this purpose.

The species assemblage

When we consider an observed and recorded sample of interactions on a particular assemblage of A_{obs} and P_{obs} species (or a set of replicated samples) as a reference sample (Chao *et al.*, 2014) we may have three sources of undersampling error that are ignored by treating a reference sample as a true representation of the interactions in well-defined assemblage: 1) some animal species are actually present but not observed (zero abundance or incidence in the interactions in the reference sample), A_0 ; 2) some plant species are actually present but not observed (zero abundance or incidence in the interactions in the reference sample), P_0 ; 3) some unobserved links (the zeroes in the adjacency matrix, UL) may actually occur but not recorded. Thus a first problem is determining if A_{obs} and P_{obs} truly represent the actual species richness interacting in the assemblage. To this end we might use the replicated reference samples to estimate the true number of interacting animal A_{est} and plant P_{est} species as in traditional diversity estimation analysis (Chao *et al.*, 2014). If there are no uniques (species seen on only one day), then A_0 and P_0 will be zero, and we have A_{obs} and P_{obs} as robust estimates of the actual species richness of the assemblage. If A_0 and P_0 are not zero they estimate the minimum number of undetected animal and plant species that can be expected with a sufficiently large number of replicates, taken from the same assemblage/locality by

the same methods in the same time period. We can use extrapolation methods (Colwell, Dunn & Harris, 2012) to estimate how many additional replicate surveys it would take to reach a specified proportion g of A_{est} and P_{est} .

The interactions

We are then faced with assessing the sampling of interactions I . Table 1 summarizes the main components and targets for estimation of interaction richness. In contrast with traditional species diversity estimates, sampling networks has the paradox that despite the potentially interacting species being present in the sampled assemblage (i.e., included in the A_{obs} and P_{obs} species lists), some of their pairwise interactions are impossible to record. The reason is forbidden links. Independently of whether we sample full communities or subset communities we face a problem: some of the interactions that we can visualize in the empty adjacency matrix Δ will simply not occur. With a total of $A_{obs}P_{obs}$ “potential” interactions (eventually augmented to $A_{est}P_{est}$ in case we have undetected species), a fraction of them are impossible to record, because they are forbidden (Jordano, Bascompte & Olesen, 2003; Olesen *et al.*, 2011).

Our goal is to estimate the true number of non-null AP interactions, including interactions that actually occur but have not been observed (I_0) from the replicated incidence frequencies of interaction types: $I_{est} = I_{obs} + I_0$. Note that I_0 estimates the minimum number of undetected plant-animal interactions that can be expected with a sufficiently large number of replicates, taken from the same assemblage/locality by the same methods in the same time period. Therefore we have two types of non-observed links: UL^* and UL , corresponding to the real

272 assemblage species richness and to the observed assemblage species richness, re-
 273 spectively (Table 1).

274 Forbidden links are non-occurrences of pairwise interactions that can be ac-
 275 counted for by biological constraints, such as spatio-temporal uncoupling (Jor-
 276 dano, 1987), size or reward mismatching, foraging constraints (e.g., accessibility)
 277 (Moré *et al.*, 2012), and physiological-biochemical constraints (Jordano, 1987). We
 278 still have very little information about the frequency of forbidden links in natural
 279 communities (Jordano, Bascompte & Olesen, 2003; Stang *et al.*, 2009; Vázquez,
 280 Chacoff & Cagnolo, 2009; Olesen *et al.*, 2011; Ibanez, 2012; Maruyama *et al.*, 2014;
 281 Vizentin-Bugoni, Maruyama & Sazima, 2014) (Table 1). Forbidden links are thus
 282 represented as structural zeroes in the interaction matrix, i.e., matrix cells that
 283 cannot get a non-zero value.

284 We might expect different types of FL to occupy different parts of the Δ ma-
 285 trix, with missing cells due to phenological uncoupling, FL_P , largely distributed
 286 in the lower-right half Δ matrix and actually missed links ML distributed in its
 287 central part (Olesen *et al.*, 2010). Yet, most of these aspects remain understud-
 288 ied. Therefore, we need to account for the frequency of these structural zeros in
 289 our matrix before proceeding. For example, most measurements of connectance
 290 $C = I/(AP)$ implicitly ignore the fact that by taking the full product AP in the
 291 denominator they are underestimating the actual connectance value, i.e., the frac-
 292 tion of actual interactions I relative to the *biologically possible* ones, not to the
 293 total maximum $I_{max} = AP$.

294 Our main problem then turns to estimate the number of true missed links,
 295 i.e., those that can't be accounted for by biological constraints and that might
 296 suggest undersampling. Thus, the sampling of interactions in nature, as the sam-

297 pling of species, is a cumulative process. In our analysis, we are not re-sampling
 298 individuals, but interactions, so we made interaction-based accumulation curves.
 299 If an interaction-based curve suggests a robust sampling, it does mean that no
 300 new interactions are likely to be recorded, irrespectively of the species, as it is a
 301 whole-network sampling approach (N. Gotelli, pers. com.). We add new, distinct,
 302 interactions recorded as we increase sampling effort (Fig. 2). We can obtain an
 303 Interaction Accumulation Curve (*IAC*) analogous to a Species Curve (*SAC*) (see
 304 Supporting Information in the online data availability repository): the observed
 305 number of distinct pairwise interactions in a survey or collection as a function of
 306 the accumulated number of observations or samples (Colwell, 2009).

307 Empirical data on Forbidden Links

308 Adjacency matrices are frequently sparse, i.e., they are densely populated with
 309 zeroes, with a fraction of them being structural (unobservable interactions) (Bas-
 310 compte & Jordano, 2014). Thus, it would be a serious interpretation error to
 311 attribute the sparseness of adjacency matrices for bipartite networks to undersam-
 312 pling. The actual typology of link types in ecological interaction networks is thus
 313 more complex than just the two categories of observed and unobserved interactions
 314 (Table 1). Unobserved interactions are represented by zeroes and belong to two
 315 categories. Missing interactions may actually exist but require additional sampling
 316 or a variety of methods to be observed. Forbidden links, on the other hand, arise
 317 due to biological constraints limiting interactions and remain unobservable in na-
 318 ture, irrespectively of sampling effort (Table 1). Forbidden links *FL* may actually
 319 account for a relatively large fraction of unobserved interactions *UL* when sam-

pling taxonomically-restricted subnetworks (e.g., plant-hummingbird pollination
 networks) (Table 1). Phenological uncoupling is also prevalent in most networks,
 and may add up to explain ca. 25-40% of the forbidden links, especially in highly
 seasonal habitats, and up to 20% when estimated relative to the total number of un-
 observed interactions (Table 2). In any case, we might expect that a fraction of the
 missing links ML would be eventually explained by further biological reasons, de-
 pending on the knowledge of natural details of the particular systems. Our goal as
 naturalists would be to reduce the fraction of UL which remain as missing links; to
 this end we might search for additional biological constraints or increase sampling
 effort. For instance, habitat use patterns by hummingbirds in the Arima Valley
 network (Table 2; Snow & Snow, 1972) impose a marked pattern of microhabitat
 mismatches causing up to 44.5% of the forbidden links. A myriad of biological
 causes beyond those included as FL in Table 2 may contribute explanations for
 UL : limits of color perception and or partial preferences, presence of secondary
 metabolites in fruit pulp and leaves, toxins and combinations of monosaccharides
 in nectar, etc. For example, aside from FL , some pairwise interactions may sim-
 ply have an asymptotically-zero probability of interspecific encounter between the
 partner species, if they are very rare. However, it is surprising that just the limited
 set of forbidden link types considered in Table 1 explain between 24.6-77.2% of
 the unobserved links. Notably, the Arima Valley, Santa Virgínia, and Hato Ratón
 networks have $> 60\%$ of the unobserved links explained, which might be related
 to the fact that they are subnetworks (Arima Valley, Santa Virgínia) or relatively
 small networks (Hato Ratón). All this means that empirical networks may have
 sizable fractions of structural zeroes. Ignoring this biological fact may contribute
 to wrongly inferring undersampling of interactions in real-world assemblages.

To sum up, two elements of inference are required in the analysis of unobserved interactions in ecological interaction networks: first, detailed natural history information on the participant species that allows the inference of biological constraints imposing forbidden links, so that structural zeroes can be identified in the adjacency matrix. Second, a critical analysis of sampling robustness and a robust estimate of the actual fraction of missing links, M , resulting in a robust estimate of I . In the next sections we explore these elements of inference, using *IACs* as analogs to *SACs* to assess the robustness of interaction sampling.

Asymptotic diversity estimates

Let's assume a sampling of the diversity in a specific locality, over relatively homogeneous landscape where we aim at determining the number of species present for a particular group of organisms. To do that we carry out transects or plot samplings across the landscape or use any other type of direct or indirect recording method, adequately replicated so we obtain a number of samples. Briefly, S_{obs} is the total number of species observed in a sample, or in a set of samples. S_{est} is the estimated number of species in the community represented by the sample, or by the set of samples, where *est* indicates an estimator. With abundance data, let S_k be the number of species each represented by exactly k individuals in a single sample. Thus, S_0 is the number of undetected species (species present in the community but not included in the sample), S_1 is the number of singleton species (represented by just one individual), S_2 is the number of doubleton species (species with two individuals), etc. The total number of individuals in the sample would be:

$$n = \sum_{k=1}^{S_{obs}} S_k$$

368

369 A frequently used asymptotic, bias corrected, non-parametric estimator is S_{Chao1}
 370 (Hortal, Borges & Gaspar, 2006; Chao, 2005; Colwell, 2013):

$$S_{Chao1} = S_{obs} + \frac{S_1(S_1 - 1)}{2(S_2 + 1)}$$

371 Another frequently used alternative is the Chao2 estimator, S_{Chao2} (Gotelli &
 372 Colwell, 2001), which has been reported to have a limited bias for small sample
 373 sizes (Colwell & Coddington, 1994; Chao, 2005). Instead of using counts it uses
 374 incidence frequencies (Q_k) among samples (number of species present in just one
 375 sample, in two samples, etc.):

$$S_{Chao2} = S_{obs} + \frac{Q_1(Q_1 - 1)}{2(Q_2 + 1)}$$

376 A plot of the cumulative number of species recorded, S_n , as a function of some
 377 measure of sampling effort (say, n samples taken) yields the species accumulation
 378 curve (SAC) or collector's curve (Colwell & Coddington, 1994). Similarly, inter-
 379 action accumulation curves (IAC), analogous to SACs, can be used to assess the
 380 robustness of interactions sampling for plant-animal community datasets (Jordano,
 381 1987; Jordano, Vázquez & Bascompte, 2009; Olesen *et al.*, 2011), as discussed in
 382 the next section.

Assessing sampling effort when recording interactions

The basic method we can propose to estimate sampling effort and explicitly show the analogues with rarefaction analysis in biodiversity research is to vectorize the interaction matrix AP so that we get a vector of all the potential pairwise interactions (I_{max} , Table 1) that can occur in the observed assemblage with A_{obs} animal species and P_{obs} plant species. The new “species” we aim to sample are the pairwise interactions (Table 3). So, if we have in our community *Turdus merula* (Tm) and *Rosa canina* (Rc) and *Prunus mahaleb* (Pm), our problem will be to sample 2 new “species”: $Tm - Rc$ and $Tm - Pm$. In general, if we have $A = 1...i$, animal species and $P = 1...j$ plant species (assuming a complete list of species in the assemblage), we’ll have a vector of “new” species to sample: $A_1P_1, A_1P_2, ...A_2P_1, A_2P_2, ...A_iP_j$. We can represent the successive samples where we can potentially get records of these interactions in a matrix with the vectorized interaction matrix and columns representing the successive samples we take (Table 3). This is simply a vectorized version of the interaction matrix. This is analogous to a biodiversity sampling matrix with species as rows and sampling units (e.g., quadrats) as columns (Jordano, Vázquez & Bascompte, 2009). The package *EstimateS* (Colwell, 2013) includes a complete set of functions for estimating the mean IAC and its unconditional standard deviation from random permutations of the data, or subsampling without replacement (Gotelli & Colwell, 2001); it further reports asymptotic estimators for the expected number of distinct pairwise interactions included in a given reference sample of interaction records (see also the `specaccum` function in library `vegan` of the R Package)(R Development Core Team, 2010; Jordano, Vázquez & Bascompte,

2009; Olesen *et al.*, 2011). In particular, we may take advantage of replicated in-
 cidence data, as it takes account of heterogeneity among samples (days, censuses,
 etc.; R.K Colwell, pers. comm.) (see also Colwell, Mao & Chang, 2004; Colwell,
 Dunn & Harris, 2012; Chao *et al.*, 2014).

In this way we effectively extend sampling theory developed for species diversity
 to the sampling of ecological interactions. Yet future theoretical work will be
 needed to formally assess the similarities and differences in the two approaches
 and developing biologically meaningful null models of expected interaction richness
 with added sampling effort.

Diversity-accumulation analysis (Magurran, 1988; Hortal, Borges & Gaspar,
 2006) comes up immediately with this type of dataset. This procedure plots
 the accumulation curve for the expected number of distinct pairwise interactions
 recorded with increasing sampling effort (Jordano, Vázquez & Bascompte, 2009;
 Olesen *et al.*, 2011). Asymptotic estimates of interaction richness and its associ-
 ated standard errors and confidence intervals can thus be obtained (Hortal, Borges
 & Gaspar, 2006) (see Supplementary Online Material). It should be noted that
 the asymptotic estimate of interaction richness explicitly ignores the fact that,
 due to forbidden links, a number of pairwise interactions among the I_{max} number
 specified in the adjacency matrix Δ cannot be recorded, irrespective of sampling
 effort.

We may expect undersampling specially in moderate to large sized networks
 with multiple modules (i.e., species subsets requiring different sampling strategies)
 (Jordano, 1987; Olesen *et al.*, 2011; Chacoff *et al.*, 2012); adequate sampling may be
 feasible when interaction subwebs are studied (Olesen *et al.*, 2011; Vizentin-Bugoni,
 Maruyama & Sazima, 2014), typically with more homogeneous subsets of species

(e.g., bumblebee-pollinated flowers). In any case the sparseness of the Δ matrix is by no means an indication of undersampling whenever the issue of structural zeroes in the interaction matrices is effectively incorporated in the estimates.

For example, mixture models incorporating detectabilities have been proposed to effectively account for rare species (Mao & Colwell, 2005). In an analogous line, mixture models could be extended to samples of pairwise interactions, also with specific detectability values. These detection rate/odds could be variable among groups of interactions, depending on their specific detectability. For example, detectability of flower-pollinator interactions involving bumblebees could have a higher detectability than flower-pollinator pairwise interactions involving, say, nictidulid beetles. These more homogeneous groupings of pairwise interactions within a network define modules (Bascompte & Jordano, 2014), so we might expect that interactions of a given module (e.g., plants and their hummingbird pollinators; Fig. 1a) may share similar detectability values, in an analogous way to species groups receiving homogeneous detectability values in mixture models (Mao & Colwell, 2005). In its simplest form, this would result in a sample with multiple pairwise interactions detected, in which the number of interaction events recorded for each distinct interaction found in the sample is recorded (i.e., a column vector in Table 3, corresponding to, say, a sampling day). The number of interactions recorded for the i_{th} pairwise interaction (i.e., $A_i P_j$ in Table 3), Y_i could be treated as a Poisson random variable with a mean parameter λ_i , its detection rate. Mixture models (Mao & Colwell, 2005) include estimates for abundance-based data (their analogs in interaction sampling would be weighted data), where Y_i is a Poisson random variable with detection rate λ_i . This is combined with the incidence-based model, where Y_i is a binomial random variable (their analogous in interaction sampling

would be presence/absence records of interactions) with detection odds λ_i . Let T be the number of samples in an incidence-based data set. A Poisson/binomial density can be written as (Mao & Colwell, 2005):

$$g(y; \lambda) = \begin{cases} \frac{\lambda^y}{y!e^\lambda} & [1] \\ \binom{T}{y} \frac{\lambda^y}{(1+\lambda)^T} & [2] \end{cases}$$

where [1] corresponds to a weighted network, and [2] to a qualitative network.

The detection rates λ_i depend on the relative abundances ϕ_i of the interactions, the probability of a pairwise interaction being detected when it is present, and the sample size (the number of interactions recorded), which, in turn, is a function of the sampling effort. Unfortunately, no specific sampling model has been developed along these lines for species interactions and their characteristic features. For example, a complication factor might be that interaction abundances, ϕ_i , in real assemblages are a function of the abundances of interacting species that determine interspecific encounter rates; yet they also depend on biological factors that ultimately determine if the interaction occurs when the partner species are present. For example, λ_i should be set to zero for all FL . In its simplest form, ϕ_i could be estimated from just the product of partner species abundances, an approach recently used as a null model to assess the role of biological constraints in generating forbidden links and explaining interaction patterns (Vizentin-Bugoni, Maruyama & Sazima, 2014). Yet more complex models (e.g., Wells & O'hara 2012) should incorporate not only interspecific encounter probabilities, but also interaction detectabilities, phenotypic matching and incidence of forbidden links. Mixture models are certainly complex and for most situations of evaluating sam-

478 pling effort better alternatives include the simpler incidence-based rarefaction and
 479 extrapolation (Colwell, Dunn & Harris, 2012; Chao *et al.*, 2014).

480 The *real* missing links

481 Given that a fraction of unobserved interactions can be accounted for by for-
 482 bidden links, what about the remaining missing interactions? We have already
 483 discussed that some of these could still be related to unaccounted constraints, and
 484 still others would be certainly attributable to insufficient sampling. Would this
 485 always be the case? Multispecific assemblages of distinct taxonomic relatedness,
 486 whose interactions can be represented as bipartite networks (e.g., host-parasite,
 487 plant-animal mutualisms, plant-herbivore interactions- with two distinct sets of
 488 unrelated higher taxa), are shaped by interspecific encounters among individuals
 489 of the partner species (Fig. 2). A crucial ecological aspect limiting these inter-
 490 actions is the probability of interspecific encounter, i.e., the probability that two
 491 individuals of the partner species actually encounter each other in nature.

492 Given log-normally distributed abundances of the two species groups, the ex-
 493 pected probabilities of interspecific encounter (*PIE*) would be simply the product
 494 of the two lognormal distributions. Thus, we might expect that for low *PIE* values,
 495 pairwise interactions would be either extremely difficult to sample, or simply do
 496 not occur in nature. Consider the Nava de las Correhuelas interaction web (NCH,
 497 Table 2), with $A = 36$, $P = 25$, $I = 181$, and almost half of the unobserved inter-
 498 actions not accounted for by forbidden links, thus $M = 53.1\%$. Given the robust
 499 sampling of this network (Jordano, Vázquez & Bascompte, 2009), a sizable fraction
 500 of these possible but missing links would be simply not occurring in nature, most

likely due to extremely low PIE , in fact asymptotically zero. Given the vectorized list of pairwise interactions for NCH, I computed the PIE values for each one by multiplying element-wise the two species abundance distributions. The $PIE_{max} = 0.0597$, being a neutral estimate, based on the assumption that interactions occur in proportion to the species-specific local abundances. With $PIE_{median} < 1.4 \cdot 10^{-4}$ we may safely expect (note the quantile estimate $Q_{75\%} = 3.27 \cdot 10^{-4}$) that a sizable fraction of these missing interactions may not occur according to this neutral expectation (Jordano, 1987; Olesen *et al.*, 2011) (neutral forbidden links, *sensu* Canard *et al.*, 2012).

When we consider the vectorized interaction matrix, enumerating all pairwise interactions for the AP combinations, the expected probabilities of finding a given interaction can be estimated with a Good-Turing approximation (Good, 1953). The technique, developed by Alan Turing and I.J. Good with applications to linguistics and word analysis (Gale & Sampson, 1995) has been recently extended in novel ways for ecological analyses (Chao *et al.*, 2015). It estimates the probability of recording an interaction of a hitherto unseen pair of partners, given a set of past records of interactions between other species pairs. Let a sample of N interactions so that n_r distinct pairwise interactions have exactly r records. All Good-Turing estimators obtain the underlying frequencies of events as:

$$P(X) = \frac{(N_X + 1)}{T} \left(1 - \frac{E(1)}{T}\right) \quad (1)$$

where X is the pairwise interaction, N_X is the number of times interaction X is recorded, T is the sample size (number of distinct interactions recorded) and $E(1)$ is an estimate of how many different interactions were recorded exactly once.

Strictly speaking Equation (1) gives the probability that the next interaction type recorded will be X , after sampling a given assemblage of interacting species. In other words, we scale down the maximum-likelihood estimator $\frac{n}{T}$ by a factor of $\frac{1-E(1)}{T}$. This reduces all the probabilities for interactions we have recorded, and makes room for interactions we haven't seen. If we sum over the interactions we have seen, then the sum of $P(X)$ is $1 - \frac{1-E(1)}{T}$. Because probabilities sum to one, we have the left-over probability of $P_{new} = \frac{E(1)}{T}$ of seeing something new, where new means that we sample a new pairwise interaction. Note, however, that Good-Turing estimators, the traditional asymptotic estimators, do not account in our case for the forbidden interactions.

Discussion

Recent work has inferred that most data available for interaction networks are incomplete due to undersampling, resulting in a variety of biased parameters and network patterns (Chacoff *et al.*, 2012). It is important to note, however, that in practice, most surveyed networks to date have been subnets of much larger networks. This is also true for protein interaction, gene regulation, and metabolic networks, where only a subset of the molecular entities in a cell have been sampled (Stumpf, Wiuf & May, 2005). Despite recent attempts to document whole ecosystem meta-networks (Pocock, Evans & Memmott, 2012), it is likely that most ecological interaction networks will illustrate just major ecosystem compartments. Due to their high generalization, high temporal and spatial turnover, and high complexity of association patterns, adequate sampling of ecological interaction networks is challenging and requires extremely large sampling effort. Undersam-

pling of ecological networks may originate from the analysis of assemblage subsets (e.g., taxonomically or functionally defined), and/or from logistically-limited sampling effort. It is extremely hard to robustly sample the set of biotic interactions even for relatively simple, species-poor assemblages; thus, we need to assess how robust is the characterization of the adjacency matrix Δ . Concluding that an ecological network dataset is undersampled just by its sparseness would be unrealistic. The reason stems from a biological fact: a sizeable fraction of the maximum, potential links that can be recorded among two distinct sets of species is simply unobservable, irrespective of sampling effort (Jordano, 1987). In addition, sampling effort needs to be explicitly gauged because of its potential influence on parameter estimates for the network.

Missing links are a characteristic feature of all plant-animal interaction networks, and likely pervade other ecological interactions. Important natural history details explain a fraction of them, resulting in unrealizable interactions (i.e., forbidden interactions) that define structural zeroes in the interaction matrices and contribute to their extreme sparseness. Sampling interactions is a way to monitor biodiversity beyond the simple enumeration of component species and to develop efficient and robust inventories of functional interactions. Yet no sampling theory for interactions is available. Focusing just on the realized interactions or treating missing interactions as the expected unique result of sampling bias would miss important components to understand how all sorts of interactions coevolve within complex webs of interdependence among species.

Contrary to species inventories, a sizable fraction of non-observed pairwise interactions cannot be sampled, due to biological constraints that forbid their occurrence. Moreover, recent implementations of inference methods for unobserved

species (Chao *et al.*, 2015) or for individual-based data (Wells & O’Hara, 2012) can be combined with the forbidden link approach. They do not account either for the existence of these ecological constraints, but can help in estimating their relative importance, simply by the difference between the asymptotic estimate of interaction richness *in a robustly-sampled* assemblage and the maximum richness I_{max} of interactions.

Ecological interactions provide the wireframe supporting the lives of species, and they also embed crucial ecosystem functions which are fundamental for supporting the Earth system. We still have a limited knowledge of the biodiversity of ecological interactions, and they are being lost (extinct) at a very fast pace, frequently preceding species extinctions (Valiente-Banuet *et al.*, 2014). We urgently need robust techniques to assess the completeness of ecological interactions networks because this knowledge will allow the identification of the minimal components of their ecological complexity that need to be restored to rebuild functional ecosystems after perturbations.

Acknowledgements

I am indebted to Jens M. Olesen, Alfredo Valido, Jordi Bascompte, Thomas Lewinshon, John N. Thompson, Nick Gotelli, Carsten Dormann, and Paulo R. Guimarães Jr. for useful and thoughtful discussion at different stages of this manuscript. Jeferson Vizentin-Bugoni kindly helped with the Sta Virgínia data. Jens M. Olesen kindly made available the Grundvad dataset; together with Robert K. Colwell, Néstor Pérez-Méndez, JuanPe González-Varo, and Paco Rodríguez provided most useful comments to a final version of the ms. Robert Colwell shared a

number of crucial suggestions that clarified my vision of sampling ecological interactions, and the final manuscript was greatly improved with comments from three anonymous referees. The study was supported by a Junta de Andalucía Excellence Grant (RNM-5731), as well as a Severo Ochoa Excellence Award from the Ministerio de Economía y Competitividad (SEV-2012-0262). The Agencia de Medio Ambiente, Junta de Andalucía, provided generous facilities that made possible my long-term field work in different natural parks.

Data accessibility

This review does not use new raw data, but includes some re-analyses of previously published material. All the original data supporting the paper, R code, supplementary figures, and summaries of analytical protocols is available at the author's GitHub repository (https://github.com/pedroj/MS_Network-Sampling), with DOI: 10.5281/zenodo.29437.

References

- Banasek-Richter, C., Cattin, M. & Bersier, L. (2004) Sampling effects and the robustness of quantitative and qualitative food-web descriptors. *Journal of Theoretical Biology*, **226**, 23–32.
- Bascompte, J. & Jordano, P. (2014) *Mutualistic networks*. Monographs in Population Biology, No. 53. Princeton University Press, Princeton, NJ.
- Bersier, L., Banasek-Richter, C. & Cattin, M. (2002) Quantitative descriptors of food-web matrices. *Ecology*, **83**, 2394–2407.

- 615 Blüthgen, N. (2010) Why network analysis is often disconnected from community
616 ecology: A critique and an ecologist's guide. *Basic And Applied Ecology*, **11**,
617 185–195.
- 618 Bosch, J., Martín González, A.M., Rodrigo, A. & Navarro, D. (2009) Plant-
619 pollinator networks: adding the pollinator's perspective. *Ecology Letters*, **12**,
620 409–419.
- 621 Brose, U., Martinez, N. & Williams, R. (2003) Estimating species richness: Sen-
622 sitivity to sample coverage and insensitivity to spatial patterns. *Ecology*, **84**,
623 2364–2377.
- 624 Bunge, J. & Fitzpatrick, M. (1993) Estimating the number of species: a review.
625 *Journal of the American Statistical Association*, **88**, 364–373.
- 626 Canard, E., Mouquet, N., Marescot, L., Gaston, K.J., Gravel, D. & Mouillot,
627 D. (2012) Emergence of structural patterns in neutral trophic networks. *PLoS*
628 *ONE*, **7**, e38295.
- 629 Chacoff, N.P., Vazquez, D.P., Lomascolo, S.B., Stevani, E.L., Dorado, J. & Padrón,
630 B. (2012) Evaluating sampling completeness in a desert plant-pollinator network.
631 *Journal of Animal Ecology*, **81**, 190–200.
- 632 Chao, A. (2005) Species richness estimation. *Encyclopedia of Statistical Sciences*,
633 pp. 7909–7916. Oxford University Press, New York, USA.
- 634 Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K. & Elli-
635 son, A.M. (2014) Rarefaction and extrapolation with Hill numbers: a framework

- 636 for sampling and estimation in species diversity studies. *Ecological Monographs*,
637 **84**, 45–67.
- 638 Chao, A., Hsieh, T.C., Chazdon, R.L., Colwell, R.K. & Gotelli, N.J. (2015) Un-
639 veiling the species-rank abundance distribution by generalizing the Good-Turing
640 sample coverage theory. *Ecology*, **96**, 1189–1201.
- 641 Coddington, J.A., Agnarsson, I., Miller, J.A., Kuntner, M. & Hormiga, G. (2009)
642 Undersampling bias: the null hypothesis for singleton species in tropical arthro-
643 pod surveys. *Journal of Animal Ecology*, **78**, 573–584.
- 644 Cohen, J.E. (1978) *Food webs and niche space*. Princeton University Press, Prince-
645 ton, New Jersey, US.
- 646 Cohen, J.E., Beaver, R.A., Cousins, S.H., DeAngelis, D.L., Goldwasser, L., Heong,
647 K.L., Holt, R.D., Kohn, A.J., Lawton, J.H., Martinez, N., O'Malley, R., Page,
648 L.M., Patten, B.C., Pimm, S.L., Polis, G., Rejmanek, M., Schoener, T.W.,
649 Schenly, K., Sprules, W.G., Teal, J.M., Ulanowicz, R., Warren, P.H., Wilbur,
650 H.M. & Yodis, P. (1993) Improving food webs. *Ecology*, **74**, 252–258.
- 651 Colwell, R. & Coddington, J. (1994) Estimating terrestrial biodiversity through ex-
652 trapolation. *Philosophical Transactions Of The Royal Society Of London Series*
653 *B-Biological Sciences*, **345**, 101–118.
- 654 Colwell, R.K. (2009) Biodiversity: concepts, patterns, and measurement. *The*
655 *Princeton Guide to Ecology* (ed. S.A. Levin), pp. 257–263. Princeton University
656 Press, Princeton.
- 657 Colwell, R.K. (2013) EstimateS: Biodiversity Estimation. -, pp. 1–33.

- 658 Colwell, R.K., Dunn, R.R. & Harris, N.C. (2012) Coextinction and persistence of
 659 dependent species in a changing world. *Annual Review of Ecology Evolution and*
 660 *Systematics*, **43**, 183–203.
- 661 Colwell, R.K., Mao, C.X. & Chang, J. (2004) Interpolating, extrapolating, and
 662 comparing incidence-based species accumulation curves. *Ecology*, **85**, 2717–2727.
- 663 Dorado, J., Vazquez, D.P., Stevani, E.L. & Chacoff, N.P. (2011) Rareness and
 664 specialization in plant-pollinator networks. *Ecology*, **92**, 19–25.
- 665 Dormann, C.F., Frund, J., Bluthgen, N. & Gruber, B. (2009) Indices, graphs and
 666 null models: Analyzing bipartite ecological networks. *Open Ecology Journal*, **2**,
 667 7–24.
- 668 Dupont, Y.L., Trøjelsgaard, K. & Olesen, J.M. (2011) Scaling down from species
 669 to individuals: a flower-visitation network between individual honeybees and
 670 thistle plants. *Oikos*, **120**, 170–177.
- 671 Dupont, Y.L., Trøjelsgaard, K., Hagen, M., Henriksen, M.V., Olesen, J.M., Ped-
 672 ersen, N.M.E. & Kissling, W.D. (2014) Spatial structure of an individual-based
 673 plant-pollinator network. *Oikos*, **123**, 1301–1310.
- 674 Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N.P.,
 675 Dalsgaard, B., de Sassi, C., Galetti, M., Guimaraes, P.R., Lomáscolo, S.B.,
 676 Martín González, A.M., Pizo, M.A., Rader, R., Rodrigo, A., Tylianakis, J.M.,
 677 Vazquez, D.P. & Allesina, S. (2013) The dimensionality of ecological networks.
 678 *Ecology Letters*, **16**, 577–583.

- 679 Elberling, H. & Olesen, J.M. (1999) The structure of a high latitude plant-flower
680 visitor system: the dominance of flies. *Ecography*, **22**, 314–323.
- 681 Frund, J., McCann, K.S. & Williams, N.M. (2015) Sampling bias is a challenge
682 for quantifying specialization and network structure: lessons from a quantitative
683 niche model. *Oikos*, pp. n/a–n/a.
- 684 Gale, W.A. & Sampson, G. (1995) Good-Turing frequency estimation without
685 tears. *Journal of Quantitative Linguistics*, **2**, 217–237.
- 686 Gibson, R.H., Knott, B., Eberlein, T. & Memmott, J. (2011) Sampling method
687 influences the structure of plant–pollinator networks. *Oikos*, **120**, 822–831.
- 688 González-Varo, J.P., Arroyo, J.M. & Jordano, P. (2014) Who dispersed the seeds?
689 The use of DNA barcoding in frugivory and seed dispersal studies. *Methods in*
690 *Ecology and Evolution*, **5**, 806–814.
- 691 Good, I.J. (1953) The population frequencies of species and the estimation of
692 population parameters. *Biometrika*, **40**, 237–264.
- 693 Gotelli, N.J. & Colwell, R.K. (2011) Estimating species richness. *Biological Di-*
694 *versity Frontiers in Measurement and Assessment* (eds. A.E. Magurran & B.J.
695 McGill), pp. 39–54. Oxford University Press, Oxford, UK.
- 696 Gotelli, N. & Colwell, R. (2001) Quantifying biodiversity: procedures and pitfalls
697 in the measurement and comparison of species richness. *Ecology Letters*, **4**,
698 379–391.
- 699 Hortal, J., Borges, P. & Gaspar, C. (2006) Evaluating the performance of species

- richness estimators: sensitivity to sample grain size. *Journal of Animal Ecology*,
75, 274–287.
- Ibanez, S. (2012) Optimizing size thresholds in a plant–pollinator interaction web:
towards a mechanistic understanding of ecological networks. *Oecologia*, **170**,
233–242.
- Jordano, P. (1987) Patterns of mutualistic interactions in pollination and seed dis-
persal: connectance, dependence asymmetries, and coevolution. *The American*
Naturalist, **129**, 657–677.
- Jordano, P., Bascompte, J. & Olesen, J. (2003) Invariant properties in coevolu-
tionary networks of plant-animal interactions. *Ecology Letters*, **6**, 69–81.
- Jordano, P., Vázquez, D. & Bascompte, J. (2009) Redes complejas de interac-
ciones planta—animal. *Ecología y evolución de interacciones planta-animal* (eds.
R. Medel, R. Dirzo & R. Zamora), pp. 17–41. Editorial Universitaria, Santiago,
Chile.
- Jurado-Rivera, J.A., Vogler, A.P., Reid, C.A.M., Petitpierre, E. & Gomez-Zurita,
J. (2009) DNA barcoding insect-host plant associations. *Proceedings Of The*
Royal Society B-Biological Sciences, **276**, 639–648.
- Magurran, A. (1988) *Ecological diversity and its measurement*. Princeton Univer-
sity Press, Princeton, US.
- Mao, C. & Colwell, R.K. (2005) Estimation of species richness: mixture models,
the role of rare species, and inferential challenges. *Ecology*, **86**, 1143–1153.

- 721 Martinez, N.D. (1993) Effects of resolution on food web structure. *Oikos*, **66**,
722 403–412.
- 723 Martinez, N. (1991) Artifacts or attributes? Effects of resolution on food-web
724 patterns in Little Rock Lake food web. *Ecological Monographs*, **61**, 367–392.
- 725 Maruyama, P.K., Vizentin-Bugoni, J., Oliveira, G.M., Oliveira, P.E. & Dalsgaard,
726 B. (2014) Morphological and spatio-temporal mismatches shape a neotropical
727 savanna plant-hummingbird network. *Biotropica*, **46**, 740–747.
- 728 Memmott, J., Alonso, D., Berlow, E., Dobson, A., Dunne, J.A., Solé, R.V. &
729 Weitz, J.S. (2006) Biodiversity loss and ecological network structure. *Ecological*
730 *Networks Linking Structure to Dynamics in Food Webs* (eds. M. Pascual & J.A.
731 Dunne), pp. 325–347. Oxford University Press, Oxford, UK.
- 732 Moré, M., Amorim, F.W., Benitez-Vieyra, S., Medina, A.M., Sazima, M. &
733 Cocucci, A.A. (2012) Armament Imbalances: Match and Mismatch in Plant-
734 Pollinator Traits of Highly Specialized Long-Spurred Orchids. *PLoS ONE*, **7**,
735 e41878.
- 736 Morris, R.J., Gripenberg, S., Lewis, O.T. & Roslin, T. (2013) Antagonistic inter-
737 action networks are structured independently of latitude and host guild. *Ecology*
738 *Letters*, **17**, 340–349.
- 739 Nielsen, A. & Bascompte, J. (2007) Ecological networks, nestedness and sampling
740 effort. *Journal of Ecology*, **95**, 1134–1141–1141.
- 741 Olesen, J.M., Bascompte, J., Dupont, Y.L., Elberling, H. & Jordano, P. (2011)

- 742 Missing and forbidden links in mutualistic networks. *Proceedings Of The Royal*
 743 *Society B-Biological Sciences*, **278**, 725–732.
- 744 Olesen, J.M., Dupont, Y.L., O’gorman, E., Ings, T.C., Layer, K., Melin, C.J.,
 745 Trjelsgaard, K., Pichler, D.E., Rasmussen, C. & Woodward, G. (2010) From
 746 Broadstone to Zackenberg. *Advances in Ecological Research*, **42**, 1–69.
- 747 Olesen, J. & Jordano, P. (2002) Geographic patterns in plant-pollinator mutualistic
 748 networks. *Ecology*, **83**, 2416–2424.
- 749 Olito, C. & Fox, J.W. (2014) Species traits and abundances predict metrics of
 750 plant-pollinator network structure, but not pairwise interactions. *Oikos*, **124**,
 751 428–436.
- 752 Ollerton, J. & Cranmer, L. (2002) Latitudinal trends in plant-pollinator interac-
 753 tions: are tropical plants more specialised? *Oikos*, **98**, 340–350.
- 754 Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes,
 755 R.J., Bruford, M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops,
 756 N., Dulloo, E., Faith, D., Freyhof, J., Gregory, R.D., Heip, C., Hoft, R., Hurtt,
 757 G., Jetz, W., Karp, D.S., Mcgeoch, M., Obura, D., Onoda, Y., Pettorelli, N.,
 758 Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S., Turak, E., Walpole, M.
 759 & Wegmann, M. (2013) Essential biodiversity variables. *Science*, **339**, 277–278.
- 760 Pocock, M.J.O., Evans, D.M. & Memmott, J. (2012) The Robustness and Restora-
 761 tion of a Network of Ecological Networks. *Science*, **335**, 973–977.
- 762 Preston, F. (1948) The commonness, and rarity, of species. *Ecology*, **29**, 254–283.

- 763 R Development Core Team (2010) *R: A language and environment for statis-*
764 *tical computing*. R Foundation for Statistical Computing. Vienna, Austria.
765 <http://www.R-project.org>, Vienna, Austria.
- 766 Rivera-Hutinel, A., Bustamante, R.O., Marín, V.H. & Medel, R. (2012) Effects of
767 sampling completeness on the structure of plant-pollinator networks. *Ecology*,
768 **93**, 1593–1603.
- 769 Schleuning, M., Frund, J., Klein, A.M., Abrahamczyk, S., Alarcón, R., Albrecht,
770 M., Andersson, G.K.S., Bazarian, S., Böhning-Gaese, K., Bommarco, R., Dals-
771 gaard, B., Dehling, D.M., Gotlieb, A., Hagen, M., Hickler, T., Holzschuh, A.,
772 Kaiser-Bunbury, C.N., Kreft, H., Morris, R.J., Sandel, B., Sutherland, W.J.,
773 Svenning, J.C., Tscharntke, T., Watts, S., Weiner, C.N., Werner, M., Williams,
774 N.M., Winqvist, C., Dormann, C.F. & Blüthgen, N. (2012) Specialization of
775 mutualistic interaction networks decreases toward tropical latitudes. *Current*
776 *Biology*, **22**, 1925–1931.
- 777 Snow, B. & Snow, D. (1972) Feeding niches of hummingbirds in a Trinidad valley.
778 *Journal of Animal Ecology*, **41**, 471–485.
- 779 Snow, B. & Snow, D. (1988) *Birds and berries*. Poyser, Calton, UK.
- 780 Stang, M., Klinkhamer, P., Waser, N.M., Stang, I. & van der Meijden, E. (2009)
781 Size-specific interaction patterns and size matching in a plant-pollinator inter-
782 action web. *Annals Of Botany*, **103**, 1459–1469.
- 783 Strogatz, S. (2001) Exploring complex networks. *Nature*, **410**, 268–276.
- 784 Stumpf, M.P.H., Wiuf, C. & May, R.M. (2005) Subnets of scale-free networks are

- not scale-free: Sampling properties of networks. *Proceedings of the National Academy of Sciences USA*, **102**, 4221–4224.
- Thébault, E. & Fontaine, C. (2010) Stability of ecological communities and the architecture of mutualistic and trophic networks. *Science*, **329**, 853–856.
- Valiente-Banuet, A., Aizen, M.A., Alcántara, J.M., Arroyo, J., Cocucci, A., Galetti, M., García, M.B., García, D., Gomez, J.M., Jordano, P., Medel, R., Navarro, L., Obeso, J.R., Oviedo, R., Ramírez, N., Rey, P.J., Traveset, A., Verdú, M. & Zamora, R. (2014) Beyond species loss: the extinction of ecological interactions in a changing world. *Functional Ecology*, **29**, 299–307.
- Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009) Evaluating multiple determinants of the structure of plant-animal mutualistic networks. *Ecology*, **90**, 2039–2046.
- Vazquez, D.P., Ramos-Jiliberto, R., Urbani, P. & Valdovinos, F.S. (2015) A conceptual framework for studying the strength of plant-animal mutualistic interactions. *Ecology Letters*, **18**, 385–400.
- Vazquez, D., Morris, W. & Jordano, P. (2005) Interaction frequency as a surrogate for the total effect of animal mutualists on plants. *Ecology Letters*, **8**, 1088–1094.
- Vizentin-Bugoni, J., Maruyama, P.K. & Sazima, M. (2014) Processes entangling interactions in communities: forbidden links are more important than abundance in a hummingbird-plant network. *Proceedings Of The Royal Society B-Biological Sciences*, **281**, 20132397–20132397.
- Vizentin-Bugoni, J., Maruyama, P.K., Debastiani, V.J., Duarte, L.d.S., Dalsgaard,

- 807 B. & Sazima, M. (2016) Influences of sampling effort on detected patterns and
808 structuring processes of a Neotropical plant–hummingbird network. *Journal of*
809 *Animal Ecology*, **85**, 262–272. ISSN 1365-2656.
810 URL <http://dx.doi.org/10.1111/1365-2656.12459>
- 811 Wells, K. & O'Hara, R.B. (2012) Species interactions: estimating per-individual
812 interaction strength and covariates before simplifying data into per-species eco-
813 logical networks. *Methods in Ecology and Evolution*, **4**, 1–8.
- 814 Wirta, H.K., Hebert, P.D.N., Kaartinen, R., Prosser, S.W., Várkonyi, G. & Roslin,
815 T. (2014) Complementary molecular information changes our perception of food
816 web structure. *Proceedings of the National Academy of Sciences USA*, **111**,
817 1885–1890.

Figure captions

Figure 1. Sampling ecological interaction networks (e.g., plant-animal interactions) usually focus on different types of subsampling the full network, yielding submatrices $\Delta[m, n]$ of the full interaction matrix Δ with A and P animal and plant species. a) all the potential plants interacting with a subset of the animals (e.g., studying just the hummingbird-pollinated flower species in a community); b) all the potential animal species interacting with a subset of the plant species (e.g., studying the frugivore species feeding on figs *Ficus* in a community); and c) sampling a subset of all the potential animal species interacting with a subset of all the plant species (e.g., studying the plant-frugivore interactions of the rainforest understory).

Figure 2. Sampling species interactions in natural communities. Suppose an assemblage with $A = 3$ animal species (red, species 1–3 with three, two, and 1 individuals, respectively) and $P = 3$ plant species (green, species a-c with three individuals each) (colored balls), sampled with increasing effort in steps 1 to 6 (panels). In Step 1 we record animal species 1 and plant species 1 and 2 with a total of three interactions (black lines) represented as two distinct interactions: $1 - a$ and $1 - b$. As we advance our sampling (panels 1 to 6, illustrating e.g., additional sampling days) we record new distinct interactions. Note that we actually sample and record interactions among individuals, yet we pool the data across species to get a species by species interaction matrix. Few network analyses have been carried out on individual data (Dupont *et al.*, 2014).

842 **Figures**

Figure 1:





Figure 2:



Jordano - Figure 1

Table captions

Table 1. A taxonomy of link types for ecological interactions (Olesen *et al.* 2011).

A , number of animal species; P , number of plant species; I , number of observed links; $C = 100I/(AP)$, connectance; FL , number of forbidden links; and ML , number of missing links. As natural scientists, our ultimate goal is to eliminate ML from the equation $FL = AP - I - ML$, which probably is not feasible given logistic sampling limitations. When we, during our study, estimate ML to be negligible, we cease observing and estimate I and FL .

Table 2. Frequencies of different type of forbidden links in natural plant-animal interaction assemblages. AP , maximum potential links, I_{max} ; I , number of observed links; UL , number of unobserved links; FL , number of forbidden links; FL_P , phenology; FL_S , size restrictions; FL_A , accessibility; FL_O , other types of restrictions; ML , unknown causes (missing links). Relative frequencies (in parentheses) calculated over $I_{max} = AP$ for I , ML , and FL ; for all forbidden links types, calculated over FL . References, from left to right: Olesen *et al.* 2008; Olesen & Myrthue unpubl.; Snow & Snow 1972 and Jordano *et al.* 2006; Vizentin-Bugoni *et al.* 2014; Jordano *et al.* 2009; Olesen *et al.* 2011.

Table 3. A vectorized interaction matrix.

Table 4. Sampling statistics for three plant-animal interaction networks (Olesen *et al.* 2011). Symbols as in Table 1; N , number of records; $Chao1$ and ACE are asymptotic estimators for the number of distinct pairwise interactions I (Hortal

867 *et al.* 2006), and their standard errors; C , sample coverage for rare interactions
868 (Chao & Jost 2012). Scaled asymptotic estimators and their confidence intervals
869 (CI) were calculated by weighting *Chao1* and *ACE* with the observed frequencies
870 of forbidden links.

871

872 Tables

Table 1:

Link type	Formulation	Definition
Potential links	$I_{max} = A_{obs}P_{obs}$	Size of observed network matrix, i.e. maximum number of potentially observable interactions; A_{obs} and P_{obs} , numbers of interacting animal and plant species, respectively. These might be below the real numbers of animal and plant species, A_{est} and P_{est} .
Observed links	I_{obs}	Total number of observed links in the network given a sufficient sampling effort. Number of ones in the adjacency matrix.
True links	I_{est}	Total number of links in the network given a sufficient sampling effort; expected for the augmented $A_{est}P_{est}$ matrix.
Unobserved links	$UL = I_{max} - I_{obs}$	Number of zeroes in the adjacency matrix.
True unobserved links	$UL* = I_{max} - I_{obs}$	Number of zeroes in the augmented adjacency matrix that, eventually, includes unobserved species.
Forbidden links	FL	Number of links, which remain unobserved because of linkage constraints, irrespectively of sufficient sampling effort.
Observed Missing links	$ML = A_{obs}P_{obs} - I_{obs} - FL$	Number of links, which may exist in nature but need more sampling effort and/or additional sampling methods to be observed.
True Missing links	$ML* = A_{est}P_{est} - I_{est} - FL$	Number of links, which may exist in nature but need more sampling effort and/or additional sampling methods to be observed. Augments ML for the $A_{est}P_{est}$ matrix.

Table 2:

Link type	Pollination			Seed dispersal		
	Zackenber	Grundvad	Arima Valley	Sta. Virginia	Hato Ratón	Nava Correhuelas
I_{max}	1891	646	522	423	272	825
I	268 (0.1417)	212 (0.3282)	185 (0.3544)	86 (0.1042)	151 (0.4719)	181 (0.2194)
UL	1507 (0.7969)	434 (0.6718)	337 (0.6456)	337 (0.4085)	169 (0.5281)	644 (0.7806)
FL	530 (0.3517)	107 (0.2465)	218 (0.6469)	260 (0.7715)	118 (0.6982)	302 (0.4689)
FL_P	530 (1.0000)	94 (0.2166)	0 (0.0000)	120 (0.1624)	67 (0.3964)	195 (0.3028)
FL_S	... (...)	8 (0.0184)	30 (0.0890)	140 (0.1894)	31 (0.1834)	46 (0.0714)
FL_A	... (...)	5 (0.0115)	150 (0.445) ^a	... (...)	20 (0.1183)	61 (0.0947)
FL_O	... (...)	... (...)	38 (0.1128) ^b	... (...)	... (...)	363 (0.5637)
ML	977 (0.6483)	327 (0.7535)	119 (0.3531)	77 (0.1042)	51 (0.3018)	342 (0.5311)

^a, Lack of accessibility due to habitat uncoupling, i.e., canopy-foraging species vs. understory species.

^b, Colour restrictions, and reward per flower too small relative to the size of the bird.

Dots indicate no data available for the FL type.

Table 3:

Interaction	Sample 1	Sample 2	Sample 3	...	Sample i
A1 - P1	12	2	0	...	6
A1 - P2	0	0	0	...	1
...
A5 - P3	5	0	1	...	18
A5 - P4	1	0	1	...	3
...
A _i - P _i	1	0	1	...	2

Table 4:

	Hato Ratón	Nava Correhuelas	Zackenberg
A	17	33	65
P	16	25	31
I_{max}	272	825	1891
N	3340	8378	1245
I	151	181	268
C	0.917	0.886	0.707
$Chao1$	263.1 ± 70.9	231.4 ± 14.2	509.6 ± 54.7
ACE	240.3 ± 8.9	241.3 ± 7.9	566.1 ± 14.8
% <i>unobserved</i> ^a	8.33	15.38	47.80

^a, estimated with library Jade (R Core Development Team 2010, Chao *et al.* 2015)