

Nonlocal translationally invariant nuclear density


Michael Gennari

TRIUMF – Co-op from University of Waterloo

In collaboration with


Petr Navrátil, Angelo Calci, and
Matteo Vorabbi

2018-03-01



No-core shell model (NCSM)


- NCSM is an *ab initio* approach to solve the many-body Schrödinger equation for bound states (narrow resonances) starting from [high-precision NN+NNN interactions](#)
- Uses large (but finite!) expansions in HO many-body basis states



$$\Psi^A = \sum_{N=0}^{N_{max}} \sum_i c_{Ni} \Phi_{Ni}^A$$


- Translational invariance of the internal wave function is preserved when single-particle Slater Determinant (SD) basis is used with N_{max} truncation

$$\langle \vec{r}_1 \cdots \vec{r}_A \vec{\sigma}_1 \cdots \vec{\sigma}_A \vec{\tau}_1 \cdots \vec{\tau}_A | A\lambda JM \rangle_{SD} = \langle \vec{\xi}_1 \cdots \vec{\xi}_{A-1} \vec{\sigma}_1 \cdots \vec{\sigma}_A \vec{\tau}_1 \cdots \vec{\tau}_A | A\lambda JM \rangle \varphi_{000}(\vec{\xi}_0)$$




No-core shell model (NCSM)

- NCSM is an *ab initio* approach to solve the many-body Schrödinger equation for bound states (narrow resonances) starting from [high-precision NN+NNN interactions](#)
- Uses large (but finite!) expansions in HO many-body basis states




$$\Psi^A = \sum_{N=0}^{N_{max}} \sum_i c_{Ni} \Phi_{Ni}^A$$

- Translational invariance of the internal wave function is preserved when single-particle Slater Determinant (SD) basis is used with N_{max} truncation




3

Intrinsic
wavefunction



COM
wavefunction



$$\langle \vec{r}_1 \cdots \vec{r}_A \vec{\sigma}_1 \cdots \vec{\sigma}_A \vec{\tau}_1 \cdots \vec{\tau}_A | A\lambda JM \rangle_{SD} = \langle \vec{\xi}_1 \cdots \vec{\xi}_{A-1} \vec{\sigma}_1 \cdots \vec{\sigma}_A \vec{\tau}_1 \cdots \vec{\tau}_A | A\lambda JM \rangle \varphi_{000}(\vec{\xi}_0)$$

Coordinate form of the density

4

Nonlocal translationally invariant density

- arXiv:1712.02879; Phys. Rev. C, in press.

$$\begin{aligned} & \langle A\lambda_f J_f M_f | \rho_{op}(\vec{r} - \vec{R}, \vec{r}' - \vec{R}) | A\lambda_i J_i M_i \rangle \\ &= \left(\frac{A}{A-1} \right)^{\frac{3}{2}} \sum_{\hat{J}_f} \frac{1}{\hat{J}_f} (J_i M_i K k | J_f M_f) \left(Y_l^* \left(\widehat{\vec{r} - \vec{R}} \right) Y_{l'}^* \left(\widehat{\vec{r}' - \vec{R}} \right) \right)_k^{(K)} \\ & \times R_{n,l} \left(\sqrt{\frac{A}{A-1}} |\vec{r} - \vec{R}| \right) R_{n',l'} \left(\sqrt{\frac{A}{A-1}} |\vec{r}' - \vec{R}| \right) \\ & \times (M^K)_{n,l,n',l',n_1,l_1,n_2,l_2}^{-1} (-1)^{l_1+l_2+K+j_2-\frac{1}{2}} \hat{J}_1 \hat{J}_2 \hat{K} \begin{Bmatrix} j_1 & j_2 & K \\ l_2 & l_1 & 1/2 \end{Bmatrix} \\ & \times \frac{(-1)}{\hat{K}} {}_{SD} \langle A\lambda_f J_f \parallel (a_{n_1 l_1 j_1}^\dagger \tilde{a}_{n_2 l_2 j_2})^{(K)} \parallel A\lambda_i J_i \rangle_{SD} \end{aligned}$$

Coordinate form of the density

Normalization

5

Nonlocal translationally invariant density

- arXiv:1712.02879; Phys. Rev. C, in press.

$$\int d\vec{x} \langle A\lambda JM | \rho_{op}^{phys}(\vec{x}) | A\lambda JM \rangle = A$$

$$\begin{aligned} & \langle A\lambda_f J_f M_f | \rho_{op}(\vec{r} - \vec{R}, \vec{r}' - \vec{R}) | A\lambda_i J_i M_i \rangle \\ &= \left(\frac{A}{A-1} \right)^{\frac{3}{2}} \sum_{\hat{J}_f} \frac{1}{\hat{J}_f} (J_i M_i K k | J_f M_f) \left(Y_l^* \left(\widehat{\vec{r} - \vec{R}} \right) Y_{l'}^* \left(\widehat{\vec{r}' - \vec{R}} \right) \right)_k^{(K)} \\ & \times R_{n,l} \left(\sqrt{\frac{A}{A-1}} |\vec{r} - \vec{R}| \right) R_{n',l'} \left(\sqrt{\frac{A}{A-1}} |\vec{r}' - \vec{R}| \right) \\ & \times (M^K)_{n,l,n',l',n_1,l_1,n_2,l_2}^{-1} (-1)^{l_1+l_2+K+j_2-\frac{1}{2}} \hat{J}_1 \hat{J}_2 \hat{K} \begin{Bmatrix} j_1 & j_2 & K \\ l_2 & l_1 & 1/2 \end{Bmatrix} \\ & \times \frac{(-1)}{\hat{K}} {}_{SD} \langle A\lambda_f J_f \parallel (a_{n_1 l_1 j_1}^\dagger \tilde{a}_{n_2 l_2 j_2})^{(K)} \parallel A\lambda_i J_i \rangle_{SD} \end{aligned}$$

All angular dependence
factorized out for plotting

NN and 3N interactions – $N^4\text{LO}(500)+3\text{N}^2\text{LO}$



6

NN systematic from LO to $N^4\text{LO}$

- D. R. Entem, N. Kaiser, R. Machleidt, and Y. Nosyk, Phys. Rev. C 91, 014002 (2015)
- D. R. Entem, R. Machleidt, and Y. Nosyk, arXiv:1703.05454



3N at $N^2\text{LO}$

- Navrátil, 650 MeV local cut-off and 500 MeV non-local cut-off




Density of ground state ${}^4\text{He}$, ${}^{12}\text{C}$, ${}^{16}\text{O}$ with NN- $\text{N}^4\text{LO}(500)+3\text{Nlnl}$


7



Local density




Applications to optical potentials




- We can achieve an additional step towards consistent optical potential calculations by using the nonlocal density

$$U(\vec{q}, \vec{K}) = \sum_{N=n,p} \int d\vec{P} \quad \eta(\vec{q}, \vec{K}, \vec{P}) t_{pN}(\vec{q}, \vec{K}, \vec{P}) \rho_N(\vec{q}, \vec{P})$$


Differential cross sections



Applications to optical potentials



Differential cross sections



Applications to density functional theory

10


- Kinetic density (and other related densities) is a DFT quantity we are capable of calculating from *ab initio* wavefunctions according to

$$\tau_T(\vec{r}) = \left(\vec{\nabla} \cdot \vec{\nabla}' \rho_T(\vec{r}, \vec{r}') \right) |_{\vec{r}=\vec{r}'}$$

$$\frac{d}{dr} R_{n,l}(r) = \frac{l}{r} R_{n,l}(r) - \frac{1}{b} \left[\sqrt{n+l+\frac{3}{2}} \cdot R_{n,l+1}(r) + \sqrt{n} \cdot R_{n-1,l+1}(r) \right]$$

- Effects of COM removal should be amplified in DFT quantities such as the kinetic density, due to the application of gradients on the nonlocal density

Kinetic density



Conclusions and outlook

11

- Conclusions
 - We observed significant differences in the nuclear density of light systems when the COM was removed
 - We can now use the more general, nonlocal density for optical potentials of nuclear reactions and *ab initio* calculations in DFT
 - More details on some of these results can be found in arXiv:1712.02879; Phys. Rev. C, in press.
- Outlook
 - We are now pursuing the use of these densities in calculations with natural orbitals (arxiv:1605.04976), reducing basis sizes and improving convergence
 - We will attempt to extend this to the translationally invariant one-body nuclear density matrix and further cut down basis sizes

Thank you
Merci

www.triumf.ca

Follow us @TRIUMFLab

