

RESEARCH STATEMENT

MATHILDE GERBELLI-GAUTHIER

I work in number theory and representation theory, with a focus on automorphic representations, functoriality in the Langlands program, and the cohomology of arithmetic groups. I am especially motivated by problems in which automorphic forms interact with other areas of mathematics, for example:

Topology: In my thesis [GG23] and subsequent works [DGG22, EGGG23], we bound the dimensions of the i^{th} cohomology of arithmetic manifolds, which, roughly speaking, measure the number of i -dimensional holes in the manifold, as they vary in sequences of covering spaces known as towers. A sample theorem is:

Theorem 1 ([GG23]). *Let p be a large enough prime, N be odd, and $X(p^n)$ be a tower of full-level compact arithmetic manifolds associated to $U(N-a, a)$. In the smallest nontrivial degree of cohomology a , we have:*

$$\dim H^a(X(p^n), \mathbf{Q}) \ll \text{Vol}(X(p^n))^{\frac{N}{N^2-1}}.$$

We derive such results from asymptotic bounds on automorphic representations, thereby proving several instances of the 30-year old Conjecture 1 quantifying the failure of the naïve Ramanujan conjecture.

Fourier analysis: In the collaboration [GGV], we give a new proof, using the cohomology of arithmetic groups, of the following Fourier interpolation formula of Radchenko-Viazovska:

Theorem 2 ([RV19]). *Let f be an even Schwartz function on the real line, and \hat{f} its Fourier transform. The function f is entirely determined by the values $(f(\sqrt{n}), \hat{f}(\sqrt{n}))_{n \in \mathbf{N}}$.*

The starting point for this strategy is the observation that the space \mathcal{S} of Schwartz functions carries the Weil representation of the metaplectic group. We deduce Theorem 2 from the vanishing of a cohomology group $H^1(\Gamma, \mathcal{S}^*)$ for a certain Γ , which we prove using the spectral theory of automorphic forms.

Outline. Section 1 discusses the relation between the cohomology of arithmetic groups and automorphic forms, and lays out recent progress and future directions in the program to use Langlands functoriality to understand their asymptotics. Section 2 concerns cohomology of arithmetic groups with infinite-dimensional coefficients, and applications to interpolation theorems. Finally, Section 3 describes two recent collaborative projects in other directions: one on mod ℓ gamma factors, and one on statistics of Fourier coefficients.

1. GROWTH OF COHOMOLOGY AND NON-TEMPERED REPRESENTATIONS

A classical measure of the topological complexity of a space X are the cohomology groups $H^i(X, \mathbf{C})$. These groups are studied geometrically using simplices or differential forms, but also algebraically as the *group cohomology* $H^i(\pi_1(X), \mathbf{C})$ of the so-called fundamental group $\pi_1(X)$ of the space X .

An especially rich class of spaces are *arithmetic manifolds* X_Γ , whose fundamental groups Γ are arithmetic groups, i.e. the integer entries $G(\mathbf{Z})$ of a (reductive) matrix group G such as GL_N . Some key properties are:

- The spaces X_Γ often parameterize objects of interest, such as quadratic forms or abelian varieties.
- The fundamental groups Γ have natural sequences of *congruence subgroups* $\Gamma(n)$, consisting of matrices congruent to the identity modulo n . The spaces $X_{\Gamma(n)}$ are *covering spaces* of X_Γ .
- Natural families of operators T_p act on $H^i(\Gamma, \mathbf{C})$ and their eigenvalues are expected to be connected to Galois representations. As such, the cohomology is the generalization of *modular forms*.

The cohomology of arithmetic manifolds, though well-studied, is in general very hard to compute. A more tractable problem is to consider the question asymptotically.

Question 1. In a tower $X_{\Gamma(n)}$ of arithmetic manifolds, how does $H^i(\Gamma(n), \mathbf{C})$ grow as a function of n ?

Arithmetic manifolds are Riemannian: their cohomology can be computed using harmonic differential forms, which carry $G(\mathbf{R})$ -representations. For X_Γ compact, Matsushima [Mat67] leveraged this to relate cohomology and representations:

$$(1) \quad H^*(\Gamma, \mathbf{C}) = \oplus_\pi m(\pi, \Gamma) H^*(\mathfrak{g}, K; \pi).$$

This equality expresses the cohomology in terms of the multiplicity $m(\pi, \Gamma)$ in $L^2(\Gamma \backslash G)$ of the irreducible *cohomological* unitary representations π of $G(\mathbf{R})$ (by definition, those such that $H^*(\mathfrak{g}, K; \pi) \neq 0$). A given group G has only finitely many cohomological representations [VZ84], allowing us to rephrase Question 1.

Question 2. Given a representation π of $G(\mathbf{R})$, how does $m(\pi, n) := m(\pi, \Gamma(n))$ grow as a function of n ?

First progress towards this question was made by deGeorge-Wallach [dGW78], who compared $m(\pi, n)$ to the volume of the arithmetic manifolds $X_{\Gamma(n)}$:

$$(2) \quad \lim_{n \rightarrow \infty} \frac{m(\pi, n)}{\text{Vol}(X_{\Gamma(n)})} = \begin{cases} k(\pi) > 0 & \text{if } \pi \hookrightarrow L^2(G(\mathbf{R})) \\ 0 & \text{otherwise.} \end{cases}$$

In particular, they show that the multiplicity $m(\pi, n)$ is always $O(\text{Vol}(X_{\Gamma(n)}))$, or, equivalently, $O(n^{\dim G})$. Additionally, integrability properties of π dictate two regimes for the rate of growth. Building on this second fact, Sarnak-Xue [SX91] conjectured growth rates for unitary representations $\pi \not\hookrightarrow L^2(G(\mathbf{R}))$:

Conjecture 1 (Sarnak-Xue). Let $p(\pi) = \inf\{p \geq 2 \mid \pi \hookrightarrow L^p(G(\mathbf{R}))\}$. Then

$$m(\pi, n) \ll_\epsilon \text{Vol}(X_{\Gamma(n)})^{2/p(\pi)+\epsilon}.$$

Representations such that $p(\pi) > 2$ are said to be *non-tempered*. To discuss them, we adopt a global perspective and consider automorphic representations, i.e. irreducible $G(\mathbf{A})$ -subrepresentations of

$$(3) \quad L^2(G(\mathbf{Q}) \backslash G(\mathbf{A})) = \oplus_\pi m(\pi) \pi$$

where \mathbf{A} are the adèles of \mathbf{Q} . Automorphic representations factor as tensor products $\pi = \otimes'_v \pi_v$ of representations of the local groups $G(\mathbf{Q}_v)$; to underscore our shift in perspective, representations of the Lie group $G(\mathbf{R})$ are now denoted π_∞ . The global and classical perspectives are related since up to finite index,

$$\Gamma(n) \backslash G(\mathbf{R}) = G(\mathbf{Q}) \backslash G(\mathbf{A}) / K_f(n)$$

for suitable subgroups $K_f(n) \subset G(\mathbf{A})$. The multiplicity counts of Question 2 can then be rewritten by considering the finite part $\pi_f := \otimes'_{v < \infty} \pi_v$ of the representations, and studying their fixed vectors $\pi_f^{K_f(n)}$. For π_∞^0 a fixed $G(\mathbf{R})$ -representation, the multiplicity is then $m(\pi_\infty^0, n) := \sum_{\pi = \pi_\infty^0 \otimes \pi_f} m(\pi) \dim \pi_f^{K_f(n)}$.

Arithmetically, the richest part of the decomposition (3) and of generalizations to non-compact quotients is the *cuspidal* part, which is the subject a central conjecture in the analytic theory of automorphic forms:

Conjecture 2 (Ramanujan-Petersson). If G is split and π is cuspidal, the constituents π_v of π are tempered.

This conjecture has analytic implications in number theory and beyond [Li20]. Yet for general G , it fails: the first counterexamples appeared in [Kur78]. But not all is lost: Conjecture 1 can be rephrased as

The Ramanujan-Petersson Conjecture holds asymptotically as $n \rightarrow \infty$.

1.1. Results. My research program aims to prove Conjecture 1 for cohomological representations using Langlands functoriality. The main idea is:

Multiplicities of non-tempered representations grow slowly because these representations come from smaller groups.

To be more precise, we rapidly outline the web of conjecture known as the *Langlands Program*. To any group G , it attaches a dual group \hat{G} . The *Reciprocity Conjecture* then predicts a correspondence

$$(4) \quad \{\text{Automorphic representations } \pi_\rho \text{ of } G\} \leftrightarrow \{\text{Representations } \rho_\pi \text{ of } \text{Gal}(\bar{\mathbf{Q}}/\mathbf{Q}) \text{ into } \hat{G}\}.$$

If a Galois representation ρ_π factors through a subgroup $\hat{H} \subset \hat{G}$, then (4) predicts a corresponding π_ρ^H for H , and according to the Langlands *Functoriality Conjecture* the characters of π_ρ^G and π_ρ^H should be related.

A well-understood instance of these relations is the Endoscopic Classification of Representation (ECR), proved by Arthur [Art13] for SO_N and Sp_N , and extended to unitary groups in [Mok15] and [KMSW14]. It describes the decomposition (3) in terms of the contributions of smaller, *endoscopic* groups.

In the footsteps of [MS19] who use endoscopy to prove Conjecture 1 for $U(N-1, 1)$, I prove in my thesis:

Theorem 3 ([GG19],[GG23]). *Let $U(N-a, a)$ be a unitary group of odd rank defined with respect to a CM extension E/F of number fields, let \mathfrak{p} be a prime of F , and let $\Gamma(\mathfrak{p}^n)$ be a sequence of cocompact congruence subgroups. Then for the smallest degree a whose cohomology does not identically vanish,*

$$\dim H^a(\Gamma(\mathfrak{p}^n), \mathbf{C}) \ll |\mathfrak{p}^n|^N,$$

where $|\cdot| : F \rightarrow \mathbf{Q}$ is the norm. In particular, Conjecture 1 holds for the corresponding representations.

The idea of the theorem is to not only appeal to the ECR, but also repurpose its tools, namely the *stabilization of the trace formula*. This is an expression for the multiplicity $m(\pi_\infty, \mathfrak{p}^n)$ in terms of a *stable* contribution of our main group G and of “error terms” coming from the smaller endoscopic groups H .

$$(5) \quad m(\pi_\infty, \mathfrak{p}^n) = St_{\pi_\infty}^G(\mathfrak{p}^n) + \sum_{H \text{ endoscopic}} St_{\pi_H}^H(\mathfrak{p}^n).$$

The main observation powering Theorem 3 is:

For good π_∞ , one of the “error terms” in (5) is in fact the dominant term as n grows.

This, combined with the first observation after (2), allows me to bound the multiplicities in terms of the smaller rate of growth $|\mathfrak{p}^n|^{\dim H}$. In subsequent joint work with Rahul Dalal, we vastly extend the results:

Theorem 4 ([DGG22]). *Let G be a rank N unitary group defined with respect to a CM extension E/F . For any cohomological π_∞ , there is $R(\pi_\infty)$ explicit such that for any sequence of ideals \mathfrak{n} split in E :*

- (i) $m(\pi_\infty, \mathfrak{n}) \ll |\mathfrak{n}|^{R(\pi_\infty)}$ and the exponent $R(\pi_\infty)$ proves Conjecture 1 if $N > 4$.
- (ii) Under some conditions on E and π_∞ , $R(\pi_\infty)$ is sharp. For example, if $G(\mathbf{R}) = U(N-a, a)$ with N odd and a the smallest degree whose cohomology does not identically vanish, then there exist sequences of cocompact lattices $\Gamma(\mathfrak{n})$ such that

$$H^a(\Gamma(\mathfrak{n}), \mathbf{C}) \asymp \text{Vol}(X_{\Gamma(\mathfrak{n})})^{\frac{N}{N^2-1}}.$$

In fact, we prove more refined statistics in [DGG22]: we weigh multiplicities by unramified functions and show that the finite factors π_v of representations such that π_∞ and E satisfies the assumptions of (ii) distribute on average according to a Sato-Tate measure predicted by Langlands functoriality.

The proof of Theorem 4 builds on that of Theorem 3, but tackles the much more delicate cases where the G -term in (5) dominates. Inspired by [Taï17], we then analyze this term by taking seriously the intuition:

Endoscopy is the echo for classical groups of parabolic induction happening on GL_N .

In practice, this means that the main work of Theorem 4 is controlling traces of representations obtained by *twisted parabolic induction*. We give upper bounds on these in general, and exact identities in some cases.

What about non-unitary groups? I begin to tackle this question with Shai Evra and Henrik Gustafsson:

Theorem 5 ([EGGG23]). *Let F be a totally real number field, and let $G = SO_5/F$ or an inner form unramified at the finite places.*

- (i) *Conjecture 1 holds for cohomological representations of G . In particular, if $\Gamma(\mathfrak{n})$ is a sequence of cocompact lattices in $G(\mathbf{R})$, then for the unique degree d of interest we have:*

$$\dim H^d(\Gamma(\mathfrak{n}), \mathbf{C}) \ll |\mathfrak{n}|^5, \quad d = \begin{cases} 2 & \text{if } G(\mathbf{R}) = SO(3, 2) \\ 1 & \text{if } G(\mathbf{R}) = SO(4, 1). \end{cases}$$

- (ii) *If the group G is compact at the infinite places, the analogue of Conjecture 1 for finite places v also holds: the multiplicity of non-tempered π_v are negligible as the level grows.*

The generalization of Conjecture 1 in (ii) is the *Sarnak-Xue Density Hypothesis*: its proof opens new applications in which arithmetic manifolds are replaced by discrete analogues such as simplicial complexes. As a corollary of Theorem 5, we exhibit infinite families of so-called Density Ramanujan graphs: regular graphs with optimal mixing properties, which are sought after for applications to computer science [Lub18].

Theorem 5 is also proved in the framework of functoriality, but adopts a mixed approach in which the multiplicity counts are split into a *global* and a *local* components, and both are estimated independently. The estimates rely on Schmidt's explicit description of the automorphic representations of SO_5 [Sch20].

1.2. Future Work. I plan to build on the above and give sharp asymptotics on the multiplicities of cohomological representations for all U_N , SO_N , and Sp_N . The current obstacles to fulfilling this program involve foundational questions of interest in their own right, which I plan on tackling in the coming years:

Asymptotics of root numbers in higher rank. The arithmetic of the ECR is encoded in Arthur's so-called ϵ -factors: certain sign characters obtained from root numbers of local L -functions associated to the automorphic representations. Even coarse control on their distribution as the level \mathfrak{n} grows would extend the sharp results of Theorem 4 to a much wider class of representations. With Rahul Dalal, our current strategy to compute these statistics uses the trace formula, and exploits the fact that the sign of the functional equation is equal to the eigenvalue of an Atkin-Lehner type operator on the space of newforms.

Character expansion of local representations. To generalize the local-global methods of Theorem 5 and to obtain sharp asymptotics, the key is to control the dimensions of fixed vectors in local representations π_p . As the level p^n grows, $\dim \pi_p^{K(p^n)}$ grows like $c(\pi_p)(p^n)^{\text{GKdim}}$, with GKdim the well-understood Gelfand-Kirillov dimension of π_p , but the $c(\pi_p)$ are mysterious. Bounding these constants is a tractable problem for induced representations by Mackey theory, so the work lies in the study of the building blocks: cuspidal representations. For this, a promising strategy is to rely on the recent developments [Fin21] showing that all cuspidal representations are constructed as compact-inductions of representations of finite groups.

Twisted parabolic induction. The restrictions in part (ii) of Theorem 4, as well as the obstacle to extending its methods to SO_N and Sp_N , come from difficulties in comparing twisted traces on GL_N with those on some of its Levi subgroups. The identities come in two categories: *spectral stable transfer*, in which one compares representations of GL_N to those of a product $GL_{N_1} \times \dots \times GL_{N_r}$ for $\sum N_r = N$, and *Speh transfer*, which compares non-tempered representations on GL_{Nd} to tempered representations on $(GL_N)^d$. Once again, we need only the coarsest asymptotic information to derive results.

2. COHOMOLOGY OF ARITHMETIC GROUPS AND INTERPOLATION

Recently, as part of the series of works leading up to the sphere-packing results for the E_8 and Leech lattices, Radchenko–Viazovska proved the following interpolation theorem.

Theorem 6 ([RV19]). *Let f be an even Schwartz function on the real line, and \hat{f} its Fourier transform. The function f is entirely determined by the values $(f(\sqrt{n}), \hat{f}(\sqrt{n}))_{n \in \mathbb{N}}$, and the only relation comes from the Poisson summation formula.*

2.1. Results. In joint work with Akshay Venkatesh, we give a new proof of this result in the language of the cohomology of arithmetic groups. The main insight driving this proof is:

The space \mathcal{S} of Schwartz functions carries the Weil representation of the metaplectic group.

The metaplectic group is the degree 2 cover of SL_2 , and the Weil representation is defined by:

- $\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ multiplies $f(x)$ by $e^{t\pi i x^2}$ and $\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}$ multiplies $\hat{f}(x)$ by $e^{t\pi i x^2}$, and
- the lift of $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ acts by the Fourier transform.

The extension of these formulas to the dual \mathcal{S}^* of the Weil representation, together with a Mayer-Vietoris-type argument, allow us to reduce Theorem 6 to

$$(6) \quad H^0(\Gamma, \mathcal{S}^*) = \mathbb{C}, \quad \text{and} \quad H^1(\Gamma, \mathcal{S}^*) = 0, \quad \text{for} \quad \Gamma = \langle \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \rangle \subset SL_2(\mathbb{Z}).$$

To compute the cohomology groups in (6), we extend the framework of Section 1 relating group cohomology and automorphic representations: the main difference is that the groups

$$H^*(\Gamma, \mathbb{C}) \quad \text{are replaced by} \quad H^*(\Gamma, V)$$

where the coefficient system is no longer trivial, or finite dimensional, but is an infinite-dimensional topological vector space V . For these representations and when the ambient group G is a cover of SL_2 , we extend a result of Bunke-Olbrich [BO98] and prove a Matsushima's formula (1) for infinite-dimensional coefficients.

Theorem 7 ([GGV]). *Let Γ be a lattice in a covering group of $SL_2(\mathbf{R})$, and let W be an infinite-dimensional irreducible smooth Fréchet representation of G with infinitesimal character λ . Then there is a unique complementary representation W^{cl} such that*

$$H^0(\Gamma, W^*) = m(\Gamma, W), \quad H^1(\Gamma, W^*) = m_{\text{cusp}}(\Gamma, W^{cl}),$$

where $m_{\text{cusp}}(\Gamma, V)$ denotes the multiplicity of V in the cuspidal spectrum at level Γ .

We give two proofs of this: one based on [BO98] using resolutions, and one, in the spirit of Matsushima, using (\mathfrak{g}, K) cohomology. The crux in both cases is the surjectivity of a differential operator on the space of automorphic forms, which we prove using the spectral decomposition and a truncation argument.

Specialization of this theorem to $W = \mathcal{S}$ gives the desired dimensions from (6) and allows us to deduce Theorem 6. Moreover, changing the coefficients to other spaces of functions gives new interpolation theorems:

Theorem 8 ([GGV]). *Let $\mathbf{P}^1(\mathbf{R})$ be the real projective line, with homogenous coordinates $[x : y]$. Then any $\Phi \in C^\infty(\mathbf{P}^1(\mathbf{R}))$ is entirely determined the following family of integrals:*

$$a_n(\Phi) = \int_{\mathbf{R}} \Phi(x, 1) e^{\pi i n x} dx, \quad b_n(\Phi) = \int_{\mathbf{R}} \Phi(1, y) e^{\pi i n y} dy.$$

2.2. Future Work. This work has natural extensions in several directions: we would like to prove the technical Theorem 7 for more degrees of cohomology, and more general class of groups. Additionally, it would be interesting to systematize the relation between cohomology vanishing and interpolation theorems.

Another line of work would move away from the archimedean place towards the p -adic places: can one deduce p -adic interpolation theorems from computing cohomology with values spaces of functions on p -adic manifolds? In this context, it is likely that the relevant lattices would be p -arithmetic groups: finite covolume subgroups of $SL_2(\mathbf{R}) \times SL_2(\mathbf{Q}_p)$.

3. GAMMA FACTORS AND L -FUNCTIONS

To conclude, I discuss two collaborative projects [BGGG⁺23] and [FGGHT23] on invariants of automorphic representations, namely gamma factors and L -functions.

3.1. Gamma factors and a converse theorem mod ℓ . We consider the classical problem of classifying the irreducible representations of $GL_n(\mathbf{F}_q)$. We study *gamma factors* $\gamma(\rho, \pi)$, invariants introduced by Piatetski-Shapiro for complex representations ρ of $GL_2(\mathbf{F}_q)$, with π a character of $GL_1(\mathbf{F}_q)$. The construction of $\gamma(\rho, \pi)$ is harmonic-analytic and relies on the realization of ρ as a space of functions, its *Whittaker Model*. Piatetski-Shapiro then proves:

Theorem 9 ([PS83]). *A \mathbf{C} -valued cuspidal representation of $GL_2(\mathbf{F}_q)$ is determined by its gamma factors.*

This statement is a *converse theorem*: such results are motivated by local Langlands correspondences, which are characterized by equalities of gamma factors and L -functions. Macdonald [Mac80] proved such a correspondence for representations of $GL_n(\mathbf{F}_q)$, associating them to equivalence classes of Weil group representations. Moreover, representations of $GL_n(\mathbf{F}_q)$ give rise to depth-zero representations of the corresponding p -adic GL_n , and [Mac80] is compatible with the usual local Langlands in depth zero.

3.1.1. Results. In [BGGG⁺23], we consider coefficients in any characteristic $\ell \nmid q$; in particular, ℓ may divide $|GL_n(\mathbf{F}_q)|$. In this situation, Vigneras [Vig88] has extended many results from the complex case. Using these, we first construct mod ℓ gamma factors. Then, exploiting their explicit description as Gauss sums in the case of $GL_2(\mathbf{F}_q)$, we compute counterexamples to the converse theorem on SAGE.

Example 10. When $(\ell, q) = (2, 5), (3, 7), (11, 23)$, or $(29, 59)$, there are non-isomorphic mod ℓ cuspidal representations of $GL_2(\mathbf{F}_q)$ whose gamma factors agree.

This suggests that for mod ℓ representations, one needs to modify the naïve construction. We do this by replacing the Whittaker model of ρ valued in the constants $\bar{\mathbf{F}}_\ell$ by its analogue consisting of functions valued in an Artinian algebra $R(\pi)$ of endomorphisms associated to π . With this new gamma factor, we find:

Theorem 11 ([BGGG⁺23]). *Let ρ be a mod ℓ cuspidal representation of $GL_n(\mathbf{F}_q)$. Then ρ is entirely determined by its $R(\pi)$ -valued γ factors $\gamma(\rho, \pi)$, as π runs over cuspidal representations of $GL_{n-1}(\mathbf{F}_q)$.*

The crux of the proof is establishing a completeness of $R(\pi)$ -valued Whittaker models, which generalizes linear independence of characters: we do this by analyzing ρ and π via their Bernstein-Zelevinsky derivatives.

3.1.2. Future Work.

Jacquet's conjecture mod ℓ . Our converse theorem concerns gamma factors $\gamma(\rho, \pi)$ against representations of $GL_{n-1}(\mathbf{F}_q)$. It was conjectured by Jacquet, and proved [Nie14] for \mathbf{C} coefficients, that $\gamma(\rho, \pi)$ determine ρ for representations π of $GL_m(\mathbf{F}_q)$ with $m \leq \lfloor \frac{n}{2} \rfloor$. We expect to extend our methods to prove this.

Characterization of Langlands-type correspondances. The correspondence [Mac80] preserves Godement-Jacquet gamma factors, a different construction. Vigneras [Vig94] built a correspondence mod ℓ , and recently, Li-Shotton [LS23] proved a version in families. We hope that they are characterized by equality of our (Rankin-Selberg) gamma factors: the first step is showing that $\gamma(\rho, \pi)$ are multiplicative over parabolic induction.

3.2. Signs in Fourier coefficients of modular forms and zeros of the L-function. This project concerns classical modular Hecke eigenforms f of weight k , given by a Fourier expansion

$$f(z) = \sum_{n=1}^{\infty} a_n(f) n^{\frac{k-1}{2}} e^{\pi i n z},$$

with z in the complex upper half-plane. The Fourier coefficients $a_n(f)$ are multiplicative and the Ramanujan conjecture for SL_2 , proved by Deligne [Del74] for modular forms of weight $k \geq 2$, states that

$$(7) \quad |a_n(f)| \leq \tau(n),$$

for $\tau(n)$ the number of positive divisors of n . We study cancellation between the $a_n(f)$, i.e. of the growth of

$$S(f, x) := \sum_{n \leq x} a_n(f).$$

The best bounds independent of f were proved by Lamzouri [Lam19]: under the GRH for the L -function $L(s, f)$, he showed that $S(f, x) = o(x \log x)$ as $\frac{\log x}{\log \log k} \rightarrow \infty$.

3.2.1. Results. The work [FGGHT23] with Claire Frechette, Alia Hamieh, and Naomi Tanabe, shows that the sums $S(f, x)$ are $o(x \log x)$ for $x \geq k^\epsilon$, under assumptions both weaker and more explicit than GRH:

Theorem 12 ([FGGHT23]). *Let $f \in S_k(N)$ be a primitive cusp form. Let ϵ and T be real numbers with $\epsilon \geq (\log k)^{-1/8}$ and $1 \leq T \leq (\log k)^{1/200}$. Suppose that the region*

$$R(T) = \left\{ s : \Re(s) \geq \frac{3}{4}, |\Im(s)| \leq T + \frac{1}{4} \right\}$$

contains no more than $\epsilon^2 \log k / 5000$ zeroes of $L(s, f)$. Then for all $x \geq k^\epsilon$, we have $|S(f, x)| \ll (x \log x) / T$.

Our proof relies on recent developments in analytic number theory pioneered by Granville–Soundararajan, whose analogous result [GS18] on character sums was the inspiration for the project. The framework, known as *pretentious number theory*, applies naturally to bounded functions. Recently the key input, Halazs' Theorem, was extended by Mangerel [Man23] to multiplicative functions satisfying (7).

The strategy is to compare $a_f(n)$ and n^{it} for $t \in \mathbf{R}$: the latter see very little cancellation, and for good t the twisted sums $S(f, x, t) = \sum_{n \leq x} a_f(n) n^{it}$ remain close enough to $S(f, x)$. This is useful because:

$S(f, x, t)$ and t -translates of the L-function $L(s, f)$ are related by the Fourier transform.

This allows us to use Plancherel's formula to compare $S(f, x)$ and a suitable shift $L(s + it, f)$. By assuming for contradiction that $S(f, x)$ is large, we first produce a lower bound involving $L(s + it, f)$, which we express in terms of its zeros using Hadamard's formula to obtain our result.

3.2.2. Future Work. A long-term goal using the methods of our theorem is to extend the strategy to the cuspidal representations of GL_n/\mathbf{Q} which satisfy the Ramanujan-Petersson Conjecture 2. A first project to test the robustness of the method is to bound sums of Rankin-Selberg products of modular forms.

REFERENCES

- [Art13] James Arthur. *The endoscopic classification of representations: Orthogonal and symplectic groups*, volume 61 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2013.
- [BGGG⁺23] Jacksyn Bakeberg, Mathilde Gerbelli-Gauthier, Heidi Goodson, Ashwin Iyengar, Gilbert Moss, and Robin Zhang. Mod ℓ gamma factors and a converse theorem for finite general linear groups. *arxiv:2307.07593*, 2023.
- [BO98] Ulrich Bunke and Martin Olbrich. Resolutions of distribution globalizations of Harish-Chandra modules and cohomology. *J. Reine Angew. Math.*, 497:47–81, 1998.
- [Del74] Pierre Deligne. La conjecture de Weil. I. *Inst. Hautes Études Sci. Publ. Math.*, (43):273–307, 1974.
- [DGG22] Rahul Dalal and Mathilde Gerbelli-Gauthier. Statistics of cohomological automorphic representations via the endoscopic classification. *arXiv:2212.12138*, 2022.
- [dGW78] David L. de George and Nolan R. Wallach. Limit formulas for multiplicities in $L^2(\Gamma \backslash G)$. *Ann. of Math. (2)*, 107(1):133–150, 1978.
- [EGGG23] Shai Evra, Mathilde Gerbelli-Gauthier, and Henrik P. A. Gustafsson. The cohomological sarnak-xue density hypothesis for so_5 . *arXiv:2309.12413*, 2023.
- [FGGHT23] Claire Frechette, Mathilde Gerbelli-Gauthier, Alia Hamieh, and Naomi Tanabe. Large sums of fourier coefficients of cusp forms. *arxiv:2308.06311*, 2023.
- [Fin21] Jessica Fintzen. Types for tame p -adic groups. *Ann. of Math. (2)*, 193(1):303–346, 2021.
- [GG19] Mathilde Gerbelli-Gauthier. Growth of cohomology of arithmetic groups and the stable trace formula: the case of $u(2, 1)$. *arXiv:1910.06900*, 2019.
- [GG23] Mathilde Gerbelli-Gauthier. Limit multiplicity for unitary groups and the stable trace formula. *Algebra & Number Theory*, 17(12):2181–2228, 2023.
- [GGV] Mathilde Gerbelli-Gauthier and Akshay Venkatesh. Cohomology of fuchsian groups and fourier interpolation. *In preparation*.
- [GS18] Andrew Granville and Kannan Soundararajan. Large character sums: Burgess’s theorem and zeros of L -functions. *J. Eur. Math. Soc. (JEMS)*, 20(1):1–14, 2018.
- [KMSW14] Tasho Kaletha, Alberto Minguez, Sug Woo Shin, and Paul-James White. Endoscopic classification of representations: inner forms of unitary groups. *arXiv preprint arXiv:1409.3731*, 2014.
- [Kur78] Nobushige Kurokawa. Examples of eigenvalues of Hecke operators on Siegel cusp forms of degree two. *Invent. Math.*, 49(2):149–165, 1978.
- [Lam19] Youness Lamzouri. Large sums of Hecke eigenvalues of holomorphic cusp forms. *Forum Math.*, 31(2):403–417, 2019.
- [Li20] Wen-Ching Winnie Li. The Ramanujan conjecture and its applications. *Philos. Trans. Roy. Soc. A*, 378(2163):20180441, 14, 2020.
- [LS23] Tzu-Jan Li and Jack Shotton. On endomorphism algebras of gelfand–graev representations II. *Bulletin of the London Mathematical Society*, aug 2023.
- [Lub18] Alexander Lubotzky. High dimensional expanders. In *Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018*, pages 705–730. World Scientific, 2018.
- [Mac80] I. G. Macdonald. Zeta functions attached to finite general linear groups. *Math. Ann.*, 249(1):1–15, 1980.
- [Man23] Alexander P. Mangerel. Divisor-bounded multiplicative functions in short intervals. *Res. Math. Sci.*, 10(1):Paper No. 12, 47, 2023.
- [Mat67] Yozô Matsushima. A formula for the betti numbers of compact locally symmetric riemannian manifolds. *Journal of Differential Geometry*, 1(1-2):99–109, 1967.
- [Mok15] Chung Pang Mok. Endoscopic classification of representations of quasi-split unitary groups. *Mem. Amer. Math. Soc.*, 235(1108):vi+248, 2015.
- [MS19] Simon Marshall and Sug Woo Shin. Endoscopy and cohomology in a tower of congruence manifolds for. In *Forum of Mathematics, Sigma*, volume 7, page e19. Cambridge University Press, 2019.
- [Nie14] Chufeng Nien. A proof of the finite field analogue of jacquet’s conjecture. *American Journal of Mathematics*, 136(3):653–674, 2014.
- [PS83] Ilya Piatetski-Shapiro. *Complex representations of $GL(2, K)$ for finite fields K* , volume 16. American Mathematical Soc., 1983.
- [RV19] Danylo Radchenko and Maryna Viazovska. Fourier interpolation on the real line. *Publications mathématiques de l’IHÉS*, 129(1):51–81, 2019.
- [Sch20] Ralf Schmidt. Paramodular forms in cap representations of $\mathrm{gsp}(4)$. *Acta Arith*, 194(4):319–340, 2020.
- [SX91] Peter Sarnak and Xiao Xi Xue. Bounds for multiplicities of automorphic representations. *Duke Math. J.*, 64(1):207–227, 1991.
- [Taï17] Olivier Taïbi. Dimensions of spaces of level one automorphic forms for split classical groups using the trace formula. In *Annales Scientifiques de l’école normale supérieure*, volume 50, pages 269–344, 2017.
- [Vig88] Marie-France Vignéras. Représentations modulaires de $GL(2, F)$ en caractéristique l , F corps fini de caractéristique $p \neq l$. *C. R. Acad. Sci. Paris Sér. I Math.*, 306(11):451–454, 1988.
- [Vig94] Marie-France Vignéras. Sur la conjecture locale de Langlands pour $GL(n, F)$ sur $\overline{\mathbf{F}}_l$. *C. R. Acad. Sci. Paris Sér. I Math.*, 318(10):905–908, 1994.
- [VZ84] David A. Vogan, Jr. and Gregg J. Zuckerman. Unitary representations with nonzero cohomology. *Compositio Math.*, 53(1):51–90, 1984.