
CAPACITY
a
EAC + EDC (mmol e⁻ g⁻¹)
×1.92
×1.25
×1.00
Plaque
Rhizo-sphere
Bulk soil
b
Poorly crystalline Fe (%)
75%
32%
30%
Plaque
Rhizo-sphere
Bulk soil
c
EEC relative to goethite
×1.62
Goethite
After 30 d cycling
d
Share of reactive Fe (%)
Mineral
Transition
Organic
Palsa
Bog
Fen
CONNECTIVITY
e
Fe(II) / compartment mean
Plaque
Rhizosphere
Bulk soil
Time (h)
f
Relative to plaque
Fe(II) variation
Available-P variation
Assay capacity (EAC + EDC)
0.52
0.52
0.40
0.14
0.08
Plaque (root surface)
Rhizosphere (< 4 mm)
Bulk soil (> 2 cm)
KINETICS
g
N₂O amount × time, 1–24 h (µmol N h)
Ox
SA
LA
ρ = 0.37
Terminal-record N₂ half-time (h)
h
Treated / untreated CO₂ (%)
γ-sterilized
TBA-treated
Paddy soil
Sandy soil
0.5–6 h
0.5–48 h
Integration window after rewetting
MEMORY
i
N₂O (µmol N per vial)
Ox
SA
LA
Time in common challenge (h)
j
Ratio to Ox mean
biomass and terminal yield
accumulation
SA/Ox
LA/Ox
PLFA
Terminal N₂
N₂O, 1–24 h
k
N₂O amount × time / Ox
SA/Ox
LA/Ox
1.19
0.48
Window end W (h), from 1 h