

THE EXTRAPOLATION FOR BOUNDARY FINITE ELEMENTS

BY

LI KAITAI

AND

YAN NINGNING

IMA Preprint Series # 287

January 1987

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS

UNIVERSITY OF MINNESOTA

514 Vincent Hall

206 Church Street S.E.

Minneapolis, Minnesota 55455

#	Author(s)	Title	Recent IMA Preprints (continued)		
#	Author(s)	Title	#	Author(s)	Title
165	D. Arnold and R.S. Falk,	Continuous Dependence on the Elastic Coefficients for a Class of Anisotropic Materials	203	G. Buttazzo, G. Dal Maso and U. Mosco,	A Derivation Theorem for Capacities with Respect to a Radon Measure
166	I.J. Bakelman,	The Boundary Value Problems for Non-linear Elliptic Equation and the Maximum Principle for Euler-Lagrange Equations	204	S. Cowin, M. Mehrabadi,	On the Identification of Material Symmetry for Anisotropic Elastic Materials
167	Ingo Müller,	Gases and Rubbers	205	R.W.R. Darling,	Constructing Nonhomeomorphic Stochastic Flows.
168	Ingo Müller,	Pseudoelasticity in Shape Memory Alloys - an Extreme Case of Thermoelasticity	206	M. Chipot,	On the Reynolds Lubrication Equation
169	Luis Magalhães,	Persistence and Smoothness of Hyperbolic Invariant Manifolds for Functional Differential Equations	207	R.V. Kohn and G.W. Milton,	On Bounding the Effective Conductivity of Anisotropic Composites
170	A. Damiliani and M. Vogelius,	Homogenization Limits of the Equations of Elasticity in Thin Domains	208	I.J. Bakelman,	Notes Concerning the Torsion of Hardening Rods and Its N-Dimensional Generalizations
171	H.C. Simpson and S.J. Spector,	On Hadamard Stability in Finite Elasticity	209	I.J. Bakelman,	The Boundary Value Problems for Non-linear Elliptic Equations!
172	J.L. Vazquez and C. Verru,	Isolated Singularities of the Solutions of the Schrödinger Equation with a Radial Potential	210	Guanglu Gong & Mingping Qian,	On the Large Deviation Functions of Markov Chain
173	G. Dal Maso and U. Mosco,	Wiener's Criterion and L^1 -Convergence	211	Arle Leizarowitz,	Control Problems with Random and Progressively Known Target
174	John H. Maddocks,	Stability and Folds	212	R.W.R. Darling,	Ergodicity of a Measure-Valued Markov Chain Induced by Random Transformations
175	R. Hardt and D. Kinderlehrer,	Existence and Partial Regularity of Static Liquid Crystal Configurations	213	G. Gong, M. Qian & Zhongxin Zhao,	Killed Diffusions and its Conditioning
176	M. Neruker,	Construction of Smooth Ergodic Cycles for Systems with Fast Periodic Approximations	214	Arle Leizarowitz,	Controlling Diffusion Processes on Infinite Horizon with the Overtaking Criterion
177	J.L. Ericksen,	Stable Equilibrium Configurations of Elastic Crystals	215	Milard Beatty,	The Poisson Function of Finite Elasticity
178	Patricio Aviles,	Local Behavior of Solutions of Some Elliptic Equations	216	David Ternan,	Travelling Wave Solutions Arising From a Combustion Model
179	S.-N. Chow and R. Lauterbach,	A Bifurcation Theorem for Critical Points of Variational Problems	217	Yuh-Jia Lee,	Sharp Inequalities and Regularity of Heat Semi-Group on Infinite Dimensional Spaces
180	R. Pego,	Phase Transitions: Stability and Admissibility in One Dimensional Nonlinear Viscoelasticity	218	D. Stroock,	Lecture Notes
181	Mariano Giaquinta,	Quadratic Functions and Partial Regularity	219	Claudio Csanuto,	Spectral Methods and Maximum Principle
182	J. Bona,	Fully Discrete Galerkin Methods for the Korteweg De Vries Equation	220	Thomas O'Brien,	A Two Parameter Family of Pension Contribution Functions and Stochastic Optimization
183	J. Maddocks and J. Keller,	Mechanics of Robes	221	Takeyuki Hida,	Analysis of Brownian Functionals
184	F. Bernis,	Qualitative Properties for some nonlinear higher order Nonlinear Higher Order Parabolic Equations with Absorption	222	Leonid Murwicz,	On Informational Decentralization and Efficiency of Resource Allocation Mechanisms
185	F. Bernis,	Finite Speed of Propagation and Asymptotic Rates for some Nonlinear Higher Order Parabolic Equations with Minimal Strategy Spaces	223	E.B. Fabes and D.W. Stroock,	A New Proof of Moser's Parabolic Harnack Inequality via the Old Ideas of Nash
186	S. Reichelstein and S. Reiter,	Game Forms with Minimal Strategy Spaces	224	Minoru Murata,	Structure of Positive Solution to $(-\Delta + V)u = 0$ in R^n
187	An Answer to Littlewood's Problem on Boundedness		225	Paul Dupuis,	Large Deviations Analysis of Reflected Diffusion Constrained Stochastic Approximation Algorithms in Convex Sets
188	J. Rubinstein and R. Maurl,	Dispersion and Convection in Periodic Media	226	F. Bernis,	Existence Results for Doubly Nonlinear Higher Order Parabolic Equations on Unbounded Domains.
189	W.H. Fleming and P.E. Souganidis,	Asymptotic Series and the Method of Vanishing Viscosity	227	S. Orey and S. Pelikan,	Large Deviations Principles for Stationary Processes of Minimal Surfaces.
190	H. Beirão Da Veiga,	Existence and Asymptotic Behavior for Strong Solutions of Navier-Stokes Equations in the Whole Space	228	R. Gulliver and S. Hildebrandt,	Boundary Configurations Spanning Continua
191	L.A. Caffarelli, J.L. Vazquez, and N.I. Wolanski,	Lipschitz Continuity of Solutions and Interfaces of the N-Dimensional Porous Medium Equation	229	J. Baxter, G. Dal Maso & U. Mosco,	Stopping Times and T-Convergence.
192	R. Johnson,	m-Functions and Floquet Exponents for Linear Differential Systems	230	Juifio Bouillet,	Self-Similar Solutions, Having Jumps and Intervals of Constancy of a Diffusion-heat Conduction Equation
193	F.V. Atkinson and L.A. Peletier,	Ground States and Dirichlet Problems for $-\Delta = F(U)$ in R^n	231	R. Hardt, D. Kinderlehrer & F.-H. Lin,	A Remark About the Stability of Smooth Equilibrium Configurations of Static Liquid Crystal
194	G. Dal Maso, U. Mosco,	The Wiener Modulus of a Radial Measure	232	M. Chipot and M. Luskin,	The Compressible Reynolds Lubrication Equation
195	H. A. Levine and H.F. Weinberger,	Inequalities between Dirichlet and Neumann Eigenvalues	233	J.H. Maddocks,	A Model for Disclinations in Nematic Liquid Crystal
196	J. Rubinstein,	On the Macroscopic Description of Slow Viscous Flow Past a Dirichlet Problem	234	C. Fofas, G.R. Seif and R. Temam,	Inertial Manifolds for Nonlinear Evolutionary Equations
197	G. Dal Maso and U. Mosco,	Wiener Criteria and Energy Decay for Relaxed Dirichlet Problems	235	P.L. Chow,	Expectation Functionals Associated with Some Stochastic Evolution
198	V. Olfker and P. Walfman,	On the Monge-Ampere Equation Arising in the Reflect Mapping Problem	236	Giuseppe Buttazzo,	Reinforcement by a Thin Layer with Oscillating Thickness
199	M. Chipot, D. Kinderlehrer and L. Caffarelli,	Some Smoothness Properties of Linear Laminaes	237	W.H. Fleming, S.J. Sheu and H.M. Soner,	On Existence of the Dominant Eigenfunction and its Application to the Large Deviation Properties of a
200	Y. Giga and R. Kohn,	Characterizing Blow-up Using Similarity Variables	238	R. Jensen and P.E. Souganidis,	A Regularity Result for Viscosity Solutions of Hamilton-Jacobi Equations in one Space Dimension
201	P. Cannarsa and H. M. Soner,	On the Singularities of the Viscosity Solutions to Hamilton-Jacobi-Bellman Equations	239	B. Boczar-Karakl, J.L. Bona and D.L. Cohen,	Interaction of Shallow-Water Waves and Bottom Topography
202	Andrew Majda,	Nonlinear Geometric Optics for Hyperbolic Systems of Conservation Laws	240	F. Colonius and W. Klemann,	Infinite Time Optimal Control and Periodicity Dirichlet Problems

#	Author(s)	Title	Recent IMA Preprints (continued)	#	Author(s)	Title
241	Harry Kesten,	Scaling Relations for 2D-Percolation	277 Cockburn, B.	The Quasi-Monotone Schemes for Scalar Conservation Laws Part II		
242	A. Leizarowitz,	Infinite Horizon Optimization for Markov Process with Finite States Spaces	278 M. Chipot, T. Sideris,	On the Abelian Higgs Model		
243	Louis H.-Y. Chen,	The Rate of Convergence in A Central Limit Theorem for Dependent Random Variables with Arbitrary Index Set	279 C. Foias, B. Nicolaenko, G.R.Sell, R. Temam,	Inertial Manifolds for the Kuramoto-Sivashinsky Equation and an Estimate of their Lowest Dimension		
244	G. Kaliljanpur,	Stochastic Differential Equations in Duals of Nuclear Spaces with some Applications	280 R. Duran,	On the Approximation of Miscible Displacement in Porous Media by Method of Characteristics Combined with a Mixed Method		
245	Tzuu-Shuh Chang,	Yunshong Chow and Yuh-Jia Lee, Evaluation of Certain Functional Integrals	281 H. Axläng, Zhang Bo ,	The Convergence for Nodal Expansion Method		
246	L. Karp and M. Pinsky,	The First Eigenvalue of a Small Geodesic Ball in a Riemannian Manifold	282 V. Twersky,	Dispersive Bulk Parameters for Coherent Propagation in Correlated Random Distributions		
247	Chl-Sing Man,	Towards An Acoustoelastic Theory for Measurement of Residual Stress	283 W.Th.F. den Hollander,	Mixing Properties for Random Walk in Random Scenery		
248	Andreas Stoll,	Invariance Principles for Brownian Intersection	284 H.R. Jauslin,	Nondifferentiable Potentials for Nonequilibrium Steady States		
249	R.W.R. Darling,	Time and Polymer Measures	285 K.Meyer, G.R.Sell,	Homoclinic Orbits and Bernoulli Bundles in Almost Periodic Systems		
250	R. Cohen, R. Hardt, D. Kinderlehrer, S.-Y.Lin, M. Luskin,	Minimum Energy for Liquid Crystals: Computational Results	286 J. Douglas,Jr., Y. Yuan,	Finite Difference Methods for the Transient Behavior of a Semiconductor Device		
251	Suzanne M. Lenhart,	Viscosity Solutions for Weakly Coupled Systems of First Order PDEs	287 Li Kaitai, Yan Ningling,	The Extrapolation for Boundary Finite Elements		
252	M. Cranston, E. Fabes, Z. Zhao,	Condition Gauge and Potential Theory for the Schrödinger Operator	288 R. Durrett, R.H. Schonmann,	Stochastic Growth Models		
253	H. Brezis, J.-M.Coron, E.H. Ljeb,	Harmonic Maps with Defects	289 David Kinderlehrer,	Remarks about Equilibrium Configurations of Crystals		
254	A. Carverhill,	Flows of Stochastic Dynamical Systems: Nontriviality of the Lyapunov Spectrum	290 D.G. Aronson, J.L. Vazquez,	Eventual C^∞ -Regularity and Concavity for Flows in One-Dimensional Porous Media		
255	A. Carverhill,	Conditioning A 'Lifted' Stochastic System in a Product Case	291 L.R. Scott, J.M. Boyle, B. Bagheri,	Distributed Data Structures for Scientific Computation		
256	R.J. Williams,	Local Time and Excursions of Reflected Brownian Motion	292 J. Douglas,Jr., P.J.Paes Leme, T. Arbogast, T. Schmitt,	Simulation of Flow in Naturally Fractured Petroleum Reservoirs		
257	H. Föllmer, S. Orey,	Large Deviations for the Empirical Field of a Gibbs Measure	293 D.G. Aronson, L.A. Caffarelli,	Optimal Regularity for One-Dimensional Porous Medium Flow		
258	A. Leizarowitz,	Characterization of Optimal Trajectories On an Infinite Horizon	294 Haim Brezis,	Liquid Crystals and Energy Estimates for S^2 -Valued Maps		
259	Y.Giga, T. Miyakawa, H. Osada,	Two Dimensional Navier Stokes Flow with Measures As Initial Vorticity				
260	M. Chipot, V. Ollker,	Sur Une Propriete Des Fonctions Propres De L'operateur De Laplace Beltrami				
261	V. Perez-Abreu,	Decompositions of Semimartingales On Duals of Countably Nuclear Spaces				
262	J.M. Ball,	Does Rank-One Convexity Imply Quasiconvexity				
263	B. Cockburn,	The Quasi-Monotone Schemes for Scalar Conservation Laws. Part I				
264	K.A. Perlick-Spector,	On Radially Symmetric Simple Waves In Elasticity				
265	P.N. Shivakumar, Chl-Sing Man, Simon W. Rabkin,	Modelling of the Heart and Pericardium at End-Diastole				
266	Jose-Luis Menaldi,	Probabilistic View of Estimates for Finite Difference Methods				
267	Robert Hardt, Harold Rosenberg,	Open Book Structures and Unicity of Minimal Submanifolds				
268	Bernardo Cockburn,	The Quasi-Monotone Schemes for Scalar Conservation Laws				
269	H.R. Jauslin, W. Zimmermann,Jr.,	Dynamics of a Model for an Ac Josephson Effect				
270	A.K. Kapila,	Introductory Lecture on Reacting Flows				
271	J.C. Taylor,	Do Minimal Solutions of Heat Equations Characterize Diffusions?				
272	J.C. Taylor,	The Minimal Eigenfunctions Characterize the Ornstein-Uhlenbeck Process				
273	Chl-Sing Man, Quan-Xin Sun,	On the Significance of Normal Stress Effects in the Flow of Glaciers				
274	Omar Hijab,	On Partially Observed Control of Markov Processes				
275	Lawrence Gray,	The Behavior of Processes with Statistical Mechanical Properties				
276	R. Hardt, D. Kinderlehrer, M. Luskin,	Remarks About the Mathematical Theory of Liquid Crystals				

THE EXTRAPOLATION FOR BOUNDARY FINITE ELEMENTS

Li Kaitai* Yan Ningning

(Institute for Computational and Applied Mathematics

Xian Tiaolong University, XIAN, CHINA)

I. Introduction

It has been shown in [1]-[3] that the Richardson extrapolation can be applied to the elliptic Ritz projection with linear finite elements and increase the second-order accuracy of linear finite elements

$$U_h(z) = u(z) + O(h^2 |\ln 1/h|)$$

in mesh point z at least to order three or four

$$U_h(z) \equiv \frac{1}{3} (4 U_{h/2}(z) - U_h(z)) = u(x) + O(h^3 |\ln 1/h|) \text{ (or } O(h^4 |\ln 1/h|))$$

Where T_h is uniform triangulation and $T_{h/2}$ is generated from T_h by dividing each triangle as usual into four congruent subtriangles.

In the present paper above basic results have been extended to boundary finite element and the coupling of finite elements and boundary finite elements.

* Research undertaken during visit to the Institute for Mathematics and its Applications

2. The Extrapolation for Boundary Finite Elements

Let us consider the following boundary value problem

$$-\Delta u = f \quad \text{in } \Omega \quad (2.1)$$

$$u = u_0 \quad \text{on } \Gamma$$

where $\Omega \subset \mathbb{R}^2$ is a bounded domain with Lipschitz boundary. The equivalent integral equation is that

$$\int_{\Gamma} q(x) E(x; y) ds_x = \int_{\Gamma} u_0(x) \frac{\partial E(x; y)}{\partial n} ds_x + \frac{1}{2} u_0(y) - \int_{\Omega} f(x) E(x; y) dx \quad (2.2)$$

$$\forall y \in \Gamma$$

$$u(y) = \int_{\Omega} f(x) E(x; y) dx + \int_{\Gamma} q(x) E(x; y) ds_x - \int_{\Gamma} u_0(x) \frac{\partial}{\partial n} E(x; y) ds_x \quad (2.3)$$

$$\forall y \in \Omega$$

Where $E(x; y)$ is a fundamental solution of Laplace equation, i.e.

$$\Delta_x E(x; y) + \delta(x-y) = 0$$

and $q = \frac{\partial u}{\partial n} \Big|_{\Gamma}$. For two dimensional problem it is known

$$E(x; y) = -\frac{1}{2\pi} \ln |x-y| \quad (2.4)$$

The Galerkin variational formulation of (2.2) is that

$$\text{find } q \in H^{-1/2}(\Gamma) \text{ such that} \quad (2.5)$$

$$b(q, q^*) = \mathfrak{L}((u_0, f), q^*) \quad \forall q^* \in H^{-1/2}(\Gamma)$$

where

$$b(q, q^*) = \int_{\Gamma} \int_{\Gamma} q(x) q^*(y) E(x; y) ds_x ds_y$$

is a symmetry and coercive bilinear form on $H^{-1/2}(\Gamma) \times H^{-1/2}(\Gamma)$

$$\begin{aligned} \mathfrak{L}((u_0, f), q^*) &= \int_{\Gamma} \int_{\Gamma} u_0(x) q^*(y) \frac{\partial}{\partial n_x} E(x; y) ds_x ds_y + \frac{1}{2} \int_{\Gamma} u_0(x) ds_x \\ &\quad - \int_{\Gamma} \int_{\Omega} f(x) q^*(y) E(x; y) dx ds_y \end{aligned}$$

is a bounded linear functional on $H^{-1/2}(\Gamma)$.

The approximation problem corresponding to (2.5) can be expressed as the following

$$\text{find } q_h \in V_h \subset H^{-1/2}(\Gamma) \text{ such that} \quad (2.6)$$

$$b(q_h, q_h^*) = \mathfrak{L}((u_0, f), q_h^*), \quad \forall q_h^* \in V_h$$

Where V_h is a boundary finite element subspace on Γ , and h is mesh size parameter. Later on, we suppose V_h is a linear elements subspace on Γ .

Lemma 1 Assume $E(x;y)$ is a fundamental solution defined by (2.4). Then there exists $\tilde{E}(x;y) \in V_h$ such that

$$\begin{aligned} \|\tilde{E}(\cdot;y)\|_{0,2,\Gamma} &\leq c |\ln h|^{1/2} \\ \|\tilde{E}(\cdot;y)\|_{1/2,\Gamma} &\leq c |\ln h|^{1/2} \\ \|\tilde{E}(\cdot;y) - E(\cdot,y)\|_{0,1,\Gamma} &\leq c h |\ln h| \end{aligned} \quad (2.7)$$

where c is a positive constant independent of h and y .

Proof. Suppose

$$M_y = \{x \mid \|x-y\| \leq 1\}, \quad \Gamma_1 = \Gamma \cap M_y, \quad \Gamma_2 = \Gamma \setminus \Gamma_1$$

Because of $E(x;y)$ satisfying

$$-\Delta E(x;y) = \delta(x-y) \quad \text{in } M_y$$

$$E(x;y) = 0 \quad \text{on } \partial M_y$$

we have $E(x;y) \in H_0^{1-\varepsilon}(M_y)$ and

$$(\nabla E(x;y), \nabla v) = v(y) \quad \forall v \in H_0^{1-\varepsilon}(M_y)$$

By [5], there exists $E_h(x;y) \in W_h \subset H_0^1(M_y)$ (where W_h is piecewise linear interpolation finite element subspace of $H_0^1(M_y)$), such that

$$(\nabla E_h(\cdot; y), \nabla v) = v(y) \quad \forall v \in W_h$$

and

$$\|E_h(\cdot; y)\|_{1,2,M_y} \leq c |1nh|^{1/2}$$

$$\|E_h(\cdot; y) - E(x; y)\|_{1,1,M_y} \leq ch |1nh|$$

so
$$\|E_h(\cdot; y)\|_{1,2,M_{y \cap \Omega}} \leq c |1nh|^{1/2}$$

$$\|E(\cdot; y) - E_h(\cdot; y)\|_{1,1,M_{y \cap \Omega}} \leq ch |1nh|$$

By virtue of trace theory, we can get

$$\|E_h(\cdot; y)\|_{0,2,\partial(M_y \cap \Omega)} \leq c |1nh|^{1/2}$$

$$\|E_h(\cdot; y) - E(\cdot; y)\|_{0,1,\partial(M_y \cap \Omega)} \leq ch |1nh|$$

In view of $\Gamma_1 \subset (M_y \cap \Omega)$, we have

$$\|E_h(\cdot; y)\|_{0,2,\Gamma_1} \leq c |1nh|^{1/2}$$

$$\|E_h(\cdot; y) - E(\cdot; y)\|_{0,1,\Gamma_1} \leq ch |1nh|$$

Because of $\|x-y\| > 1, \quad \forall x \in \Gamma_2$

so
$$\|E(\cdot; y)\|_{2,\infty,\Gamma_2} \leq 1$$

Suppose $E^1(\cdot; y)$ is a piecewise linear interpolation of $E(x; y)$, then

$$\|E^I(\cdot; y) - E(\cdot; y)\|_{0,1,\Gamma_2} \leq ch^2 \|E(\cdot; y)\|_{2,1,\Gamma_2} \leq \tilde{c}h^2$$

$$\|E^I(\cdot; y)\|_{0,\Gamma_2} \leq c$$

$$\|E^I(\cdot; y)\|_{1,\Omega} (My \quad \Omega) \leq c$$

Setting

$$\tilde{E}(x; y) = \begin{cases} E_h(x; y) & x \in \Gamma_1 \\ E^I(x; y) & x \in \Gamma_2 \end{cases}$$

then

$$\|\tilde{E}(\cdot; y)\|_{0,r} \leq \|E_h(\cdot; y)\|_{0,\Gamma_1} + \|E^I(\cdot; y)\|_{0,\Gamma_2} \leq c|\ln h|^{1/2}$$

$$\|\tilde{E}(\cdot; y)\|_{1,\Omega} \leq c|\ln h|^{1/2}$$

$$\|\tilde{E}(\cdot; y)\|_{1/2,\Gamma} = \inf\{\|v\|_{1,\Omega}, \gamma_0^* \tilde{E}(t, y) = \gamma_0 v\}$$

$$\leq \|E(\cdot; y)\|_{1,\Omega} \leq c|\ln h|^{1/2}$$

where γ_0 is the trace operator of 0-order on the Γ . In addition,

$$\|\tilde{E}(\cdot; y) - E(\cdot; y)\|_{0,1,\Gamma} \leq \|\tilde{E}(\cdot; y) - E(\cdot; y)\|_{0,1,\Gamma_1}$$

$$+ \|\tilde{E}(\cdot; y) - E(\cdot; y)\|_{0,1,\Gamma_2} \leq ch\|\ln h\|$$

The proof ends.

Remark 1 In view of (7) it is clear that

$$\|\tilde{E}(\cdot; y)\|_{0,1,\Gamma} \leq c |\ln h|^{1/2}$$

Lemma 2 Suppose G_z is the Green function of boundary integral equation (2.2) ,

$$b(q, G_z) = q(z) \quad \forall q \in H^{-1/2}(\Gamma)$$

$G_{zh} \in V_h$ is the approximation of G_z

$$b(q, G_{zh}) = q(z) \quad \forall q \in V_h \subset H^{-1/2}(\Gamma)$$

Then we have

$$\|G_{zh}\|_{-3/2,\Gamma} \leq c |\ln h|^{1/2} \quad (2.8)$$

$$\|G_{zh} - G_z\|_{-3/2,\Gamma} \leq c h$$

where c is a positive constant independent of h and z .

Proof. (1). $\forall g \in H^{3/2}(\Gamma)$, there exists a $q_g \in V_h \subset H^{1/2}(\Gamma)$

such that

$$b(q_g, q') = \langle g, q' \rangle \quad \forall q' \in V_h$$

and

$$\|q_g\|_{1/2,\Gamma} \leq c \|g\|_{3/2,\Gamma}$$

By [6] ,

$$\langle g, G_{zh} \rangle = b(q_g, G_{zh}) = q_g(z) \leq \|q_g\|_{0,\infty,\Gamma} \leq \quad ([6])$$

$$c | \ln h |^{1/2} \|g_g\|_{1/2, \Gamma} \leq c | \ln h |^{1/2} \|g\|_{3/2, \Gamma}$$

So

$$\|G_{Zh}\|_{-3/2, \Gamma} = \sup_{g \in H^{3/2}(\Gamma)} \frac{\langle g, G_{Zh} \rangle}{\|g\|_{3/2, \Gamma}} \leq c | \ln h |^{1/2}$$

(2) In similar manner, $\forall g \in H^{3/2}(\Gamma)$, there exist $q_g \in H^{1/2}(\Gamma)$, $q_h \in V_h \subset H^{1/2}(\Gamma)$ such that

$$b(q_g, q') = \langle g, q' \rangle \quad \forall q' \in H^{-1/2}(\Gamma)$$

$$b(q_h, q') = \langle g, q' \rangle \quad \forall q' \in V_h$$

By [11],

$$\|q_g - q_h\|_{0, \Gamma} \leq ch^{3/2} \|q_g\|_{1/2, \Gamma} \leq c h^{3/2} \|g\|_{3/2, \Gamma}$$

So

$$\begin{aligned} \langle g, G_Z - G_{ZR} \rangle &= b(q_g - q_h, G_Z - G_{Zh}) = b(q_g - q_h, G_Z) \\ &= q_g(Z) - q_h(Z) \leq \|q_g - q_h\|_{0, \infty, \Gamma} \leq ch^{-1/2} \|q_g - q_h\|_{0, \Gamma} \\ &\leq c h \|g\|_{3/2, \Gamma} \end{aligned}$$

$$\|G_Z - G_{Zh}\|_{-3/2, \Gamma} = \sup_{g \in H^{3/2}(\Gamma)} \frac{\langle g, G_Z - G_{Zh} \rangle}{\|g\|_{3/2, \Gamma}} \leq ch \quad \#$$

Assume $F(\Gamma_h)$ is a finite element regular family on Γ which satisfies that the distance from any point on the arc $S_i S_{i+1}$ to line $\overline{S_i S_{i+1}}$ is shorter than $\theta(h^2)$. Then we have

Lemma 3 Suppose $q \in C^3(\Gamma)$, $q^I \in V_h$ is the interpolation of q , then

$$b(q^I - q, G_{Zh}) = \frac{h^2}{12} b(q'', G_{Zh}) + O(h^3 |\ln h|^{3/2}) \|q\|_{3, \infty, \Gamma} \quad (2.9)$$

Proof. In fact, on the $\overline{S_i S_{i+1}}$

$$\begin{aligned} q^I(s) &= \frac{S_{i+1} - s}{h} q(S_i) + \frac{s - S_i}{h} q(S_{i+1}) \\ &= \frac{S_{i+1} - s}{h} (q(s) + (S_i - s) q'(s) + \frac{1}{2} (S_i - s)^2 q''(s) + \frac{1}{6} (S_i - s)^3 q'''(\xi)) \\ &\quad + \frac{s - S_i}{h} (q(s) + (S_{i+1} - s) q'(s) + \frac{1}{2} (S_{i+1} - s)^2 q''(s) + \frac{1}{6} (S_{i+1} - s)^3 q'''(\eta)) \\ &= q(s) + \frac{1}{2} (S_{i+1} - s)(s - S_i) q''(s) + O(h^3) \|q\|_{3, \infty, \Gamma} \end{aligned}$$

Therefore we have

$$\begin{aligned} I(y) &= \int_{\Gamma} (q^I(s) - q(s)) E(x; y) ds_x = \int_{\Gamma} (q^I - q) \tilde{E}(x; y) ds_x \\ &\quad + \int_{\Gamma} (q^I - q) (E(x; y) - \tilde{E}(x; y)) ds_x = I_1 + I_2 \end{aligned} \quad (2.10)$$

By lemma 1 and Remark 1 we have

$$I_2 \leq C h^2 \|q\|_{2, \infty, \Gamma} \int_{\Gamma} |\tilde{E}(x; y) - E(x; y)| dS_x = O(h^3 |\ln h|) \|q\|_{2, \infty, \Gamma}$$

$$I_1 = \sum_i \int_{\overline{S_i S_{i+1}}} (q^I - q) \tilde{E}(x; y) ds_x$$

$$= \sum_i \int_{\overline{S_i S_{i+1}}} (q^I - q) \tilde{E}(x; y) ds_x + \sum_i \left(\int_{\overline{S_i S_{i+1}}} - \int_{\overline{S_i S_{i+1}}} \right) (q^I - q) \tilde{E}(x; y) ds_x = L_1 + L_2$$

$$L_2 \leq c h^2 \int_{\Gamma} \left| \frac{\partial}{\partial n} ((q^I - q) \tilde{E}(x; y)) \right| dS_x$$

$$= ch^2 \{ \|q - q\|_{1, \infty, \Gamma} \|\tilde{E}\|_{1, 1, \Gamma} + \|q^I - q\|_{1/2, \Gamma} \|\frac{\partial \tilde{E}}{\partial n}\|_{1/2, \Gamma} \}$$

$$\leq ch^3 |\ln h|^{1/2} \|q\|_{2, \infty, \Gamma} + ch^{3+1/2} \|q\|_{2, \Gamma} \|\tilde{E}\|_{1/2, \Gamma} \leq ch^3 |\ln h|^{1/2} \|q\|_{2, \infty, \Gamma} \quad (2.11)$$

on the $\overline{S_i S_{i+1}}$, by virtue of $\tilde{E}(x; y) \in V_h$,

$$\tilde{E}(x; y) = \frac{S_{i+1} - S}{h} \tilde{E}(x_i; y) + \frac{S - S_i}{h} \tilde{E}(x_{i+1}; y) = N_i(s) \tilde{E}(x_i; y) + N_{i+1}(S) \tilde{E}(x_{i+1}; y)$$

$$\int_{\overline{S_i S_{i+1}}} N_i(s) ds = \frac{1}{h} \int_{S_i}^{S_{i+1}} (S_{i+1} - s) ds = h/2$$

$$\int_{\overline{S_i S_{i+1}}} N_{i+1}(s) ds = h/2$$

Hence

$$L_1 = \sum_i \int_{\overline{S_i S_{i+1}}} (q^I - q) \tilde{E}(x; y) ds_x = \sum_i \int_{\overline{S_i S_{i+1}}} \left(\frac{1}{2} (S_{i+1} - S)(S - S_i) q''(s) \right) ds_x$$

$$(N_i(s) \tilde{E}(x_i; y) + N_{i+1}(s) \tilde{E}(x_{i+1}; y)) ds_x + O(h^3) \|q\|_{3, \infty, \Gamma} \int_{\Gamma} |\tilde{E}(x; y)| ds_x$$

$$= \sum_i q''(\xi) \left(\frac{h^3}{24} \tilde{E}(x_i; y) + \frac{h^3}{24} \tilde{E}(x_{i+1}; y) \right) + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma}$$

$$= \sum_i \frac{h^2}{12} \{ q''(\xi) \left(\int_{S_i S_{i+1}} (N_i(s) \tilde{E}(x_i; y) + N_{i+1}(s) \tilde{E}(x_{i+1}; y)) ds_x \right) \} + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma}$$

$$\begin{aligned} L_1 &= \frac{h^2}{12} \int_{\Gamma} q''(s) \tilde{E}(x; y) ds_x + \frac{h^2}{12} \sum_i \int_{S_i S_{i+1}} (q''(\xi) - q''(s)) \tilde{E}(x; y) ds_x \\ &\quad + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \end{aligned}$$

but

$$\int_{S_i S_{i+1}} (q''(\xi) - q''(s)) \tilde{E}(x; y) ds_x = O(h |\ln h|^{1/2})$$

So

$$\begin{aligned} L_1 &= \frac{h^2}{12} \int_{\Gamma} q''(s) \tilde{E}(x; y) ds_x + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \\ &= \frac{h^2}{12} \int_{\Gamma} q''(s) E(x; y) ds_x + \frac{h^2}{12} \int_{\Gamma} q''(s) (\tilde{E}(x; y) - E(x; y)) ds_x \\ &\quad + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \quad (\text{by lemma 1}) \end{aligned}$$

$$= \frac{h^2}{12} \int_{\Gamma} q''(s) E(x; y) ds_x + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \quad (2.12)$$

Consequently,

$$I(y) = I_1 + I_2 = L_1 + L_2 + I_2 = \frac{h^2}{12} \int_{\Gamma} q''(s) E(x; y) ds_x + O(h^3 |\ln h|) \|q\|_{3, \infty, \Gamma} \quad (2.13)$$

Therefore,

$$\begin{aligned} b(q^I - q_{1, G_{Zh}}) &= \int_{\Gamma} I(y) G_{Zh}(y) ds_y = \frac{h^2}{12} \int_{\Gamma} \int_{\Gamma} q''(x) G_{Zh}(y) E(x; y) ds_x ds_y \\ &+ O(h^3 |\ln h|) \|q\|_{3, \infty, \Gamma} \|G_{Zh}\|_{-3/2, \Gamma} \\ &= \frac{h^2}{12} \int_{\Gamma} \int_{\Gamma} q''(x) G_Z(y) E(x; y) ds_x ds_y + \frac{h^2}{12} \int_{\Gamma} \int_{\Gamma} q''(x) (G_{Zh}(y) - G_Z(y)) E(x; y) ds_x ds_y \\ &+ O(h^3 |\ln h|^{3/2}) \|q\|_{3, \infty, \Gamma} \\ &= \frac{h^2}{12} b(q'', G_Z) + O(h^2 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \|G_{Zh} - G_Z\|_{0, 1, \Gamma} + O(h^3 |\ln h|^{3/2}) \end{aligned}$$

Owing to that

$$\|G_{Zh} - G_Z\|_{0, 1, \Gamma} \leq |\Omega| \|G_{Zh} - G_h\|_{-3/2} \leq c h$$

Finally, we obtain

$$b(q^I - q, G_{Zh}) = \frac{h^2}{12} b(q'', G_Z) + O(h^3 |\ln h|^{3/2}) \|q\|_{3, \infty, \Gamma}$$

Theorem 1 Suppose q is the solution of (2.5), $q \in C^3(\Gamma)$, $q_h \in V_h \subset H^{-1/2}(\Gamma)$ is a solution of (2.6), then for any mesh point Z we have

$$\frac{1}{3} (4 q_{h/2}(Z) - q_h(Z)) = q(Z) + O(h^3 |\ln h|^{3/2}) \|q\|_{3, \infty, \Gamma} \quad (2.14)$$

Proof since

$$b(q - q_h, G_{2h}) = 0$$

$$b(q, G) = q(z), \quad b(q_h, G_{zh}) = q_h(z)$$

∀ mesh point z we have

$$q(z) - q_h(z) = q^I(z) - q_h(z) = b(q^I - q_h, G_{zh})$$

$$= b(q^I - q, G_{zh}) = \frac{h^2}{12} b(q'', G_{zh}) + O(h^3 |\ln h|^{3/2}) \|q\|_{3, \infty, \Gamma}$$

Hence (2.14) is obtained.

3. Extrapolation for the Collocation Method

The integral equation (2.2) can be expressed as the following

find $q \in H^{-1/2}(\Gamma)$ such that

$$\int_{\Gamma} q(x) E(x;y) dS_x = F(y) \quad (3.1)$$

Where

$$F(y) = \int_{\Gamma} u_0(x) \frac{\partial}{\partial n} E(x;y) ds_x + \frac{1}{2} u_0(y) - \int_{\Omega} f(x) E(x;y) dx$$

The collocation method for (3.1) is that

find $q_h \in V_h \subset H^{-1/2}(\Gamma)$ such that

$$\int_{\Gamma} q_h(x) E(x;y_j) ds_x = F(y_j) \quad j = 1, 2, \dots, n \quad (3.2)$$

where y_j are mesh points of Γ_h .

From (3.1) and (3.2) we have

$$\int_{\Gamma} (q(x) - q_h(x)) E(x;y_j) ds_x = 0, \quad j = 1, 2, \dots$$

Hence

$$\int_{\Gamma} (q^I(x) - q_h(x)) E(x;y_j) ds_x = \int_{\Gamma} (q^I(x) - q(x)) E(x;y_j) ds_x, \quad j=1, 2, \dots, n \quad (3.3)$$

i.e.

$$\int_{\Gamma} (q^I(x) - q_h(x)) E(x;y_j) ds_x = I(y_j) \quad j = 1, 2, 3, \dots, n \quad (3.4)$$

where $I(y)$ is defined by (2.10). By virtue of (2.13) we have

$$\int_{\Gamma} [q^I(x) - q_h(x) - \frac{h^2}{12} q''(s)] E(x; y_j) ds_x = O(h^3 \|q\|_{3, \infty, \Gamma}) \quad (3.5)$$

$$j = 1, 2, \dots, n.$$

Let N_i, N_{i+1} are shape function of lineare element in one dimensional space.

$$N_i(s) = \frac{S_{i+1} - s}{h}, \quad N_{i+1}(s) = \frac{s - S_i}{h}$$

Then

$$(q^I(x) - q_h(x) - \frac{h^2}{12} q''(s)) \Big|_{S_i S_{i+1}} = \alpha_i N_i(s) - \alpha_{i+1} N_{i+1}(s)$$

The algebraic equations (3.5) can be expressed as

$$AX = F$$

$$\text{where } X^T = \{\alpha_1, \alpha_2, \dots, \alpha_n\}, \quad F^T = \{f_1, f_2, \dots, f_n\}$$

$$A = (a_{ij}),$$

$$a_{ij} = \int_{S_{i-1} S_i} N_{i+1}(s) E(x; y_j) ds_x + \int_{S_i S_{i+1}} N_i(s) E(x; y_j) ds_x$$

$$f_i = O(h^3 \|q\|_{3, \infty, \Gamma})$$

It is clear that

$$\left| \int_{S_i S_{i+1}} N_i(s) E(x; y_j) ds_x \right| = \left| \int_{S_i S_{i+1}} N_i(s) (E(x, y_j) - \tilde{E}(x_i y_j)) ds_y \right|$$

$$+ \int_{s_i s_{i+1}} N_i(s) \tilde{E}(x; y_j) ds_x| \leq c h |\ln h| + c |\ln h|^{1/2}$$

Hence

$$a_{ij} = O(h^{1/2})$$

$$\alpha_i = O\left(\frac{(\ln h)^{1/2}}{(\ln h)^{1/2}}\right)^{n-1} h^3 |\ln h| \|q\|_{3, \infty, \Gamma}$$

$$= O(h^3 |\ln h|^{1/2})$$

Finally, we obtain: \forall mesh point z

$$q^I(z) - q_h(z) = \frac{h^2}{12} q''(x) + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma}$$

By virtue of $q^I(z) = q(z)$ we have

Lemma 4 Suppose $q \in C^\infty(\Omega)$ is a solution of (2.5) and q_h is a collocation solution of (3.2), then

$$q(z) - q_h(z) = \frac{h^2}{12} q''(x) + O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \quad (3.6)$$

for any mesh points z

From lemma 4 we deduce the extrapolation formula for collocation method

$$\frac{1}{3} (4 q_{h/2}(z) - q_h(z)) - q(z) = O(h^3 |\ln h|^{1/2}) \|q\|_{3, \infty, \Gamma} \quad (3.7)$$

4. Numerical Test

Let us consider the following problems

$$-\Delta u(x) = -2\pi\delta(x) \quad \forall x \in \Omega = \{x \mid |x| \leq 0.5\}.$$

$$u|_{\Gamma} = \ln(0.5) \quad \text{on } \Gamma = \partial\Omega$$

Exact and Approximation solutions

$\tilde{q}_h(x) = \frac{1}{3} (4q_{h/2}(x) + q_h(x))$	$q_h(x)$	$q_{h/2}(x)$	$q(x)$
1.9999950	1.9986870	1.9996680	2.0000000
1.9999770	1.9986910	1.9996560	2.0000000
1.9999810	1.9986830	1.9996570	2.0000000
1.9999740	1.9986800	1.9996520	2.0000000
1.9999790	1.9986900	1.9996570	2.0000000
1.9999790	1.9986830	1.9996550	2.0000000
1.9999700	1.9986910	1.9996500	2.0000000
1.9999800	1.9986860	1.9996570	2.0000000
1.9999640	1.9986880	1.9996450	2.0000000
1.9999880	1.9986890	1.9996630	2.0000000
1.9999650	1.9986860	1.9996450	2.0000000
1.9999800	1.9986890	1.9996580	2.0000000
1.9999720	1.9986830	1.9996500	2.0000000
1.9999800	1.9986920	1.9996580	2.0000000
1.9999550	1.9986830	1.9996370	2.0000000
1.9999860	1.9986880	1.9996620	2.0000000
1.9999720	1.9986880	1.9996510	2.0000000

$q_h(x) = \frac{1}{3} (4q_{h/2}(x) + q_h(x))$	$q_h(x)$	$q_{h/2}(x)$	$q(x)$
1.9999510	1.9986830	1.9996340	2.0000000
1.9999760	1.9986910	1.9996550	2.0000000
1.9999840	1.9986860	1.9996590	2.0000000
1.9999930	1.9986870	1.9996660	2.0000000
2.0000000	1.9986920	1.9996730	2.0000000
1.9999860	1.9986880	1.9996610	2.0000000
1.9999890	1.9986800	1.9996610	2.0000000
1.9999580	1.9986900	1.9996410	2.0000000
1.9999960	1.9986870	1.9996680	2.0000000
1.9999810	1.9986880	1.9996580	2.0000000
1.9999690	1.9986880	1.9996490	2.0000000
1.9999740	1.9986860	1.9996520	2.0000000
1.9999830	1.9986850	1.9996580	2.0000000
1.9999720	1.9986910	1.9996520	2.0000000
1.9999820	1.9986860	1.9996580	2.0000000

Error

$q(x) - \tilde{q}_h(x)$	$q(x) - q_h(x)$	$q(x) - q_{h/2}(x)$
.0000049	.0013126	.0003319
.0000225	.0013094	.0003443
.0000190	.0013167	.0003433
.0000263	.0013115	.0003477
.0000210	.0013096	.0003432

.0000215	.0013167	.0003452
.0000302	.0013092	.0003500
.0000196	.0013143	.0003432
.0000362	.0013115	.0003550
.0000119	.0013114	.0003368
.0000352	.0013142	.0003549
.0000197	.0013107	.0003425
.0000283	.0013169	.0003505
.0000197	.0013076	.0003417
.0000452	.0013165	.0003630
.0000137	.0013118	.0003382
.0000285	.0013118	.0003493
.0000491	.0013167	.0003660
.0000240	.0013086	.0003451
.0000161	.0013142	.0003406
.0000070	.0013133	.0003335
.0000001	.0013083	.0003272
.0000144	.0013117	.0003388
.0000114	.0013201	.0003387
.0000423	.0013101	.0003593
.0000044	.0013126	.0003315
.0000190	.0013118	.0003421
.0000305	.0013120	.0003510
.0000257	.0013144	.0003480
.0000170	.0013150	.0003415
.0000283	.0013087	.0003483
.0000185	.0013142	.0003424

References

1. H. Blum, Q. Lin and R. Rannacher, Asymptotic Error Expansion and Richardson Extrapolation for linear Finite Elements, Numer. Math. 49, 11-37 (1986).
2. Lin Qun, Wang Junping, Some Expansions of the Finite Element Approximation IMS --15, 1984, 12.
3. Zhu Qiding, Lin Qun, Single Directional of Finite Difference and Finite Element, Eng. Math., 1984.2.
4. Brebbia, C.A., The Boundary Element Method for Engineers, 1978.
5. Zhu Jialin, The Boundary Integral Method of Elliptic Boundary Problem.
6. Zhu Qiding, The Conforming Super Convergence about Finite Element Derivative, Compu. Math., 1983.
7. Coupling Method of Finite Element Method for Radiation Problems. (To appear).
8. Scott. R. (1976), Optimates for the Finite Element Method on Irregular Meshes, Math, Compu., 30, 681 --697.
9. Claes Johnson and J. Claude Nedelec, On the Coupling of Boundary Integral and Finite Element Method, Math. Compu. Vol. 35, 1063 --1079, 1980.
10. J. C. Nedelec, Curved Finite Element Methods for the Solution of Singular Integral Equations on Surfaces in R^3 , Comput. Methods App. Mech. Engi., Vol. 8, 1976, 61-80.