

# Exact analytic solution for slope flows with spatially varying eddy viscosity and diffusivity

M. G. Giometto · R. Grandi · J. Fang ·  
M.B. Parlange

Received: DD Month YEAR / Accepted: DD Month YEAR

**Abstract** An exact analytic solution of the steady-state Prandtl model equations is derived, valid for spatially varying eddy diffusivities (O’Briens type) and Prandtl number of unity. In this formulation profiles show significant variations in both phase and amplitude of minima-maxima with respect to the classic constant eddy viscosity model and the more recent, approximate, WKB solution. The near wall region is characterized by a relatively stronger surface inversion and velocity gradients, the low-level-jet is further displaced toward the wall, and its peak velocity strongly depends on the model parameter, suggesting a tighter coupling between dynamics and thermodynamics.

## 1 Introduction

Natural convection of turbulent stratified flows over sloping surfaces is ubiquitous in nature and engineering, and is of interest not only as a fundamental problem in itself, but also because of the important role it plays over a broad

---

M.G. Giometto  
School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland  
E-mail: marco.giometto@epfl.ch

Riccardo Grandi  
Mathematics Institute for Geometry and Applications, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

J. Fang  
School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

M.B. Parlange  
Civil Engineering, Faculty of Applied Sciences, University of British Columbia, Vancouver, BC, Canada

range of scales and applications. On a local scale, for instance, it governs local weather conditions, affecting atmospheric transport processes and dispersion of scalars such as heat and humidity (Monti et al., 2002; Nylen et al., 2004). On a regional scale, it drives the development of deep-convective circulations (Egger, 1985; Parish and Bromwich, 1998) and is responsible for intense cyclonic vorticity in the middle and upper troposphere (Parish, 1992). Further, recent experimental field campaigns (Chu, 1987; Oerlemans and Vugts, 1993) have shown that persistent katabatic winds characterize the atmospheric boundary layer over snow-ice surfaces and glaciers, and therefore an accurate characterization of such flows is an essential component of understanding and modeling of the weather and climate. However, the complex dynamics (e.g. occurrence of intermittency, waves, Kelvin-Helmholtz instabilities and low-level jets) and the lack of a satisfactory similarity theory for such flows (Nadeau et al., 2013) pose a heavy burden in terms of computational requirements for numerical modellers; in most cases the required resolution is prohibitively costly (Shapiro and Fedorovich, 2004b,a; Fedorovich and Shapiro, 2009b,a; Spalart et al., 2011). Because of this, conceptual models are still of great interest, and represent a valid tool for the characterization of such systems. A cornerstone in the understanding of natural convection of stratified fluid along a sloping surface is represented by the classic Prandtl analytic model (Prandtl, 1942), and its more recent extensions/generalizations to include the effects of Coriolis force (Gutman and Malbakhov, 1964), external winds (Lykosov and Gutman, 1972), surface heterogeneity (Egger, 1985; Shapiro and Fedorovich, 2007; Oldroyd et al., 2014). The Prandtl model approximates the atmosphere in a Boussinesq sense and describes a steady flow over a thermally perturbed unbounded planar sloping surface that lies within a stratified environment, given by the following system of equations:

$$N^2 u(z) \sin \alpha = d(K_H db/dz)/dz, \quad (1)$$

$$b(z) \sin \alpha = -d(K_M du/dz)/dz, \quad (2)$$

where  $z$  denotes the normal-to-slope coordinate directions,  $u$  is the downslope velocity,  $b$  is buoyancy,  $N$  is the buoyancy frequency characterizing the system (related to the background stratification),  $\alpha$  is the slope angle and  $K_M$  and  $K_H$  denote the eddy viscosity and diffusivity (an eddy viscosity/diffusivity model has been used to parametrize turbulent fluxes of momentum and buoyancy). Equations are integrated in  $z \in [0, H]$  with boundary conditions  $u(0) = 0$ ,  $u(H) = 0$ ,  $b(0) = b_s$  and  $b(H) = 0$  ( $b_s > 0$  for upslope flows, whereas  $b_s < 0$  for downslope flows).

The flow is assumed to be invariant in the downslope direction and the model can be used to determine its vertical structure (Nappo and Shankar, 1987). Eqs. 1 and 2 state that downslope (upslope) convection of cold (warm) air is balanced by momentum flux divergence, and that along-slope generated buoyancy is balanced by buoyancy flux divergence; thus the model is applicable away from ridges and valleys, so that advection terms are negligible (Nappo and Shankar, 1987). The Prandtl constant-K model is known to be overdisipative in the near surface regions, and therefore not able to represent the

observed strong surface gradients of temperature and momentum that commonly characterize katabatic and anabatic flows (Oerlemans, 1998; Grisogono and Oerlemans, 2001). Simple variations in the eddy diffusivity profiles were introduced in an exact analytic solution by Gutman (1983), and more recently Grisogono and Oerlemans (2001) considered general variations in the vertical structure of the eddy diffusivities, and derived solutions based on the WKB approximation. The WKB solution is able to account for additional dynamics while still retaining an elegant form. However, that theory is only applicable when the model parameters ( $K_M, K_H$ ) vary more slowly than the solution ( $u, b$ ), and the validity of such a constraint for slope flows has been the subject of great debate (Grisogono and Oerlemans, 2002). In the following, we derive a closed-form solution to the Prandtl-model equations, valid for O'Brien's type (O'Brien, 1970) eddy diffusivities and representing an exact alternative to the WKB solution – for the chosen form of the model parameter.

## 2 The Analytic Solution

We consider the one-dimensional Prandtl model equations 1 and 2. Assuming  $Pr = 1$  (i.e.  $K_M = K_H = K$ ) and assigning length, velocity and buoyancy scales as  $L = \max(K)^{1/2} N^{-1/2} \sin^{-1/2} \alpha$ ,  $U = b_s N^{-1}$  and  $B = b_s$  Eqs. 1 and 2 can be reduced to:

$$u^+ = d(K^+ db^+ / dz^+) / dz^+, \quad \text{in } [0, H^+] \quad (3a)$$

$$b^+ = -d(K^+ du^+ / dz^+) / dz^+, \quad \text{in } [0, H^+] \quad (3b)$$

respectively, where  $K^+$  is a normalized eddy viscosity,  $z^+ = zL^{-1}$ ,  $u^+ = uU^{-1}$ ,  $b^+ = bB^{-1}$ , with boundary conditions  $u^+(0) = 0$ ,  $u^+(H^+) = 0$ ,  $b^+(0) = \pm 1$  and  $b^+(H^+) = 0$ . Further, we introduce the following variables

$$f_1 = b^+ - iu^+ \quad \text{and} \quad f_2 = b^+ + iu^+, \quad (4)$$

and upon substitution of Eqs. 3a and 3b into Eqs. 4, the system can be decoupled to obtain two (independent) complex ordinary differential equations (ODEs) for the canonical variables  $f_1$  and  $f_2$ :

$$i \cdot f_1 = d(K^+ df_1 / dz^+) / dz^+, \quad \text{in } [0, H^+] \quad (5a)$$

$$i \cdot f_2 = -d(K^+ df_2 / dz^+) / dz^+, \quad \text{in } [0, H^+] \quad (5b)$$

with boundary conditions  $f_i(0) = \pm 1$  and  $f_i(H^+) = 0$ ,  $i = 1, 2$ .

We assume eddy diffusivities in line with the classic O'Brien's type (O'Brien, 1970), viz.  $K^+ = A^+(z^+ + \epsilon^+)(z^+ - H^+ - \epsilon^+)^2$ , where  $\epsilon^+$  is an additional parameter that we introduce in order to shift the zeroes of  $K^+$  outside the interval of integration  $[0, H^+]$ .  $\epsilon^+$  can be tuned to provide the desired value of viscosity / diffusivity at the wall.

In order to solve Eqs. 5a and 5b, we observe that both have the form

$$f = \mp i \cdot d/dz^+ (K^+ df / dz^+),$$

and therefore can be solved using the same method. First, we rewrite the equation in canonical form:

$$f'' + Pf' + Qf = 0, \quad \text{in } [0, H^+] \quad (6)$$

where  $P(z^+) = (K^+)/K^+$  and  $Q(z^+) = 1/(\pm iK^+)$ . Second, it is an easy computation to show that, if  $f$  solves Eq. 6 then  $g(z^+) := f((H^+ + 2\epsilon^+)z^+ - \epsilon^+)$  solves:

$$g'' + \tilde{P}g' + \tilde{Q}g = 0, \quad \text{in } [0, 1] \quad (7)$$

where  $\tilde{P}(z^+) = \gamma'(z^+)/\gamma(z^+)$  and  $\tilde{Q}(z^+) = 1/(\pm i \cdot A^+ \cdot (H^+ + 2\epsilon^+)\gamma(z^+))$ , with  $\gamma(z^+) = z^+(z^+ - 1)^2$ . Eqn. 7 is a second order ODE with three regular singular points at  $z^+ = 0, 1$  and  $\infty$ , as in Morse and Feshbach (1953). This special case is known as the *equation of Papperitz* and its general solution is:

$$g(z^+) = \alpha(1-z^+)^{\mu} {}_2F_1(\mu, 1-\mu', 1+\mu-\mu', 1-z^+) + \beta(1-z^+)^{\mu'} {}_2F_1(\mu', 1-\mu, 1+\mu'-\mu, 1-z^+), \quad (8)$$

where  ${}_2F_1$  are *Gaussian hypergeometric functions* and  $\mu, \mu'$  are the solutions to the degree two equation

$$x^2 + x + 1/(\pm iA^+(H^+ + 2\epsilon^+)) = 0,$$


depending on which of Eqs. 7 we are solving. Note that the general solution to Eq. 7 is also often written using the *Riemann symbol*:

$$\left\{ \begin{array}{ccc} 0 & 1 & \infty \\ \lambda & \mu & \nu \\ \lambda' & \mu' & \nu' \end{array} \right\}$$


where, in this case,  $\mu$  and  $\mu'$  are as above,  $\lambda = \lambda' = 0$  and  $\nu = 0, \nu' = 2$ . To retrieve the general solution to Eq. 6 it is sufficient to substitute  $(z^+ + \epsilon^+)/(H^+ + 2\epsilon^+)$  in place of  $z^+$  in Eq. 8. As the integration constants  $\alpha$  and  $\beta$  are specified through the imposition of boundary conditions, back-substituting  $b^+ = (f_1 + f_2)/2$  and  $u^+ = -(f_1 - f_2)/2i$  yields the solution in terms of  $u^+$  and  $b^+$ .

### 3 Examples

In Fig. 1 we compare the new analytic solution (A1 in the following) against the constant- $K$  (A2), the WKB (A3) and a numerical solution (N1). The constant- $K$  value is fixed to  $K_{A2}^+ = \max(K_{A1}^+)/3$ . Profiles for N1 are derived by solving monolithically the system of ODEs defined by Eqs. 3a and 3b, adopting a second-order accurate centered finite difference discretization with over 40,000 collocation points and stretching the grid via a quadratic coordinate transform (at such resolution profiles are invariant to further refinements down to machine precision). The excellent match between A1 and



**Fig. 1** Comparison of the new analytic solution (A1) against the constant- $K$  (A2), the WKB (A3) and a numerical 1D solution (N1). Analytic profiles of normalized velocity ( $u^+$ ) are denoted with dot-dashed lines whereas analytic profiles of normalized buoyancy ( $b^+$ ) are denoted by dashed lines. Here  $K^+ = A^+ \cdot (z^+ + \epsilon^+)(z^+ - H^+ - \epsilon^+)^2$  where  $A^+ = 6.75e - 4$ ,  $\epsilon^+ = 1.5 \times 10^{-3}$  and  $H^+ = 10$ . A corresponding dimensional profile is characterized by  $\max(K) = 0.1 \text{ m}^2\text{s}^{-1}$  and  $K(z = 0) = 10^{-5} \text{ m}^2\text{s}^{-1}$ , in agreement with common atmospheric values.




**Fig. 2** Sensitivity of A1 (left) and A3 (right) profiles to variations in the diffusivity coefficient  $K^+$ . Normalized velocity values ( $u^+$ ) are denoted with dot-dashed lines whereas normalized buoyancy values ( $b^+$ ) with dashed lines. Equations are integrated in the interval  $z^+ \in [0, 10]$ , which is fully shown in the plot. Here we considered  $A^+ = 6.75 \times 10^{-4}$  and  $\epsilon^+ = (7.4, 1.5, 0.30, 0.06) \times 10^{-3}$ . Under the constraint  $K(z = 0) = 10^{-5} \text{ m}^2\text{s}^{-1}$ , the corresponding set of dimensional kinematic diffusivities satisfy  $\max(K) = 0.002, 0.01, 0.05, 0.25 \text{ m}^2\text{s}^{-1}$ .

**Table 1** Set of parameters for re-normalization of the reference solution. The reference normalized solution is computed adopting  $H^+ = 10$ ,  $A^+ = 6.75 \times 10^{-4}$ ,  $\epsilon^+ = 1.5 \times 10^{-3}$ .

Value	Ref.	$\alpha_1$	$\alpha_2$	$N_1$	$N_2$	$b_{s,1}$	$b_{s,2}$
$\alpha$ (deg)	30	15	60	30	30	30	30
$N$ (Hz)	0.01	0.01	0.01	0.005	0.02	0.01	0.01
$b_s$ (ms $^{-2}$ )	0.30	0.30	0.30	0.30	0.30	0.15	0.60

N1 certifies the quality of results for both methods. In comparison to its analytical counterparts, A1 shows a remarkably strong inversion in the near surface regions ( $z^+ < 1$ ), suggesting overdiffusive behaviour of both A2 and A3, which are incapable of providing such strong buoyancy gradients. For instance, normalized surface buoyancy gradients of simulation A1 are over an order of magnitude larger than those of A2 ( $(db/dz)_{A1}^+ / (db/dz)_{A2}^+ = \mathcal{O}(10)$  as  $z^+ \rightarrow 0$ ). Despite this, we observe a relatively good match between  $u_{A1}^+$  and  $u_{A3}^+$  in the interval  $z^+ \in [0, 1]$ , which confirms the enhanced dissipative properties of the A3 solution, when compared against the constant- $K$  approach. To understand the importance of a decreasing eddy viscosity in the near surface regions, it is worth noting that  $(du/dz)_{A1}^+ / (du/dz)_{A2}^+ = \mathcal{O}(10)$ , whereas  $(du/dz)_{A1}^+ / (du/dz)_{A3}^+ = \mathcal{O}(1)$  as  $z^+ \rightarrow 0$ . Overall the new solution exhibits significant variations when compared against A2 and A3, in both amplitude and location of maxima-minima. For instance, both the low-level jet height and the peak normalized velocity are significantly reduced, features that are of great importance for an accurate representation of the stable boundary layer and from a parameterization perspective (Mahrt, 1998). Further, from Fig. 2, it is clear that  $\max(u_{A1}^+)$  now depends on the diffusive properties of the flow, i.e.  $\max(u_{A1}^+) = f(K^+)$ , in contrast to its analytic counterparts, which predict  $\max(u_{A2}^+) = \max(u_{A3}^+) = \sqrt{2}/2[(e^{(\pi/4)})/\sqrt{Pr}] = 0.32$ , regardless of  $K^+$ .  $u_{A1}^+$  and  $b_{A1}^+$  also show a stronger dependence on the  $K^+$  parameter with respect to A3 profiles, in particular in the  $z^+ \in [0, 1]$  interval. Overall, A1 predicts significantly reduced mass and buoyancy (horizontal) fluxes, viz.  $\int_0^{H^+} u^+ dz^+$ ,  $\int_0^{H^+} b^+ dz^+$ , with respect to A3. To get acquainted with the proposed normalization Fig. 3 proposes a set of re-dimensionalized velocity profiles  $u$  (ms $^{-1}$ ) from a reference normalized solution. We propose a reference dimensional solution, characterized by  $\alpha = 30$  deg,  $N = 0.01$  Hz and  $b_s = 0.3$  ms $^{-2}$ , and consider variations in both sloping angle  $\alpha$ , imposed buoyancy frequency  $N$  and imposed surface buoyancy  $b_s$  (see Table 1). Variations in the sloping angle  $\alpha$  result in a modification of the normalization constant  $L \propto \sin^{-1/2}$  which acts as a stretching of the independent variable  $z$  (no effects on  $\max(u)$ ). In particular, the steeper the angle, the smaller the characteristic scales of the flow will become. Variations in the background stratification, viz.  $N$ , will modify both  $L \propto N^{-1/2}$  and  $U \propto N^{-1}$ , yielding changes in the system, whose lengths and velocities will tend to decrease (increase) with increasing (decreasing)  $N$ . Further, variations in the prescribed surface buoyancy  $b_s$  will scale both buoyancy and velocity profiles accordingly, since  $b(z) \propto b_s$  and  $u(z) \propto b_s$ . As in both the classic Prandtl and WKB linear models,  $K^+$  needs



**Fig. 3** Family of dimensional velocity profiles derived from re-dimensionalization of a reference normalized solution. The normalized solution is obtained in  $z^+ \in [0, 10]$  for  $A^+ = 6.75 \times 10^{-4}$  and  $\epsilon^+ = 1.5 \times 10^{-3}$  (assuming  $K(z=0) = 10^{-5} \text{ m}^2\text{s}^{-1}$ , the corresponding dimensional kinematic diffusivities satisfy  $\max(K) = 0.01 \text{ m}^2\text{s}^{-1}$ ). The considered set of parameters are reported in Table 1. Decreasing values of the parameters (with respect to the reference ones) are denoted with solid lines, increasing values by dashed lines.

to be assigned a-priori; it is not coupled and does not feed back into the solution, which represents the main weakness of such linear approaches (Grisogono and Oerlemans, 2001).

## 4 Conclusions

We have derived a closed-form analytic solution of the Prandtl model equations, based on an ad hoc decoupling of the system and specific choice of the model parameter, resulting in a set of *equations of Papperitz*, characterized by three regular singular points. The new solution is valid for O'Brien-type eddy diffusivities (cubic polynomials) and Prandtl number of unity. For the considered settings the new solution represents an improvement with respect to both the classic constant- $K$  and the more recent WKB solutions. New profiles show significant variations in both phase and amplitude of minima-maxima, stronger surface gradients are combined with overall reduced horizontal fluxes and the LLJ is further displaced toward the wall. The new solution suggests a somewhat different coupling between velocity and buoyancy – when compared against its analytical counterparts, showing a diffusivity dependent peak velocity and a remarkably strong inversion layer in the near surface location. In addition to a theoretical insight, the proposed solution can be used to improve future stable boundary layer parameterizations when coupled with other parts of the boundary layer physics.

## 5 Acknowledgements

This research was primarily funded by the Swiss National Science Foundation (SNSF-200021-134892) and by the Competence Center for Environmental Sustainability (CCES-SwissEx) of the ETH domain. We are also grateful for funding from the NSERC Discovery Grant program.

## References

- Chu PC (1987) An instability theory of ice-air interaction for the formation of ice edge bands. *Journal of Geophysical Research* 92(7):6966, DOI 10.1029/JC092iC07p06966
- Egger J (1985) Slope Winds and the Axisymmetric Circulation over Antarctica. *Journal of the Atmospheric Sciences* 42:1859–1867, DOI 10.1175/1520-0469(1985)042<1859:SWATAC>2.0.CO;2
- Fedorovich E, Shapiro A (2009a) Structure of numerically simulated katabatic and anabatic flows along steep slopes. *Acta Geophysica* 57(4):981–1010, DOI 10.2478/s11600-009-0027-4, URL <http://www.springerlink.com/index/10.2478/s11600-009-0027-4>
- Fedorovich E, Shapiro A (2009b) Turbulent natural convection along a vertical plate immersed in a stably stratified fluid. *Journal of Fluid Mechanics* 636:41, DOI 10.1017/S0022112009007757
- Grisogono B, Oerlemans J (2001) Katabatic Flow: Analytic Solution for Gradually Varying Eddy Diffusivities. *Journal of the Atmospheric Sciences* 58(21):3349–3354, DOI [http://dx.doi.org/10.1175/1520-0469\(2001\)058<3349:KFASFG>2.0.CO;2](http://dx.doi.org/10.1175/1520-0469(2001)058<3349:KFASFG>2.0.CO;2)
- Grisogono B, Oerlemans J (2002) Justifying the WKB approximation in pure katabatic flows. *Tellus, A* 54(5):453–462, DOI 10.1034/j.1600-0870.2002.201399.x
- Gutman LN (1983) On the theory of the katabatic slope wind. *Tellus, A* 35(3):332–335, DOI 10.3402/tellusa.v35i3.11434
- Gutman LN, Malbakhov VM (1964) On the Theory of the Katabatic Winds of Antarctica. University of Melbourne, Meteorology Department
- Lykosov VN, Gutman LN (1972) Turbulent boundary layer above a sloping underlying surface. *Proceedings of the USSR Academy of Sciences* 8:799–809, URL [papers2://publication/uuid/B2EC0692-4251-4224-8185-A1AE8D9B00E9](http://papers2://publication/uuid/B2EC0692-4251-4224-8185-A1AE8D9B00E9)
- Mahrt L (1998) Stratified Atmospheric Boundary Layers and Breakdown of Models. In: *Theoretical and Computational Fluid Dynamics*, vol 11, pp 263–279, DOI 10.1007/s001620050093, URL <http://link.springer.com/10.1007/s001620050093>
- Monti P, Fernando HJS, Princevac M, Chan WC, Kowalewski TA, Pardyjak ER (2002) Observations of Flow and Turbulence in the Nocturnal Boundary Layer over a Slope. *Journal of the Atmospheric Sciences* 59:2513–2534, DOI 10.1175/1520-0469(2002)059<2513:OOFATI>2.0.CO;2



- Morse PM, Feshbach H (1953) *Methods of Theoretical Physics*. McGraw-Hill
- Nadeau DF, Pardyjak ER, Higgins CW, Parlange MB (2013) Similarity Scaling Over a Steep Alpine Slope. *Boundary-Layer Meteorology* 147(3):401–419, DOI 10.1007/s10546-012-9787-5, URL <http://link.springer.com/10.1007/s10546-012-9787-5>
- Nappo CJ, Shankar RK (1987) A model study of pure katabatic flows. *Tellus A* 39(1):61–71, DOI 10.3402/tellusa.v39i1.11740, URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1600-0870.1987.tb00289.x/abstract>
- Nylen TH, Andrew G F, Doran PT (2004) Climatology of katabatic winds in the McMurdo dry valleys, southern Victoria Land, Antarctica. *Journal of Geophysical Research* 109:1–9, DOI 10.1029/2003JD003937
- O’Brien JJ (1970) A Note on the Vertical Structure of the Eddy Exchange Coefficient in the Planetary Boundary Layer. *Journal of the Atmospheric Sciences* 27:1213–1215, DOI 10.1175/1520-0469(1970)027<1213:ANOTVS>2.0.CO;2, URL [http://dx.doi.org/10.1175/1520-0469\(1970\)027<1213:ANOTVS>2.0.CO;2](http://dx.doi.org/10.1175/1520-0469(1970)027<1213:ANOTVS>2.0.CO;2)
- Oerlemans J (1998) The atmospheric boundary layer over melting glaciers. In: *Clear and Cloudy Boundary Layers*, vol 48, Royal Netherlands Academy of Arts and Sciences, pp 129–153, URL [http://igitur-archive.library.uu.nl/phys/2007-0716-203217/oerlemans\\_07\\_atmosphericboundarylayer.pdf](http://igitur-archive.library.uu.nl/phys/2007-0716-203217/oerlemans_07_atmosphericboundarylayer.pdf)
- Oerlemans J, Vugts HF (1993) A Meteorological Experiment in the Melting Zone of the Greenland Ice Sheet. *Bulletin of the American Meteorological Society* 74(3):355–365, DOI 10.1175/1520-0477(1993)074<0355:AMEITM>2.0.CO;2
- Oldroyd HJ, Katul G, Pardyjak ER, Parlange MB (2014) Momentum balance of katabatic flow on steep slopes covered with short vegetation. *Geophysical Research Letters* 41:4761–4768, DOI 10.1002/2014GL060313
- Parish TR (1992) On the Role of Antarctic Katabatic Winds in Forcing Large-Scale Tropospheric Motions. *Journal of the Atmospheric Sciences* 49:1374–1385, DOI 10.1175/1520-0469(1992)049<1374:OTROAK>2.0.CO;2
- Parish TR, Bromwich DH (1998) A Case Study of Antarctic Katabatic Wind Interaction with Large-Scale Forcing. *Monthly Weather Review* 126:199–209, DOI 10.1175/1520-0493(1998)126<0199:ACSOAK>2.0.CO;2
- Prandtl L (1942) *Führer durch die Strömungslehre*. Vieweg & Sohn, Braunschweig
- Shapiro A, Fedorovich E (2004a) Prandtl number dependence of unsteady natural convection along a vertical plate in a stably stratified fluid. *International Journal of Heat and Mass Transfer* 47(22):4911–4927, DOI 10.1016/j.ijheatmasstransfer.2004.04.035, URL <http://linkinghub.elsevier.com/retrieve/pii/S0017931004002157>
- Shapiro A, Fedorovich E (2004b) Unsteady convectively driven flow along a vertical plate immersed in a stably stratified fluid. *Journal of Fluid Mechanics* 498:333–352, DOI 10.1017/S0022112003006803, URL [http://www.journals.cambridge.org/abstract\\_S0022112003006803](http://www.journals.cambridge.org/abstract_S0022112003006803)

- 286 Shapiro A, Fedorovich E (2007) Katabatic flow along a differentially  
287 cooled sloping surface. *Journal of Fluid Mechanics* 571:149, DOI 10.  
288 1017/S0022112006003302, URL [http://www.journals.cambridge.org/  
289 abstract\\_S0022112006003302](http://www.journals.cambridge.org/abstract_S0022112006003302)
- 290 Spalart P, Strelets M, Garbaruk A (2011) Quality and Reliability of Large-  
291 Eddy Simulations II. In: *Evaluating Subgrid-Scale Models for Large-Eddy  
292 Simulation of Turbulent Katabatic Flow*, vol 16, pp 253–267, DOI 10.  
293 1007/978-94-007-0231-8, URL [http://www.springerlink.com/index/10.  
294 1007/978-94-007-0231-8](http://www.springerlink.com/index/10.1007/978-94-007-0231-8)