

Mufaddal M. Gombera, MD
Orthopedic Surgery and Sports Medicine
Fondren Orthopedic Group

PREOPERATIVE REHABILIATION (PREHAB) PROTOCOL FOR HIP ARTHROSCOPY PATIENTS

GOALS

- Physical therapy for 2 weeks prior to surgery
- Evaluation and establishment of goals after surgery
- Advise on maintaining postoperative ROM restrictions (see below)
- Instruct on 20# WB with flat-foot pattern with crutches, stair use
- Identify if additional equipment needed (e.g. seated toilet, grasper, only on as needed basis)
- Exercise program for core and gluteal strengthening
- Schedule post-op physical therapy visit for 4-7 days after surgery

EXERCISES

- Core muscle strengthening, gluteus strengthening
- Single leg calf-raises
- Single leg squats
- Side-lying hip abduction/adduction
- Four point kneeling hip rocks
- Kneeling hip flexor stretches

POST OP RESTRCTIONS

- No sitting with hip flexed >90 degrees to avoid hip flexor irritation
- Hip brace when ambulating (typically for 4 weeks, occasionally for 6 weeks)
- 20lb Weight bearing with foot flat gait pattern (can range from 2 to 6 weeks)
- ROM restrictions: Abduction to 45 degrees, no extension past 0, IR to neutral,

FOLLOW-UP

- Schedule first post-op therapy visit 4-7 days after surgery

7401 Main Street
Houston, TX 77030
www.fondren.com
www.gomberamd.com
Office: 713-794-3457
info@gomberamd.com