Probabilistic Machine Learning

Lecture 02: Probability and Statistics

Prof. Dr. Michael Granitzer

Where We Left Off

Lecture 01 ended on an ideal — model the joint, condition on the data, read off answers — and on the shortcuts most methods take. Today builds the language all of them are written in:

Model the joint
Random variables and their joint distribution

→


Condition on the data
For events (Conditional Probability) — for a parameter (Statistics and Estimation)

→

Read off answers
A peak, a mean, an interval (Statistics and Estimation)

It also left a claim unpaid: that **epistemic** uncertainty is the kind more data removes. Statistics and Estimation turns that slogan into a rate.

Argument Map



Probability by Counting

> ***Guiding Question:*** *A hundred students, two courses, one exam each. How far do we get with nothing but counting — before any definition?*

Example: Students, Courses and Exams

Probabilistic ML — 40 students
a quarter of them fail the exam


Databases — 60 students
one in ten fails the exam

Each of 100 students sat exactly one of the two exams.

- **Forward:** how many students fail overall?
- **Backward:** a student failed — how probable is it that they took PML?

Joint, Marginal, Conditional — One Table

? Of the PML students, what share failed?




P(fail | PML)

= **10** / **40** = 0.25

conditional: a cell, out of its own column

Sum Rule and Product Rule — On the Table

? 40% took PML, and a quarter of them failed. What share failed PML?



… of those, failed: P(fail | PML) = **10** / **40**

Product rule

P(PML, fail) = P(PML) · P(fail | PML)

= 40/100 · 10/40 = 10/100

a share of a share — the 40s cancel

The Two Rules

Sum rule

$$P(X) = \sum_{Y} P(X, Y)$$

Add along a row or column to remove a variable.

Product rule


$$P(X, Y) = P(Y \mid X) P(X)$$

A cell is its column total times its share of that column.

Almost all of probability theory is these two rules applied repeatedly.

Bayes' Theorem — On the Table

? And which course did they most probably take?



P(PML | fail) = **10** / **16** ≈ 0.63

P(DB | fail) = **6** / **16** ≈ 0.37

PML — odds 10 : 6

same normaliser: comparing needs only the cells

Bayes' Theorem, in Advance

Combining the two rules and the symmetry $P(X, Y) = P(Y, X)$:

$$P(Y \mid X) = \frac{P(X \mid Y)P(Y)}{P(X)}, \qquad P(X) = \sum_{Y} P(X \mid Y)P(Y)$$

$$P(\text{PML} \mid \text{fail}) = \frac{0.25 \cdot 0.4}{0.25 \cdot 0.4 + 0.1 \cdot 0.6} = \frac{0.10}{0.16} \approx 0.63$$

Bayes' Theorem, in Advance

Combining the two rules and the symmetry $P(X, Y) = P(Y, X)$:

$$P(Y \mid X) = \frac{P(X \mid Y)P(Y)}{P(X)}, \qquad P(X) = \sum_{Y} P(X \mid Y)P(Y)$$

$$P(\text{PML} \mid \text{fail}) = \frac{0.25 \cdot 0.4}{0.25 \cdot 0.4 + 0.1 \cdot 0.6} = \frac{0.10}{0.16} \approx 0.63$$

And if one in four DB students failed too?

Bayes' Theorem, in Advance

Combining the two rules and the symmetry $P(X, Y) = P(Y, X)$:

$$P(Y \mid X) = \frac{P(X \mid Y)P(Y)}{P(X)}, \qquad P(X) = \sum_Y P(X \mid Y)P(Y)$$

$$P(\text{PML} \mid \text{fail}) = \frac{0.25 \cdot 0.4}{0.25 \cdot 0.4 + 0.1 \cdot 0.6} = \frac{0.10}{0.16} \approx 0.63$$

And if one in four DB students failed too?

Then $P(\text{PML} \mid \text{fail}) = 0.4$: the result says nothing about the course.

Takeaway

> Two rules — **sum** to remove a variable, **product** to split a joint — generate joint, marginal, conditional probabilities and Bayes' theorem. What follows makes them precise and extends them beyond counting.

Random Experiments and Events

> ***Guiding Question:*** *Before any number: what exactly is uncertain, and which questions about it can be asked at all?*

What Is a Random Experiment?

A coin flip, tomorrow's rainfall, whether a user clicks a link. What makes each of them a *random experiment* — and what would not be one?

What Is a Random Experiment?

A coin flip, tomorrow's rainfall, whether a user clicks a link. What makes each of them a *random experiment* — and what would not be one?

> **Definition (Random experiment).** A procedure that can — at least in theory — be repeated infinitely often, characterised by:
>
> 1. **Configuration** — a precisely specified system that can be reconstructed.
> 2. **Procedure** — an instruction for executing the experiment.
> 3. **Unpredictability** of the outcome.

Coin flip: a 1-Euro coin (configuration), placed on the thumb and snapped into the air (procedure), record heads or tails (outcome).


Randomness as Missing Information

> A random experiment is still causal. Its unpredictability reflects **missing information about the causal chain**, not an absence of causes.
>
> Consequence: a random experiment can become a deterministic process once new information is available.

A coin flip filmed at 10,000 fps and measured precisely enough stops being random. What does that imply for a model that calls something "random"?

Sample Space and Events

Definition (Sample space, event space). A set $\Omega = \{\omega_1, \ldots, \omega_n\}$ is the **sample space** of a random experiment if every outcome is associated with exactly one element $\omega \in \Omega$, an **outcome**.
For finite Ω, every subset $A \subseteq \Omega$ is an **event**; A **occurs** iff the observed $\omega \in A$. All events together, $\mathcal{P}(\Omega)$, form the **event space**.




Ω	sample space: all outcomes
ω = 6	an outcome: an element
A = {4, 5, 6}	an event: a subset

ω ∈ A: the event "at least 4" occurs

An outcome is an element; an event is a subset — a *question you can ask* about the result.


Important Event Types



$\varnothing$

impossible event


no outcome belongs to it



Ω

certain event


every outcome belongs to it



$\bar{A} = \Omega \setminus A$

complementary event


A = {1, 2}, $\bar{A}$ = {3, 4, 5, 6}



$|A| = 1$

elementary event


A = {5}: one outcome



$A \subseteq B$

A entails B

A = {1, 2}, B = {1, 2, 4, 5}




$A \cap B = \varnothing$

incompatible events

A = {1, 2}, B = {3, 6}

Examples


One coin flip



E = {H}: "heads"

events: ∅, {H}, {T}, {H, T}


Two coin flips: the set



E = {HH, HT, TH}: "at least one head"

4 outcomes, $2^4 = 16$ events


… as a grid: 1st × 2nd



same outcomes, arranged by flip

The same event, twice: as a subset of Ω, and as cells of the 1st × 2nd grid. The sample space fixes which questions can be asked.

The Event Space Grows Fast



$|\Omega| = n \rightarrow 2^n$ events

1	2
2	4
3	8
6	64
10	1,024
20	1,048,576

lines: one element added

For $|\Omega| = n$ the event space has 2^n elements — far too many to list, which is why probability is done by counting, not by enumeration.

Not Every Sample Space Is Finite

Dosage of a drug: Ω = (0, ∞)



uncountably many outcomes — probability can only live on intervals, not on points

Race with 7 teams: all orderings




7! = 5,040 outcomes; shaded: the event "team 3 wins"

Choosing Ω is a modelling decision: it fixes what counts as an outcome, and therefore which questions can be asked at all.

Venn Diagrams

Venn diagrams are the graphical representation of events:




- the **rectangle** is the sample space Ω
- a **circle** is an event $A \subseteq \Omega$
- every **region** is a set — and every set operation is a region you can point at

Important Event Types

Before each step: which region will light up?

$A \cap B = \emptyset$



A and B are disjoint


Algebra of Events

Law	Union form	Intersection form
Commutative	$A \cup B = B \cup A$	$A \cap B = B \cap A$
Associative	$(A \cup B) \cup C = A \cup (B \cup C)$	$(A \cap B) \cap C = A \cap (B \cap C)$
Distributive	$(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$	$(A \cap B) \cup C = (A \cup C) \cap (B \cup C)$

Events form a set algebra — the same laws you know, but now every identity is a picture.

Set Operations as Regions

A ∪ B ∪ C




Union of three events

Mini-Exercise


Are $(A \cup B) \cap C$ and $(A \cap B) \cup C$ the same region?

Mini-Exercise

Are $(A \cup B) \cap C$ and $(A \cap B) \cup C$ the same region?



inside C only



all of C, plus the lens A ∩ B


No. The first lies entirely inside C; the second is all of C plus the lens $A \cap B$. The distributive laws expand them into different expressions.

The Probability Measure

> ***Guiding Question:*** *We need a language that can say "how strongly", not just "true or false". What must such a language satisfy — and who decides the numbers?*

An Intuition: Probability as Area

Throw 40 stones at random into Ω



P(A) = area of A / area of Ω ≈ 11 of 40 stones


counted out of

Draw the sample space Ω on the ground, with each event as a region. Throw a stone at random.

How likely does it land in A?

An Intuition: Probability as Area

Throw 40 stones at random into Ω



P(A) = area of A / area of Ω ≈ 11 of 40 stones

counted out of

Draw the sample space Ω on the ground, with each event as a region. Throw a stone at random.

How likely does it land in A?

$$P(A) = \frac{\text{area of } A}{\text{area of } \Omega}$$


provided that

1. the stone always lands in Ω, and
2. every part of Ω is equally likely.

Probability measures **how much of** Ω an event occupies.

An Intuition: Probability as Area

Throw 40 stones at random into Ω



P(A) = area of A / area of Ω ≈ 11 of 40 stones

counted out of

Draw the sample space Ω on the ground, with each event as a region. Throw a stone at random.

How likely does it land in A?

$$P(A) = \frac{\text{area of } A}{\text{area of } \Omega}$$

provided that

1. the stone always lands in Ω, and
2. every part of Ω is equally likely.

Probability measures **how much of** Ω an event occupies.

> Assumption 2 is the one we will have to give up.

Two Ways to Read a Probability

Frequentist
The probability of an elementary event a is the fraction of repetitions of the experiment that yield a.
Requires the experiment to be repeatable.

Bayesian
The probability of a is the degree of certainty that a will happen — even without conducting the experiment.
Requires a stated state of knowledge.

Both interpretations yield the same mathematics. They are mental models for what the number *means*, not different theories.

The Classical (Laplace) Definition

Definition (Classical / Laplace probability). If every elementary event in Ω is assigned the same probability, then

$$P(A) = \frac{|A|}{|\Omega|} = \frac{\text{number of cases favourable for } A}{\text{number of possible outcomes}}$$

An experiment whose configuration and procedure make outcomes equiprobable — by definition or by construction — is a **Laplace experiment**; its probabilities need not be estimated, they are implied by the setup.

Why That Definition Is Not Enough

> **The circularity.** "Laplace probability" defines probability using *equiprobability* — that is, using another notion of probability. Strictly speaking it is a circular definition.

The frequentist repair — defining probability as the limit of relative frequencies — fails too: such a limit exists only in mathematical settings, never in an experiment that can actually be repeated.

So what do we do? Stop trying to *define* what probability is, and instead specify what it must *do*.

The Kolmogorov Axioms

Definition (Probability measure, Kolmogorov 1933). Let Ω be a sample space and $\mathcal{P}(\Omega)$ its event space. A function $P : \mathcal{P}(\Omega) \to \mathbb{R}$ is a **probability measure** if

I	$P(A) \geq 0$	non-negative
II	$P(\Omega) = 1$	normalised
III	$A \cap B = \emptyset \Rightarrow P(A \cup B) = P(A) + P(B)$	additive

INTUITION

Probability is **one kilogram of sand** spread over Ω; $P(A)$ is the sand lying in A.

- **I** — no region holds negative sand.
- **II** — all the sand lies somewhere in Ω.
- **III** — two regions that do not overlap hold their sand together.

The axioms say how *any* assignment must behave — not which probabilities the events get. How the sand is spread is the modeller's choice.

For infinite Ω the event space is a σ-algebra and Axiom III holds for countably many disjoint events — measure theory; nothing in this course depends on it.

Probability Space

Definition. Let Ω be a sample space, $\mathcal{P}(\Omega)$ an event space, and P a probability measure. The triple $(\Omega, \mathcal{P}(\Omega), P)$ — also written (Ω, P) — is a **probability space**.

INTUITION

Three separate modelling choices: Ω — **what can happen** (the ground); $\mathcal{P}(\Omega)$ — **what may be asked** (the regions one may draw); P — **how likely each answer is** (how the sand is spread).

Same Ω, different P

A fair and a loaded die share $\Omega = \{1, \ldots, 6\}$ but are two probability spaces.

Finite Ω

P is fixed by the elementary events: $p_\omega \geq 0$, $\sum_\omega p_\omega = 1$, and $P(A) = \sum_{\omega \in A} p_\omega$.

What the Axioms Already Give Us

Property	Statement	From	Intuitively
Complement	$P(\overline{A}) = 1 - P(A)$	II, III	*the sand not in A lies in $\overline{A}$*
Empty event	$P(\emptyset) = 0$	complement, $A = \Omega$	*no region, no sand*
Bounds	$0 \leq P(A) \leq 1$	I, complement	*no region holds more than all the sand*
Monotonicity	$A \subseteq B \Rightarrow P(A) \leq P(B)$	I, III	*a larger region holds at least as much*
Inclusion–exclusion	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$	III	*the overlap was weighed twice*
Union bound	$P(A \cup B) \leq P(A) + P(B)$	I, inclusion–exclusion	*weighing twice never weighs less*
Finite additivity	$P(A_1 \cup \ldots \cup A_p) = \sum_i P(A_i)$ for disjoint A_i	III, repeated	*disjoint regions add up*

Takeaway

> Probability is not defined by counting or by repetition — it is any assignment of numbers to events that is **non-negative, normalised, and additive**. The Laplace formula is one such assignment, not the definition.

Conditional Probability

> ***Guiding Question:*** *We learn that B happened. What becomes of every other probability — and which rules follow from that one step?*

Motivating Example

You roll a 6-sided die twice. **After the first roll you see a 3.** Knowing this, what is the probability that both rolls sum to 8?

Motivating Example

You roll a 6-sided die twice. **After the first roll you see a 3.** Knowing this, what is the probability that both rolls sum to 8?


- Before the first roll: $\Omega = \{(1,1),(1,2),\ldots,(6,6)\}$, $|\Omega| = 36$
- After seeing the 3, only $\Omega_{red} = \{(3,1),\ldots,(3,6)\}$ remains
- Outcomes like $(2,4)$ have become **impossible**
- The reduced space now carries all the probability: $P(\Omega_{red}) = 1$
- "Sum is 8" inside it: $E = \{(3,5)\}$

$$P = \frac{1}{6}$$

Definition

Definition (Conditional probability of A under the condition B).


$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}, \qquad \text{if } P(B) > 0$$



Back to the stone: once it is known to lie in B, B **is the whole world**.

Conditioning, Drawn

Conditioning on B = keep B, stretch it back to width 1



P(A) = 0.40

P(B) = 0.10 + 0.06 = 0.16

P(A ∩ B) = 0.10

P(A | B) = 0.10 / 0.16 = 0.63


P(A | B) = 0.63

Before the last click: what share of B is A?

Then: set $P(A \cap B) = P(B)$. Now $P(A \mid B)$?

Conditioning, Drawn

Conditioning on B = keep B, stretch it back to width 1



P(A) = 0.40

P(B) = 0.10 + 0.06 = 0.16

P(A ∩ B) = 0.10

P(A | B) = 0.10 / 0.16 = 0.63

P(A | B) = 0.63


Before the last click: what share of B is A?

$P(A \cap B)/P(B) \approx 0.63$

Then: set $P(A \cap B) = P(B)$. Now $P(A \mid B)$?

Conditioning, Drawn

Conditioning on B = keep B, stretch it back to width 1



P(A) = 0.40

P(B) = 0.10 + 0.06 = 0.16

P(A ∩ B) = 0.10

P(A | B) = 0.10 / 0.16 = 0.63

P(A | B) = 0.63

Before the last click: what share of B is A?

$P(A \cap B)/P(B) \approx 0.63$

Then: set $P(A \cap B) = P(B)$. Now $P(A \mid B)$?

1: all of B lies inside A.

Example: Two Coins

Two fair coins are flipped. How likely is B = **both heads**, if we learn …

(a) "The first coin is heads"

$F = \{hh, ht\}$

(b) "At least one coin is heads"

$A = \{hh, ht, th\}$

Example: Two Coins

Two fair coins are flipped. How likely is B = **both heads**, if we learn …

(a) "The first coin is heads"

$F = \{hh, ht\}$

$$P(B \mid F) = \frac{P(B \cap F)}{P(F)} = \frac{1/4}{1/2} = \frac{1}{2}$$

(b) "At least one coin is heads"

$A = \{hh, ht, th\}$

Example: Two Coins

Two fair coins are flipped. How likely is B = **both heads**, if we learn …

(a) "The first coin is heads"

$F = \{hh, ht\}$

$$P(B \mid F) = \frac{P(B \cap F)}{P(F)} = \frac{1/4}{1/2} = \frac{1}{2}$$

(b) "At least one coin is heads"

$A = \{hh, ht, th\}$

$$P(B \mid A) = \frac{P(B \cap A)}{P(A)} = \frac{1/4}{3/4} = \frac{1}{3}$$

Example: Two Coins

Two fair coins are flipped. How likely is B = **both heads**, if we learn …

(a) "The first coin is heads"

$F = \{hh, ht\}$

$$P(B \mid F) = \frac{P(B \cap F)}{P(F)} = \frac{1/4}{1/2} = \frac{1}{2}$$


(b) "At least one coin is heads"

$A = \{hh, ht, th\}$

$$P(B \mid A) = \frac{P(B \cap A)}{P(A)} = \frac{1/4}{3/4} = \frac{1}{3}$$


Same event, similar news. **Why is (b) smaller?**

Two Coins, Drawn



Both count the same outcome hh. News (a) leaves **two** possible outcomes, news (b) leaves **three**: weaker news rules out less, so hh gets a smaller share.

Two Coins, Drawn



Both count the same outcome hh. News (a) leaves **two** possible outcomes, news (b) leaves **three**: weaker news rules out less, so hh gets a smaller share.

> Pictures stop at two or three events. **Learn the rule** — it does not run out.

Total Probability


Theorem (Total probability). Let $E_1, \ldots, E_k$ be mutually exclusive with $\Omega = E_1 \cup \ldots \cup E_k$ and $P(E_i) > 0$. Then for any event B:

$$P(B) = \sum_{i=1}^{k} P(E_i) \cdot P(B \mid E_i)$$

Proof. Decompose B along the partition and use additivity:

$$P(B) = P((E_1 \cup \ldots \cup E_k) \cap B) = \sum_{i=1}^{k} P(E_i \cap B) = \sum_{i=1}^{k} P(E_i) \cdot P(B \mid E_i)$$

Total Probability, Drawn



$$P(B) = P(B \cap E_1) + P(B \cap E_2) + P(B \cap E_3) = \Sigma_i\, P(E_i) \cdot P(B \mid E_i)$$

B is cut into disjoint pieces, one per E_i; each piece is "first E_i, then B within E_i" — the product rule.

Rules That Follow

Product rule

$$P(A \cap B) = P(B) \cdot P(A \mid B) = P(A) \cdot P(B \mid A)$$

Complement, same condition

$$P(\overline{A} \mid B) = 1 - P(A \mid B)$$


But in general

$$P(A \mid \overline{B}) \neq 1 - P(A \mid B)$$

Held fixed at B, $P_B(E) = P(E \mid B)$ for every event E satisfies the Kolmogorov axioms — conditioning yields a **new probability measure** P_B.


Conditioning Gives a New Measure

$P_B(E) = P(E \cap B) / P(B)$: every event E is measured by its piece inside B




Axiom I: $P_B(E) \geq 0$

a piece of B is never negative



Axiom II: $P_B(\Omega) = 1$

all of B is the whole world




Axiom III: $P_B(E_1 \cup E_2) = P_B(E_1) + P_B(E_2)$

disjoint pieces of B add up

Inside B, the three axioms hold again — so **every rule proved for P holds for P_B**.

Chain Rule

$$P(E_1 \cap E_2 \cap \ldots \cap E_n) = P(E_1) \cdot P(E_2 \mid E_1) \cdot P(E_3 \mid E_1 \cap E_2) \cdots P(E_n \mid E_1 \cap \ldots \cap E_{n-1})$$



Each factor keeps a share of what is left. Any joint probability can be written as a chain of conditionals. Choosing a *good* order — and dropping conditions that do not matter — is exactly what graphical models will do later.

Takeaway

> Conditioning **shrinks the sample space and renormalises**. Everything else — product rule, total probability, Bayes, the chain rule — is bookkeeping on that single move.

Bayes' Theorem

> ***Guiding Question:*** *A 99%-accurate test says you are sick. Why is that not a 99% chance of being sick — and how do we turn* $P(+ \mid \text{sick})$ *into* $P(\text{sick} \mid +)$*?*

Bayes' Theorem — An Intuition First

A test for a rare disease is **99% accurate**: positive for 99% of the sick, negative for 99% of the healthy. About **1 in 1,000 people** has the disease.

Your test comes back positive. How likely is it that you are actually sick?

Bayes' Theorem — An Intuition First

A test for a rare disease is **99% accurate**: positive for 99% of the sick, negative for 99% of the healthy. About **1 in 1,000 people** has the disease.

Your test comes back positive. How likely is it that you are actually sick?

> Only about **9%** — the answer most people miss by a factor of ten.

Why? Count People, Not Percentages

Restricting to the positives is exactly the conditioning step

P(sick | positive) = 1/11 ≈ 9%



What Just Happened

Restricting attention to the people who tested positive is exactly the move from the conditional-probability definition:

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$$

The 11 positives are the new sample space B; the single sick person among them is $A \cap B$. The rarity of the disease survives the evidence because there were so many more healthy people to misclassify.

The trap

"99% accurate" describes $P(\text{positive} \mid \text{sick})$.

You wanted $P(\text{sick} \mid \text{positive})$.

These are different numbers.

Bayes' Theorem — Two Events

Theorem (Bayes, 2 events). Let $(\Omega, \mathcal{P}(\Omega), P)$ be a probability space and $A, B \in \mathcal{P}(\Omega)$ two events with $P(B) > 0$. Then:

$$P(A \mid B) = \frac{P(B \mid A) \cdot P(A)}{P(B)}$$

posterior — what we believe after

likelihood — how well A explains B

evidence — how ordinary B is

prior — belief before the evidence

Which of the four was "99% accurate", and which one did the intuition ignore?

Bayes' Theorem — Two Events

Theorem (Bayes, 2 events). Let $(\Omega, \mathcal{P}(\Omega), P)$ be a probability space and $A, B \in \mathcal{P}(\Omega)$ two events with $P(B) > 0$. Then:

$$P(A \mid B) = \frac{P(B \mid A) \cdot P(A)}{P(B)}$$

posterior — what we believe after

likelihood — how well A explains B

evidence — how ordinary B is


prior — belief before the evidence

Which of the four was "99% accurate", and which one did the intuition ignore?

"99% accurate" is the **likelihood** $P(+ \mid \text{sick})$; the intuition ignored the **prior** $P(\text{sick}) = 0.001$.

Proof — Two Events

Intuition



One overlap $A \cap B$, measured against two wholes: against B it is $P(A \mid B)$, against A it is $P(B \mid A)$.

Proof

step	because
$P(A \mid B) = \frac{P(A \cap B)}{P(B)}$	definition
$= \frac{P(A \cap B)}{P(A)} \cdot \frac{P(A)}{P(B)}$	multiply and divide by $P(A)$
$= \frac{P(B \mid A)\, P(A)}{P(B)}$	definition, roles swapped

Needs $P(A), P(B) > 0$. ■

Mini-Exercise

A high-risk group. Same test: positive for **99%** of the sick, for **1%** of the healthy. Now **1 in 20** is sick. Above or below 9%? Guess, then count 2,000 people.

Mini-Exercise

A high-risk group. Same test: positive for **99%** of the sick, for **1%** of the healthy. Now **1 in 20** is sick. Above or below 9%? Guess, then count 2,000 people.

- 100 sick → **99** positive; 1,900 healthy → **19** positive.
- 99 of all **118** positives are sick: $P(\text{sick} \mid +) \approx 84\%$.

Mini-Exercise

A high-risk group. Same test: positive for **99%** of the sick, for **1%** of the healthy. Now **1 in 20** is sick. Above or below 9%? Guess, then count 2,000 people.

- 100 sick → **99** positive; 1,900 healthy → **19** positive.
- 99 of all **118** positives are sick: $P(\text{sick} \mid +) \approx 84\%$.

	everyone	high-risk group
the test, $P(+ \mid \text{sick})$ / $P(+ \mid \text{healthy})$	99% / 1%	99% / 1%
who is tested, $P(\text{sick})$	1 in 1,000	1 in 20
$P(\text{sick} \mid +)$	$\approx 9\%$	$\approx 84\%$

Mini-Exercise

A high-risk group. Same test: positive for **99%** of the sick, for **1%** of the healthy. Now **1 in 20** is sick. Above or below 9%? Guess, then count 2,000 people.

- 100 sick → **99** positive; 1,900 healthy → **19** positive.
- 99 of all **118** positives are sick: $P(\text{sick} \mid +) \approx 84\%$.

	everyone	high-risk group
the test, $P(+ \mid \text{sick})$ / $P(+ \mid \text{healthy})$	99% / 1%	99% / 1%
who is tested, $P(\text{sick})$	1 in 1,000	1 in 20
$P(\text{sick} \mid +)$	$\approx 9\%$	$\approx 84\%$

> Both answers are right. Same test, same rule — **different people tested**.

The Same Answer, by Bayes' Theorem

Which known number goes where?

$$P(\text{sick} \mid +) = \frac{\overbrace{P(+ \mid \text{sick})}^{\text{likelihood: } 0.99} \cdot \overbrace{P(\text{sick})}^{\text{prior: } 0.05}}{\underbrace{P(+ \mid \text{sick})\,P(\text{sick})}_{\text{sick and positive: } 0.99 \cdot 0.05 = 0.0495} + \underbrace{P(+ \mid \text{healthy})\,P(\text{healthy})}_{\text{healthy and positive: } 0.01 \cdot 0.95 = 0.0095}}$$

The Same Answer, by Bayes' Theorem

Which known number goes where?

$$P(\text{sick} \mid +) = \frac{\overbrace{P(+ \mid \text{sick})}^{\text{likelihood: } 0.99} \cdot \overbrace{P(\text{sick})}^{\text{prior: } 0.05}}{\underbrace{P(+ \mid \text{sick})\,P(\text{sick})}_{\text{sick and positive: } 0.99 \cdot 0.05 = 0.0495} + \underbrace{P(+ \mid \text{healthy})\,P(\text{healthy})}_{\text{healthy and positive: } 0.01 \cdot 0.95 = 0.0095}}$$

$$= \frac{0.0495}{0.0495 + 0.0095} = \frac{0.0495}{0.0590} \approx 0.84$$

The denominator is the **evidence** $P(+)$: every way to test positive.

The Same Answer, by Bayes' Theorem

Which known number goes where?

$$P(\text{sick} \mid +) = \frac{\overbrace{P(+ \mid \text{sick})}^{\text{likelihood: } 0.99} \cdot \overbrace{P(\text{sick})}^{\text{prior: } 0.05}}{\underbrace{P(+ \mid \text{sick})\,P(\text{sick})}_{\text{sick and positive: } 0.99 \cdot 0.05 = 0.0495} + \underbrace{P(+ \mid \text{healthy})\,P(\text{healthy})}_{\text{healthy and positive: } 0.01 \cdot 0.95 = 0.0095}}$$

$$= \frac{0.0495}{0.0495 + 0.0095} = \frac{0.0495}{0.0590} \approx 0.84$$

The denominator is the **evidence** $P(+)$: every way to test positive.

> Times 2,000, the fraction is **99 / 118**. Counting people and the formula are **one calculation**, written two ways.

Bayes' Theorem — Multiple Events

Theorem (Bayes, multiple events). Let $E_1, \ldots, E_k$ be mutually exclusive with $\Omega = E_1 \cup \ldots \cup E_k$ and $P(E_i) > 0$, and let B be an event with $P(B) > 0$. Then

$$P(E_i \mid B) = \frac{P(B \mid E_i) \cdot P(E_i)}{P(B)} = \frac{P(B \mid E_i) \cdot P(E_i)}{\sum_{j=1}^{k} P(B \mid E_j) \cdot P(E_j)}$$

posterior of E_i

evidence

every E_i explains B, weighted by its prior


likelihood — how well E_i explains B

prior of E_i

The denominator is the **total probability** of the evidence.

Proof — Multiple Events

Intuition



The events E_i cut B into pieces. The posterior of E_i is **its piece out of all pieces**.

Proof

step	because
$P(E_i \mid B) = \frac{P(B\|E_i)\,P(E_i)}{P(B)}$	Bayes, two events
$P(B) = \sum_j P(B \cap E_j)$	additivity
$= \sum_j P(B \mid E_j)\,P(E_j)$	product rule

Additivity holds because the E_j are disjoint and cover Ω. Insert $P(B)$ into the first line. ■

Takeaway

Bayes' theorem is the arithmetic of changing your mind: **posterior ∝ likelihood × prior**. Evidence never speaks alone — it is always weighed against how plausible the hypothesis was to begin with.

Back to Lecture 01: the Model Says "Fraud"

A fraud model is **99% accurate** in both directions. One transaction in **1,000** is fraudulent. Per 100,000 transactions:

	flagged	not flagged	total
fraud	99	1	100
legitimate	999	98,901	99,900
total	**1,098**	98,902	100,000

Back to Lecture 01: the Model Says "Fraud"

A fraud model is **99% accurate** in both directions. One transaction in **1,000** is fraudulent. Per 100,000 transactions:

	flagged	not flagged	total
fraud	99	1	100
legitimate	999	98,901	99,900
total	**1,098**	98,902	100,000

$$P(\text{fraud} \mid \text{flagged}) = \frac{99}{1{,}098} \approx 9\%$$

> Nine in ten blocked cards belong to honest customers — from a "99% accurate" model. The label "fraud" hid the **prior**; a score would at least have carried it.

Independence

> ***Guiding Question:*** *When does learning one thing tell us nothing about another — and why is that the assumption every tractable model leans on?*

Independence

Definition (Statistical independence). Two events A and B are **independent** iff

$$P(A \cap B) = P(A) \cdot P(B)$$

If additionally $0 < P(B) < 1$, this is equivalent to

$$P(A \mid B) = P(A) \qquad \text{and} \qquad P(A \mid B) = P(A \mid \overline{B})$$

Independence means: **learning B changes nothing about A.**

The Classic Confusion


> **Statistical independence does not mean the events are disjoint.**

Disjoint events with positive probability are about as **dependent** as events can be: if A occurs, B certainly did not.

$$A \cap B = \emptyset \Rightarrow P(A \mid B) = 0 \neq P(A)$$

Independence Is About Areas, Not Overlap

P(A) · P(B) = 0.30 · 0.40 = 0.12 = P(A ∩ B) → independent



P(A) = 0.30

P(B) = 0.40


P(A ∩ B) = 0.12

P(A | B) = 0.12 / 0.40 = 0.30 = P(A)

Independent: **knowing B tells nothing about A.** A takes the same share of B as of all of Ω, so its probability stays the same.

The Same Picture, Dependent

P(A) · P(C) = 0.30 · 0.40 = 0.12 ≠ 0.20 = P(A ∩ C) → dependent



P(A) = 0.30

P(C) = 0.40

P(A ∩ C) = 0.20

P(A | C) = 0.20 / 0.40 = 0.50 ≠ P(A)

Here $P(A) \cdot P(C) = 0.12$ but $P(A \cap C) = 0.20$: the intersection is **larger** than independence would predict — A and C occur together more often than chance.

More Than Two Events

Events $A_1, \ldots, A_n$ are (mutually) independent iff for **every** subset of indices $i_1, \ldots, i_l$:

$$P(A_{i_1} \cap \ldots \cap A_{i_l}) = P(A_{i_1}) \cdots P(A_{i_l})$$

Pairwise independence is **not** enough — every subset must factorise.

Takeaway

> Independence is a statement about **numbers**, not pictures: $P(A \cap B)$ equals the product of the marginals. Disjoint is not independent, and identical marginals can hide either.

Random Variables

> ***Guiding Question:*** *We rarely care which outcome occurred, only what it was worth. How do outcomes become numbers — and what is left of the probability measure?*

Why Random Variables?

Three fair coins: what matters is rarely *which* sequence occurred, but the **number of heads** Y.

Y	outcomes ω with $Y(\omega) = k$	count	$P(Y = k)$
0	TTT	1	$\frac{1}{8}$
1	HTT · THT · TTH	3	$\frac{3}{8}$
2	HHT · HTH · THH	3	$\frac{3}{8}$
3	HHH	1	$\frac{1}{8}$

> Y compresses eight outcomes into four numbers — the probabilities come along.

Definition


Definition (Random variable). A random variable X maps outcomes to a measurable range E: $X : \Omega \to E$. Its probabilities are inherited from the underlying probability space:

$$P(X \in S) = P(\{\omega \in \Omega \mid X(\omega) \in S\})$$

A random variable is neither random nor a variable — it is a **function** on the sample space. The randomness lives in ω, not in X.

The Function, Drawn

Three arrows arrive at 1 and at 2: 3/8 each — the whole shape



Examples

experiment	sample space Ω	random variable $X(\omega)$	values
two coins	$\{HH, HT, TH, TT\}$	number of heads	$0, 1, 2$
two dice	pairs $\langle n_1, n_2 \rangle$	sum $n_1 + n_2$	$2, \ldots, 12$
draw a person	a population	height of the person	an interval of $\mathbb{R}$

The same outcome can carry many random variables: one per question asked of it.

Discrete and Continuous

Discrete random variables

Countable range, usually $E \subseteq \mathbb{N}$.

- number of children of a person
- sum of two dice

Continuous random variables

Real-valued range, $E = \mathbb{R}$.

- height of a person
- horsepower of a car

The distinction decides *how* we may describe the probabilities — with a mass function or with a density.

A Random Variable With Infinitely Many Values


Flip a biased coin ($P(H) = p$) until the first head; X is the number of flips needed.

Geometric distribution, $X \sim \mathrm{Geom}(p)$:

$$P(X = 1) = p$$

$$P(X = 2) = (1 - p)p$$

$$P(X = n) = (1 - p)^{n-1}p$$



Countably infinite range, and the probabilities still sum to 1.

Takeaway

A random variable is a **function from outcomes to numbers** that carries the probability measure with it. It lets us stop talking about sample spaces and start talking about quantities — which is what the rest of the course does.

Probability Mass Function

Definition (PMF). For a discrete random variable X, the **probability mass function** is

$$p(a) = P(X = a)$$


INTUITION

$p(a)$ is how much of the total probability sits exactly on the value a.

What follows from the axioms, with nothing new assumed:

- $p(a) > 0$ for at most countably many values a
- $\sum_i p(a_i) = 1$ — the masses add up to the whole
- p defines a probability measure on the values: the **distribution** of X

Example: Sum of Two Dice



The triangular shape is counting, drawn: six of the 36 outcomes give 7, only one gives 2.

Expectation

Definition (Expected value).

$$E[X] = \sum_i a_i \cdot p(a_i)$$

INTUITION

The **average outcome** if the experiment were repeated infinitely often — the balance point of the bar chart of p.

function of X	$E[g(X)] = \sum_i g(a_i)p(a_i)$
linear transformation	$E[aX + b] = aE[X] + b$
first moment	$\mu = E[X]$; the first *central* moment $E[X - \mu] = 0$

Variance

Definition (Variance).

$$\mathrm{Var}(X) = E\big[(X-\mu)^2\big] = E[X^2] - \big(E[X]\big)^2$$

Spread

The second central moment: the expected squared distance from the mean. Larger variance, larger typical deviation.

Scale and unit

$\mathrm{Var}(aX+b) = a^2\mathrm{Var}(X)$

$\sigma = \sqrt{\mathrm{Var}(X)}$, the **standard deviation**, has the unit of X.

Takeaway


A discrete distribution is a PMF; expectation and variance compress it into **where it sits** and **how much it spreads**.

Bernoulli and Binomial

	models	PMF	$E[X]$	$\mathrm{Var}(X)$
Bernoulli(p)	one trial: $X = 1$ success, $X = 0$ failure	$p(x) = p^x(1-p)^{1-x}$	p	$p(1-p)$
Binomial $B(n, p)$	successes in n independent trials	$p(k) = \binom{n}{k} p^k (1-p)^{n-k}$	np	$np(1-p)$

$\binom{n}{k} = \frac{n!}{k!(n-k)!}$ counts the sequences with k successes; the PMF multiplies it by the probability of one such sequence — combinatorics and independence in a single line.

How the Shape Depends on the Parameters



Predict before each click: as n grows, does the distribution get wider or narrower — and what happens to its shape?

Mini-Exercise

A student guesses all **10** questions of a quiz, each with **4** options.

How many correct answers should they expect — and is 5 or more likely?

Mini-Exercise

A student guesses all **10** questions of a quiz, each with **4** options.

How many correct answers should they expect — and is 5 or more likely?

$B(10, 0.25)$: on average $10 \cdot 0.25 = 2.5$ correct. Five or more is twice the expectation — unlikely (about 8%).

Takeaway

A yes/no attribute is a **Bernoulli** variable with one parameter p; counting the yeses in n independent trials gives a **binomial**. Both are counting in probabilistic clothing — and p is the number data will have to tell us.


From Mass to Density

Definition (PDF). A continuous random variable X has a **probability density function** f with

$$P(b \leq X \leq c) = \int_b^c f(a)\,da, \qquad \int_{-\infty}^{\infty} f(a)\,da = 1$$

$f(a)$ is **not** a probability — it is probability *per unit interval*. Consequently $f(a) > 1$ is perfectly possible, and $P(X = a) = 0$ for every single value a.

Probability Is Area Under the Curve



The shaded region is $P(-1 \le X \le 1)$ — the same "probability as area" intuition from the axioms unit, now with a curve instead of a Venn diagram.

Expectation for Continuous Variables


	discrete — mass p	continuous — density f
total	$\sum_i p(a_i) = 1$	$\int_{-\infty}^{\infty} f(a)\,da = 1$
expectation	$E[X] = \sum_i a_i p(a_i)$	$E[X] = \int_{-\infty}^{\infty} a f(a)\,da$
function of X	$E[g(X)] = \sum_i g(a_i) p(a_i)$	$E[g(X)] = \int_{-\infty}^{\infty} g(a) f(a)\,da$

Only the sum turns into an integral. The rules are unchanged: $E[aX + b] = aE[X] + b$ and $\mathrm{Var}(X) = E[(X - \mu)^2]$.

Two Basic Continuous Distributions

Uniform on $[a, b]$


$$f(x) = \frac{1}{b-a} \text{ for } x \in [a, b]$$



Equal-length intervals are equally likely — the Laplace assumption, continuous.


Normal $N(\mu, \sigma^2)$

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$



$E[X] = \mu$, $\mathrm{Var}(X) = \sigma^2$; $N(0, 1)$ is the **standard** normal.

What σ Does to the Shape



The peak falls as the curve widens — the area is always 1.

Why the Normal Is Everywhere

> The **central limit theorem**: under mild conditions, the average of n independent observations is approximately normal for large n — whatever the original distribution was.

This is why the normal is the default when the true distribution is unknown — and why a sum of many independent coin flips, the binomial, looks bell-shaped.

Other Distributions You Will Meet

distribution	values	typically models
Exponential	$[0, \infty)$	time until an event occurs (e.g. the next earthquake)
Gamma	$[0, \infty)$	time until n events have occurred
Beta	$[0, 1]$	an unknown probability
Dirichlet	probability vectors	distributions *over* probabilities


Continuous distributions replace mass with **density** and sums with **integrals**. Probability remains area, expectation remains a weighted average, and the normal is the default shape because of the central limit theorem.

Joint Distributions

> ***Guiding Question:*** *Never one quantity alone. How do we describe several random variables at once — the joint the three moves start from?*

More Than One Random Variable

Univariate




p(x) — one column

Bivariate



p(x, y) — two columns

Multivariate



p(x, y, z, …) — three or more


A **joint distribution** describes the outcomes of several random variables *at the same time* — which is what every dataset with more than one column is.

Joint Mass and Density

joint PMF (discrete)	joint PDF (continuous)
$p_{X,Y}(a,b) = P(X = a, Y = b)$	$P((X,Y) \in R) = \iint_R f_{X,Y}(a,b)\,da\,db$

	G=0	G=1	G=2	G=3
B=0	.15	.1	.0875	.0375
B=1	.1	.175	.1125	0
B=2	.0875	.1125	0	0
B=3	.0375	0	0	0

one probability per cell · all sum to 1



darker = denser · volume over R ≈ 0.33

From here on we write $p(a, b, c, \ldots)$ for the joint of $X_1, X_2, X_3, \ldots$ — a mass or a density, as the variables require.


Marginalisation

Marginalisation eliminates a random variable by summing over all its states. For $p(x_1, \ldots, x_n)$, a **marginal** over a subset is

$$p(x_1) = \sum_{x_2} \cdots \sum_{x_n} p(x_1, x_2, \ldots, x_n)$$

INTUITION

Forget about a variable, but keep its influence correctly accounted for: collapse the joint onto the axis of the variable you keep.



Example: Boys and Girls

	G=0	G=1	G=2	G=3	p(B)
B=0	.15	.1	.0875	.0375	.375
B=1	.1	.175	.1125	0	.3875
B=2	.0875	.1125	0	0	.2
B=3	.0375	0	0	0	.0375
p(G)	.375	.3875	.2	.0375	total 1

Families: 15% have no children, 20% one, 35% two, 30% three. Each child is a boy or a girl with probability $\frac{1}{2}$, independently.

B boys, G girls.

> The margins of the table *are* the marginals — hence the name.

Covariance

$$\operatorname{cov}(A, B) = E\big[(A - E[A])(B - E[B])\big]$$

$\operatorname{cov} > 0$
A large when B large — positively related


$\operatorname{cov} < 0$
A large when B small — negatively related

$\operatorname{cov}(A, A) = \operatorname{Var}(A)$
variance is self-covariance

Independence implies zero covariance. **The converse is false** — covariance only detects *linear* association.

Covariance Sees Only One Kind of Structure

Slide ρ, then switch the relation. Predict the covariance first.



Predict first: with $Y = X^2 - 1$, every point is fixed by its x. The covariance?

Takeaway

The joint distribution holds everything; **marginalisation** sums a variable away, **conditioning** restricts, and **covariance** summarises linear co-variation. Zero covariance is weaker than independence.

Data as a Sample

> ***Guiding Question:*** *A table is all we ever get. Which joint distribution could have produced it — and what else went into the recorded values?*

Four Attribute Types — What the Values Allow

	type	the values…	meaningful	statistics	examples
categorical	**nominal**	are names that only distinguish objects	$=, \neq$	mode, entropy, contingency table, χ^2 test	ZIP code, employee ID, eye colour
	ordinal	carry an order	$<, >$	median, percentiles, rank correlation, sign test	grades, rankings, {good, better, best}
numeric	**interval**	have meaningful differences; the zero is arbitrary	$+, -$	mean, standard deviation, Pearson correlation, t-test	temperature in °C or °F, calendar dates
	ratio	have a true zero, so ratios are meaningful	$\times, \div$	geometric mean, harmonic mean, coefficient of variation	length, mass, counts, age, temperature in K

Each type permits everything the one above it does, plus more.



Four Attribute Types — What You May Transform

	type	permissible transformation	example
categorical	**nominal**	any one-to-one relabelling	reassigning employee IDs changes nothing
	ordinal	any strictly increasing f: $x \mapsto f(x)$	{good, better, best} is as well coded by {1, 2, 3} as by {1, 5, 9}
numeric	**interval**	$x \mapsto ax + b$, $a > 0$	°C and °F differ in zero point and unit
	ratio	$x \mapsto ax$, $a > 0$	length in metres or in feet

INTUITION

A statement is **meaningful** on a scale if no permissible transformation can change its truth. "Twice as long" survives metres → feet; "twice as warm" does not survive °C → °F.

Why the Type Decides the Method

- ✗ **The classic error:** averaging a nominal attribute. The mean of ZIP codes 10115 and 81667 is a number, and it is meaningless.
- ✗ **The subtle one:** 20 °C is not "twice as warm" as 10 °C — interval scales have no true zero, so ratios do not transfer.

> The attribute type determines which **statistics**, which **distance measures** and which **plots** are permissible. Every method in this course depends on it.

Mini-Exercise

A customer table records satisfaction (one to five stars) and year of birth. Which scale is each — and whose **mean** would you not report?

Mini-Exercise

A customer table records satisfaction (one to five stars) and year of birth. Which scale is each — and whose **mean** would you not report?

Satisfaction is **ordinal**: its mean assumes equal gaps between the stars — report the median. Year of birth is **interval**: the mean year is fine, but ratios are not ("born twice as late" means nothing); differences such as age are fine.

Takeaway

> An attribute's **scale of measurement** licenses the operations you may perform on it. Choosing a statistic before checking the scale is the most common silent error in data analysis.

Four Shapes of Data

Record data

Objects × attributes — the familiar table. Rows are *treated as* independent.

Graph data

Objects carry **links** to other objects: web pages, social networks, citations.

Fields / grids

Objects sit in a **spatial** arrangement: images, sensor lattices, simulations.

Spatio-temporal

Position **and** time: monthly land temperatures, traffic movement, trajectories.

The shape decides which methods even apply. A table whose rows are treated as independent can be shuffled without loss; a grid or a time series cannot — the arrangement *is* information. Independence of rows is an assumption, not a property of the file.

Three Data Quality Problems

Noise

Random distortion of measured values. Cannot be removed, only modelled or smoothed.

Outliers

Values far from the rest. Either an **error** to remove or the **signal** you were looking for — fraud, faults, discoveries.

Missing values

Absent entries. Handling depends on **why** they are missing, which the data itself does not tell you.

> "Outlier" is not a property of a data point; it is a judgement about a model of what is normal. Deleting outliers deletes evidence against that model.

Mini-Exercise

A weather sensor misses 8% of its temperature readings, and it fails mostly **when it is very cold**. Is deleting the rows with missing values harmless?

Mini-Exercise

A weather sensor misses 8% of its temperature readings, and it fails mostly **when it is very cold**. Is deleting the rows with missing values harmless?

> No. Deletion removes the coldest days, so every statistic afterwards — the mean temperature first — is biased upwards. Had the gaps been random transmission dropouts, deletion would be roughly harmless. The file looks the same in both cases; only the **mechanism** differs.

Mini-Exercise

A weather sensor misses 8% of its temperature readings, and it fails mostly **when it is very cold**. Is deleting the rows with missing values harmless?

No. Deletion removes the coldest days, so every statistic afterwards — the mean temperature first — is biased upwards. Had the gaps been random transmission dropouts, deletion would be roughly harmless. The file looks the same in both cases; only the **mechanism** differs.

The shape of the data decides which methods apply; the **reason** for a quality problem decides how it may be handled. Neither is visible in the file.

A Table Is a Sample From a Joint Distribution

customer	payment	satisfaction	born	income (k€)	returned?
1	card	4	1987	52	yes
2	invoice	2	1995	31	no
3	card	5	1979	78	no
…	…	…	…	…	…

Each column is a random variable

$X_{\text{payment}}, X_{\text{satisf.}}, \ldots$ — a function from customers to values, exactly as a random variable was defined.

Each row is one draw

The whole table is n draws from one joint distribution $p(x_1, \ldots, x_d)$ that nobody has written down.

The Scale Suggests the Model

attribute	scale	a first model
returned?	nominal, two values	**Bernoulli**(p) — one parameter
payment	nominal, k values	**categorical**$(p_1, \ldots, p_k)$, $\sum_j p_j = 1$
satisfaction	ordinal	categorical — the order is not used, or an ordinal model
born	interval	normal, on a restricted range
income	ratio, positive, skewed	log-normal, or normal on $\log$ income

> The scale limits the model as it limits the statistics: a normal distribution for ZIP codes is as meaningless as their mean.

What "i.i.d." Claims

The rows $\mathbf{x}_1, \ldots, \mathbf{x}_n$ are **independent and identically distributed** (i.i.d.) if

independent	$p(\mathbf{x}_1, \ldots, \mathbf{x}_n) = \prod_i p_i(\mathbf{x}_i)$	*one customer's row tells nothing about another's*
identically distributed	$p_i = p$ for every row i	*every row comes from the **same** joint distribution*

Which of these datasets break it: daily sales of one shop · two visits by the same customer · pixels of one image · a survey sent to a random sample?

What "i.i.d." Claims

The rows $\mathbf{x}_1, \ldots, \mathbf{x}_n$ are **independent and identically distributed** (i.i.d.) if

independent	$p(\mathbf{x}_1, \ldots, \mathbf{x}_n) = \prod_i p_i(\mathbf{x}_i)$	*one customer's row tells nothing about another's*
identically distributed	$p_i = p$ for every row i	*every row comes from the **same** joint distribution*

Which of these datasets break it: daily sales of one shop · two visits by the same customer · pixels of one image · a survey sent to a random sample?

Only the random survey is plausibly i.i.d. Time series, repeated measurements and spatial data all violate it — the arrangement *is* information.

Takeaway

A dataset is n **draws from an unknown joint distribution**: columns are random variables, rows are draws, and the scale of each column suggests its model. Reading a table this way rests on an assumption — **i.i.d.** — that should be checked, not presumed.

Where Does a Recorded Value Come From?

A sensor records 22.4 °C. Besides the temperature, what went into that number?

$$x = s + b + \varepsilon$$

s: the true value

b: artefact — systematic, the same every time

ε: noise — random, mean 0

Where Does a Recorded Value Come From?

A sensor records 22.4 °C. Besides the temperature, what went into that number?

$$x = s + b + \varepsilon$$

s: the true value

b: artefact — systematic, the same every time

ε: noise — random, mean 0

> A dataset records x, never s. Every data problem is a claim about the terms that were added on the way.

Three Problems on One Axis



Which of the three problems does collecting more data fix?

Three Problems on One Axis



Which of the three problems does collecting more data fix?

Only **noise**. An artefact shifts every new reading too; an outlier comes back in the same proportion.

Each Problem Is a Model Statement

problem	in the model	more data	what helps
noise	$\varepsilon \sim p(\varepsilon)$, $E[\varepsilon] = 0$	averages it out	model it; average
artefact	$b \neq 0$, systematic	does **not** help	calibration, a second instrument, domain knowledge
outlier	a draw from another q	a share π remains	robust statistics, or model q: $(1-\pi)\,p + \pi\,q$
missing	M = missing, $P(M \mid x)$	depends on why	ask for the mechanism first

None of these can be read off the data alone. Calling a value noise, an artefact or an outlier is an assumption about how it was generated.

Mini-Exercise

Patients are weighed on three scales; one of them reads 3 kg too high. Does weighing ten times as many patients bring the mean weight closer to the truth?

Mini-Exercise

Patients are weighed on three scales; one of them reads 3 kg too high. Does weighing ten times as many patients bring the mean weight closer to the truth?

> No. If a third of the patients use each scale, the mean stays about 1 kg too high however many are weighed: an **artefact**. More data shrinks only the noise.

Takeaway

A recorded value is **truth + artefact + noise**, or it does not follow the model at all. More data removes **noise only**; artefacts, outliers and missing values each need an assumption about how the data was generated.

Statistics and Estimation

> ***Guiding Question:*** *A model has a parameter nobody told us, and all we have is a sample. What does the sample say about the parameter — and how sure may we be?*


We Never Know the Distribution

Up to now we assumed a distribution was given. In practice it is **unknown**, and we must estimate its properties from data.

	what it is	its numbers	do we see them?
Population	the entire group — all customers, all transactions, all documents	**parameters** μ, σ^2	no — the truth we want
Sample	the observations we actually have	**statistics** $\overline{x}$, s^2	yes — and we estimate the parameters from them

Descriptive statistics summarise a sample, usually while assuming something about the underlying distribution.

What We Summarise



Centre

1st moment

mean, median

Dispersion

2nd central moment

variance, std. dev., IQR

Shape

3rd central moment

skewness, quantiles

Univariate statistics describe one random variable; **multivariate** statistics describe $X_1, \ldots, X_d$ together.

Means

mean	formula	use it for	example
sample	$\overline{x} = \frac{1}{n}\sum_{i=1}^{n} x_i$	values that add up	average salary
weighted	$\overline{x}_w = \frac{\sum_i w_i x_i}{\sum_i w_i}$	values that count unequally	grade over courses of different credits
geometric	$\left(\prod_{i=1}^{n} x_i\right)^{1/n}$	factors that multiply	+50% then −50%: factor 0.87 per period
harmonic	$\frac{n}{\sum_i 1/x_i}$	rates over equal amounts	60 km/h out, 40 back: 48 km/h
trimmed	average after dropping the k smallest and k largest	robustness on the cheap	judges' scores

Median

The **median** is the middle value of the sorted sample: for odd n the $\frac{n+1}{2}$-th value, for even n the average of the two middle values.

INTUITION

Half of the values lie below it, half above. Only their **order** counts, not how far out they lie.


✓ **Robust** against outliers — one extreme value moves the median by at most one position

✗ **Costlier** to compute — it needs sorting or a selection algorithm, not one pass with a running sum

For binned data, the median is interpolated within its interval from the cumulative counts, the interval's lower border and width.

One Value, Two Very Different Reactions

Drag the light point — which marker follows it?



Commit first: drag the top salary to the far right. Which marker follows?

Mini-Exercise

Salaries in a small firm (in k€): 38, 41, 42, 44, 45, 47, 260.

Compute the mean and the median. Which number should appear in a recruitment ad, and which one describes a typical employee?

Mini-Exercise — Solution

Mini-Exercise — Solution

Mean ≈ 73.9, median $= 44$.

The single large value pulls the mean above **every** salary except one. The median describes the typical employee; the mean describes the payroll divided by headcount. Both are correct — they answer different questions.

Dispersion

Sample variance and **standard deviation** of the observed data:

$$s^2 = \frac{1}{n}\sum_{i=1}^{n}(x_i - \overline{x})^2, \qquad s = \sqrt{s^2}$$

INTUITION

The average squared distance from the centre: how stretched or squeezed the data is. s brings it back to the unit of the data.

Computable in **one pass**, $O(n)$, from the running totals n, $\sum x_i$ and $\sum x_i^2$ — which matters when the data does not fit in memory.

Why the Denominator Changes

The **population variance** σ^2 is the variance over the whole population. With the true mean μ unknown, it is estimated by

$$\hat{\sigma}^2 = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \overline{x})^2$$

INTUITION

$\overline{x}$ is computed from the same data, so it sits closer to the data than μ does: the deviations come out **systematically too small**. Dividing by $n-1$ instead of n compensates — one degree of freedom was spent on the mean.

Naming is not uniform: many textbooks, and pandas, call the $n-1$ version s^2; NumPy's `var` divides by n unless told otherwise.

Takeaway

s^2 describes **the sample**; $\hat{\sigma}^2$ estimates **the population**. The $n-1$ is not a convention — it is the price of having estimated the mean from the same data.

You Computed a Number. Which Number Did You Want?

Seven salaries gave a sample mean of $\overline{x} = 73.9$ k€.

What is the probability that the population mean μ is exactly 73.9 k€?

You Computed a Number. Which Number Did You Want?

Seven salaries gave a sample mean of $\overline{x} = 73.9$ k€.

What is the probability that the population mean μ is exactly 73.9 k€?

Zero. $\overline{x}$ is not μ — it is one **estimate** of it, computed from the seven people who happened to be in the sample.

The honest question is not *what is* μ but **how far from** μ **can this number be?**

Estimator and Estimate

Definition. An **estimator** is a function of the sample, $\hat{\theta} = g(X_1, \ldots, X_n)$, used to approximate an unknown quantity θ of the population. The number it takes on for one particular sample is an **estimate**.


INTUITION

The estimator is the **recipe**, $\overline{X} = \frac{1}{n}\sum_i X_i$; the estimate is what one batch of data cooks from it, $\overline{x} = 73.9$.

The sample is random, so the estimator is a **function of random variables**: $\hat{\theta}$ is itself a random variable, with a distribution of its own.

$\overline{x}$, s^2 and $\hat{\sigma}^2$ from descriptive statistics are all estimators — used so far without saying what kind of object they are.

An Estimator Has a Distribution



Predict first: after a thousand samples, what shape will the lower picture have — and how wide?

Three Things the Picture Shows

- → The means pile up **around** μ — the estimator is not systematically off
- → The pile is **narrower than the population** — averaging cancels noise
- → The pile is **bell-shaped**, even though the population was not

> **Definition (Sampling distribution).** The distribution of an estimator $\hat{\theta}$ over all possible samples of size n. $\hat{\theta}$ is **unbiased** if $E[\hat{\theta}] = \theta$.

Use the Widget: Change the Population

Set the population to **skewed** and then to **bimodal**, keeping $n = 5$.

Which of the three observations survive a population that is not symmetric?

Use the Widget: Change the Population

Set the population to **skewed** and then to **bimodal**, keeping $n = 5$.

Which of the three observations survive a population that is not symmetric?

All three. The shape of the sampling distribution stops depending on the shape of the population as soon as n is more than a handful — which is why one formula can cover cases we never inspect.

Standard Error

Definition (Standard error). The standard deviation of an estimator's sampling distribution. For the sample mean of independent draws:

$$\mathrm{SE}(\overline{X}) = \frac{\sigma}{\sqrt{n}}$$

INTUITION

How far off a typical **estimate** is — where σ says how far apart typical **observations** are. Both in the unit of the data.

σ and SE are different quantities. More data does nothing to σ — the world stays as variable as it was — and shrinks SE.

Precision Costs Data

Multiply the data by k, and the standard error is multiplied by $1/\sqrt{k}$:

0.71	→ **½**	→ **⅕**	→ **1⁄10**
2 × the data	4 × the data	25 × the data	100 × the data

The $\sqrt{n}$ is why the last digit of an estimate is so expensive: halving the error costs four times the data, and a tenfold improvement costs a hundredfold.

Two Theorems We Have Already Seen Happen

For i.i.d. observations with mean μ and finite variance σ^2:

Law of large numbers	$\overline{X}_n \to \mu \text{ as } n \to \infty$	*the pile keeps narrowing onto the truth*
Central limit theorem	$\overline{X}_n \approx N(\mu, \sigma^2/n) \text{ for large } n$	*and we know the shape of what is left*

The first says the error goes away; the second says how much of it is left after n observations. Only the second lets us put a number on our ignorance.

From a Number to an Interval

If $\overline{X} \approx N(\mu, \sigma^2/n)$, then for 95% of samples $\overline{X}$ lands within $1.96\,\mathrm{SE}$ of μ — and therefore μ lands within $1.96\,\mathrm{SE}$ of $\overline{X}$.

$$\overline{x} \pm 1.96 \cdot \frac{\sigma}{\sqrt{n}}$$


the estimate — our best single guess

how confident we insist on being

how noisy the estimate is

What Does "95% Confident" Mean?

Change the level or n — which number follows, and which does not?



58 of 60 intervals cover μ (97%) — nominal 95%

The Sentence Almost Everyone Gets Wrong

WHAT IT DOES NOT MEAN

"There is a 95% probability that μ lies in **this** interval."

μ is a fixed number. It is either in this interval or it is not — there is no probability left to have.

WHAT IT MEANS

"95% of the intervals produced **by this procedure** contain μ."

The probability is a property of the recipe, evaluated before the data is seen.

Where This Leaves Us

Two kinds of uncertainty meet in one formula. We can now measure one of them.

- → σ — how variable the world is. **Aleatoric.** Unchanged by data.
- → $\mathrm{SE} = \sigma/\sqrt{n}$ — how badly we know our own estimate. **Epistemic.** Shrinks with data.

"More data helps" was a slogan.
It is now a rate: $1/\sqrt{n}$.

Mini-Exercise

A shop timed $n = 100$ customers at a counter: $\overline{x} = 180$ s, and past records give $\sigma = 60$ s. What is the standard error — and how many customers would halve it?

Mini-Exercise

A shop timed $n = 100$ customers at a counter: $\overline{x} = 180$ s, and past records give $\sigma = 60$ s. What is the standard error — and how many customers would halve it?

$\mathrm{SE} = 60/\sqrt{100} = 6$ s, so the 95% interval is about 180 ± 12 s. Halving the SE needs twice the $\sqrt{n}$ — **four times the data**, $n = 400$. $\sigma = 60$ s does not change: customers stay as variable as they were; only our knowledge of their average improves.

Takeaway

A statistic is a **random variable**. Its distribution over repeated samples is the sampling distribution; its spread is the **standard error**, $\sigma/\sqrt{n}$. A confidence interval reports that spread — and its percentage describes the **procedure**, not the one interval in front of you.

An Unknown Number, Ten Observations

A new feature went live. Of the first **10 users**, **3 came back**.

What is p, the probability that a user comes back — and how sure are we?

An Unknown Number, Ten Observations

A new feature went live. Of the first **10 users**, **3 came back**.

What is p, the probability that a user comes back — and how sure are we?

WHAT THE LAST UNIT CAN SAY

$\hat{p} = 0.3$, standard error ≈ 0.14, a 95% interval of roughly $[0.02, 0.58]$ — about the *procedure*, not about p.

WHAT WE WOULD LIKE TO SAY

"p is probably between 0.1 and 0.6, most likely around 0.3" — a statement **about** p.

Conditioning on the Data — for a Parameter

Treat the unknown p as a random variable θ and apply **Bayes' theorem**, with θ in place of the hypothesis and the data D as the evidence:

$$p(\theta \mid D) = \frac{p(D \mid \theta) \cdot p(\theta)}{p(D)}$$

posterior — belief about θ after the data

likelihood — how probable the data is for each θ

evidence — normalises, does not depend on θ

prior — belief about θ before

For i.i.d. yes/no data with h yeses in n trials: $p(D \mid \theta) = \theta^h(1-\theta)^{n-h}$ — read as a function of θ, with the data fixed.

A Prior That Behaves Like Counts


Beta prior. $p(\theta) \propto \theta^{a-1}(1-\theta)^{b-1}$ on $[0,1]$, written $\mathrm{Beta}(a,b)$. $\mathrm{Beta}(1,1)$ is flat: every p equally plausible.

Multiply by the likelihood $\theta^h(1-\theta)^{n-h}$:

$$p(\theta \mid D) \propto \theta^{a+h-1}(1-\theta)^{b+n-h-1} \quad \Longrightarrow \quad \theta \mid D \sim \mathrm{Beta}(a+h,\ \ b+n-h)$$

The update is **adding counts**: a and b act like yeses and noes seen before the data. A prior whose posterior stays in the same family is called **conjugate**.

Watching the Posterior Form



Predict before each click: which way does the curve move after a ✗ — and does it get narrower every time?

What the Posterior Gives You


After 3 of 10, with a flat prior: $\theta \mid D \sim \mathrm{Beta}(4, 8)$.

question	answer	from
most plausible p	$0.30 = h/n$	the peak (mode)
best guess on average	0.33	the posterior mean $\frac{a+h}{a+b+n}$
where is p, 95%?	$[0.11,\ 0.61]$	a **credible interval**
after 1,000 users, 300 back	$[0.27,\ 0.33]$	same model, more data

> The peak under a flat prior is exactly h/n. **Lecture 03** keeps only this peak and calls it maximum likelihood — the first shortcut of the course.

How Much Does the Prior Matter?

Two analysts: one with a flat prior $\mathrm{Beta}(1, 1)$, one convinced that p is near a half, $\mathrm{Beta}(10, 10)$.



With little data the prior matters; with enough data the analysts **agree**. A prior is a stated assumption — it can be argued with, and it washes out.

Mini-Exercise

Prior $\mathrm{Beta}(2, 2)$. Of 10 users, **8** come back.

Is the posterior mean above or below 0.8 — and why?

Mini-Exercise

Prior $\mathrm{Beta}(2, 2)$. Of 10 users, **8** come back.

Is the posterior mean above or below 0.8 — and why?

Below. The posterior is $\mathrm{Beta}(2 + 8,\ 2 + 2) = \mathrm{Beta}(10, 4)$, mean $10/14 \approx 0.71$. The prior counts two yeses and two noes, so it pulls towards 0.5 — and the pull fades as data grows.

Two Ways to Answer "Where Is p?"

	random	the 95% interval	needs
Frequentist	the data; p is fixed	a **procedure** that catches p in 95% of samples — about this one interval, no probability statement	nothing but the sampling model
Bayesian	p; the data is fixed	"p lies in this interval with probability 0.95" — a true sentence about p	a **prior**: what we believed about p before the data

> The axioms are the same; the question is not. The rest of the course mostly takes the second road — and Lecture 03 starts with its cheapest version: keep only the peak.

Takeaway

An unknown parameter is just another unknown: give it a **prior**, condition on the data, and the **posterior** says what the data justifies — as a peak, a mean and an interval. For yes/no data with a Beta prior the update is **adding counts**, and the prior's influence fades as data grows.

Image Credits & References

Sheldon, Ross and others (2002). A first course in probability. *Pearson Education India*.