

Algebra Definitions

Peter Kagey

May 2019

1 Groups

1.1 Notation and definitions

1.1.1 Basic definitions

Definition 1.1.1.1 (Normal subgroup). Let G be a group and K be a subgroup of G . If $gkg^{-1} \in K$ for all $k \in K$ and $g \in G$, then K is called a normal subgroup of G and is denoted $K \trianglelefteq G$.

Definition 1.1.1.2 (Simple group). A group G is called a simple group is a group whose only normal subgroups are $\{e\}$ and G .

Definition 1.1.1.3 (Semidirect product). Let $K \trianglelefteq G$ and $Q \leq G$. A group G is a semidirect product of K by Q (denoted $G = K \rtimes Q$) if there exists $Q_1 \cong Q$ such that Q_1 is a complement of K in G , that is $K \cap Q_1 = 1$ and $KQ_1 = G$.

1.1.2 Galois Theory

Definition 1.1.2.1 (Normal series). A normal series of a group G is a sequence of subgroups

$$G = G_0 \geq G_1 \geq \dots \geq G_n = 1$$

in which $G_{i+1} \trianglelefteq G_i$ for all i .

Definition 1.1.2.2 (Factor groups). The factor groups of a normal series are the groups G_i/G_{i+1} for $i = 0, 1, \dots, n-1$.

Definition 1.1.2.3 (Length). The length of a normal series is the number of nontrivial factor groups.

Definition 1.1.2.4 (Solvable group). A finite group is solvable if it has a normal series whose factor groups are cyclic of prime order.

1.1.3 Centralizer/Normalizer

Definition 1.1.3.1 (Center). The center of a group G , denoted by $Z(G)$, is the set of all $a \in G$ that commute with every element of G .

Definition 1.1.3.2 (Centralizer). The centralizer of a subset S of a group G is defined to be

$$C_G(S) = \{g \in G \mid gs = sg \text{ for all } s \in S\}.$$

Definition 1.1.3.3 (Normalizer). The centralizer of a subset S in the group G is defined to be

$$N_G(S) = \{g \in G \mid gS = Sg\}.$$

Definition 1.1.3.4 (Commutator). If $a, b \in G$, the commutator of a and b , denoted $[a, b]$, is

$$[a, b] = aba^{-1}b^{-1},$$

and the commutator subgroup of G , denoted G' , is the subgroup of G generated by all of the commutators.

Definition 1.1.3.5 (Class equation). Partition G into its conjugacy classes, with x_i the representative of the i th conjugacy class. The class equation of the finite group G is

$$|G| = |Z(G)| + \sum_i [G : C_G(x_i)].$$

1.1.4 Group Actions

Definition 1.1.4.1 (Group action). Let G be a group and X be a set. Then a group action on X is a function $\varphi: G \times X \rightarrow X$ denoted $\varphi(g, x) = g \cdot x$ and satisfying

- (i) Identity: group action by the identity is trivial for all $x \in X$: $1 \cdot x = x$.
- (ii) Compatibility: $(gh) \cdot x = g \cdot (h \cdot x)$.

And X is called a G -set.

Definition 1.1.4.2 (Orbit). The orbit of an element $x \in X$ is denoted by

$$G \cdot x = \{g \cdot x \mid g \in G\}$$

Definition 1.1.4.3 (Stabilizer subgroup). The stabilizer subgroup of G with respect to $x \in X$ is denoted

$$G_x = \{g \in G \mid g \cdot x = x\}$$

Definition 1.1.4.4 (Transitive). A group action is called transitive if for each $x, y \in X$ there exists some $g \in G$ such that $g \cdot x = y$.

1.2 Theorems

Theorem 1.2.1 (First isomorphism theorem). If $\varphi: G \rightarrow H$ is a group homomorphism then $\ker(\varphi) \trianglelefteq G$ and $G/\ker(\varphi) \cong \varphi(G)$.

Theorem 1.2.2 (Second isomorphism theorem). Let G be a group with $S \leq G$ and $N \trianglelefteq G$. Then

1. $SN \leq G$
2. $S \cap N \trianglelefteq S$, and
3. $(SN)/N \cong S/(S \cap N)$.

Strictly speaking, N does not have to be a normal subgroup as long as S is a subgroup of the normalizer of N , $S \leq N_G(N)$.

Theorem 1.2.3 (Third isomorphism theorem). Let G be a group with normal subgroup $N \trianglelefteq G$. Then

1. If $K \leq G$ (resp. $K \trianglelefteq G$) such that $N \subseteq K \subseteq G$, then $K/N \leq G/N$ (resp. $K/N \trianglelefteq G/N$).
2. Every subgroup (resp. normal subgroup) of G/N is of the form K/N , for some subgroup (resp. normal subgroup) $K \subset G$ such that $N \subseteq K \subseteq G$.
3. If $K \trianglelefteq G$ such that $N \subseteq K \subseteq G$, then $(G/N)/(K/N) \cong G/K$.

Theorem 1.2.4 (Simplicity of the A_n). A_n is simple for all $n \geq 5$.

Theorem 1.2.5 (Sylow's theorem).

- (i) If P is a Sylow p -subgroup of a finite group G , then all Sylow p -subgroups of G are conjugate to P .
- (ii) If there are r Sylow p -subgroups, then r divides $|G|$ and $r \equiv 1 \pmod{p}$.

Theorem 1.2.6 (Fundamental Theorem of Abelian Groups). If G and H are finite abelian groups, then $G \cong H$ if and only if, for all primes p , they have the same elementary divisors.