A Constructive Introduction to Quantum Field Theory

Lecture 2: The Gaussian free field

M. Gubinelli
Mathematical Institute, University of Oxford

SISSA, October 2026

Abstract

We study the first example of a Euclidean field theory: the Gaussian free field (GFF) on the torus $\mathbb{T}^d$ and on $\mathbb{R}^d$. We construct it, prove reflection positivity and recover the free relativistic field, discuss the Euclidean Källén–Lehmann representation, and define its local (Wick) operators. We prove that the Dyson–Schwinger equation characterises the GFF among probability measures on distributions, and that the GFF is the stationary one-time (resp. one-point) marginal of the parabolic (resp. elliptic) stochastic quantisation equation. Exercises are collected at the end. These notes follow the slides of Lecture 2.

Contents

Introduction **1**

3 The Gaussian free field **2**
3.1 Definition and construction 2
3.2 Reflection positivity and the free field 3
3.3 Local operators: Wick products 7
3.4 Gaussian integration by parts and the Dyson–Schwinger equation 8
3.5 The GFF as stationary marginal of parabolic stochastic quantisation 9
3.6 The GFF as one-point marginal of elliptic stochastic quantisation 11
3.7 Outlook 12

References and further reading **14**

Introduction

In Lecture 1 we saw that, after the passage to imaginary time, a relativistic quantum field is encoded in a probability measure on distributions satisfying the Osterwalder–Schrader axioms, formally $\mu(\mathrm{d}\varphi) \propto \exp(-\int[\frac{1}{2}|\nabla\varphi|^2 + \frac{1}{2}m^2\varphi^2 + V(\varphi)])\mathcal{D}\varphi$ (Eq. (2.13)). In this lecture we treat its Gaussian part rigorously. Conventions are those of Lecture 1. Sections, equations and exercises keep the numbering of the notes, so that §1–2 refer to the notes of Lecture 1.

3 The Gaussian free field

3.1 Definition and construction

Throughout, $\mathcal{M}$ is either the torus $\mathbb{T}_L^d = (\mathbb{R}/L\mathbb{Z})^d$ or $\mathbb{R}^d$, $d \geq 1$. We write $\mathcal{S}(\mathcal{M})$ for $C^\infty(\mathbb{T}_L^d)$, resp. the Schwartz space, and $\mathcal{S}'(\mathcal{M})$ for its dual. We equip $\mathcal{S}'(\mathcal{M})$ with the σ-algebra generated by the maps $\varphi \mapsto \varphi(f)$; it coincides with the Borel σ-algebra [GV64, Bog98]. Test functions are real unless stated otherwise. Let $m > 0$ and

$$A := m^2 - \Delta, \qquad C := A^{-1}. \tag{3.1}$$

C is a positive bounded operator on $L^2(\mathcal{M})$ with $\|C\| \leq m^{-2}$. Both A and C are continuous bijections of $\mathcal{S}(\mathcal{M})$, being Fourier multipliers by $(|k|^2 + m^2)^{\pm 1}$.

Definition 3.1. The *(massive) Gaussian free field* on $\mathcal{M}$ is the centred Gaussian probability measure $\mu = \mu_m$ on $\mathcal{S}'(\mathcal{M})$ with covariance C:

$$\int \varphi(f)\varphi(g)\,\mu(\mathrm{d}\varphi) = \langle f, Cg\rangle = \iint f(x)G(x,y)g(y)\,\mathrm{d}x\,\mathrm{d}y, \tag{3.2}$$

equivalently $\int \mathrm{e}^{\mathrm{i}\varphi(f)}\mu(\mathrm{d}\varphi) = \mathrm{e}^{-\frac{1}{2}\langle f, Cf\rangle}$ for all $f \in \mathcal{S}(\mathcal{M})$. Here G is the Green function of A. On $\mathbb{R}^d$, $G(x,y) = G_m(x-y)$ with $G_m(x) = \int \frac{\mathrm{e}^{\mathrm{i}k\cdot x}}{|k|^2+m^2}\frac{\mathrm{d}k}{(2\pi)^d}$. On $\mathbb{T}_L^d$, $G(x,y) = \sum_{n\in\mathbb{Z}^d} G_m(x-y+nL)$ (method of images).

Formally $\mu_m(\mathrm{d}\varphi)\text{“}= \text{”}Z^{-1}\mathrm{e}^{-\frac{1}{2}\langle\varphi, A\varphi\rangle}\mathcal{D}\varphi$, the Gaussian part of (2.13). The Green function satisfies (Exercise 3.1): $G_m(x) \sim \frac{1}{2\pi}\log\frac{1}{|x|}$ as $x \to 0$ if $d = 2$; $G_m(x) \sim \frac{\Gamma(d/2-1)}{4\pi^{d/2}}|x|^{2-d}$ if $d \geq 3$; $G_m(x) = \mathrm{e}^{-m|x|}/(2m)$ if $d = 1$; and G_m decays like $|x|^{-(d-1)/2}\mathrm{e}^{-m|x|}$ at infinity. Uniqueness of μ_m is clear, since a probability measure on $\mathcal{S}'(\mathcal{M})$ is determined by its characteristic functional.

Theorem 3.2 (Existence and regularity). *(a) Let $\mathcal{M} = \mathbb{T}_L^d$, let $(e_n)_{n\geq 0}$ be a real orthonormal basis of $L^2(\mathbb{T}_L^d)$ of eigenfunctions of $-\Delta$, $-\Delta e_n = \kappa_n e_n$, $\lambda_n := m^2 + \kappa_n$, and let (g_n) be i.i.d. $N(0,1)$. For every $s > \frac{d}{2} - 1$ the series*

$$\varphi = \sum_{n\geq 0} \lambda_n^{-1/2} g_n\, e_n \tag{3.3}$$

converges a.s. and in $L^2(\mathbb{P}; H^{-s}(\mathbb{T}_L^d))$, and its law is μ_m. In particular $\mu_m(H^{-s}) = 1$ for $s > \frac{d}{2} - 1$.
(b) On $\mathbb{R}^d$, μ_m exists by the Bochner–Minlos theorem.

Proof. (a) With $\|\varphi\|_{H^{-s}}^2 = \sum_n (1+\kappa_n)^{-s}\langle e_n, \varphi\rangle^2$,

$$\mathbb{E}\|\varphi\|_{H^{-s}}^2 = \sum_n \frac{1}{(1+\kappa_n)^s\lambda_n} \asymp \sum_{k\in\mathbb{Z}^d} (1+|k|^2)^{-s-1} < \infty \iff 2s+2 > d.$$

The partial sums form an L^2-bounded martingale with values in the Hilbert space H^{-s}, so they converge a.s. and in L^2. For $f \in \mathcal{S}(\mathcal{M})$, $\varphi(f) = \sum_n \lambda_n^{-1/2} g_n \langle e_n, f\rangle$ is centred Gaussian with variance $\sum_n \lambda_n^{-1}\langle e_n, f\rangle^2 = \langle f, Cf\rangle$.

(b) Bochner–Minlos [GV64, GJ87]: if $\Psi : \mathcal{S}(\mathbb{R}^d) \to \mathbb{C}$ is continuous, positive definite and $\Psi(0) = 1$, there is a unique probability measure on $\mathcal{S}'(\mathbb{R}^d)$ with characteristic functional Ψ. For $\Psi(f) = \mathrm{e}^{-\frac{1}{2}\langle f, Cf\rangle}$, continuity follows from $\langle f, Cf\rangle \leq m^{-2}\|f\|_{L^2}^2$. For positive definiteness, write $\Psi(f_j - f_k) = a_j a_k \mathrm{e}^{Q_{jk}}$ with $a_j = \mathrm{e}^{-\frac{1}{2}Q_{jj}}$ and $Q_{jk} = \langle f_j, Cf_k\rangle$ positive semidefinite. Then $(\mathrm{e}^{Q_{jk}})$ is positive semidefinite by the Schur product theorem. □




Figure 1: Left: a sample of the massive GFF on $\mathbb{T}^2$ ($m = 4$, Fourier cutoff 512^2, zero mode removed). It is not a function: refining the cutoff adds fluctuations at all scales. Right: the Green function G_m in $d = 1, 2, 3$: bounded, logarithmic and $|x|^{-1}$ singularities at the origin, and exponential decay at infinity.

Remark 3.3. The Cameron–Martin space of μ_m is $H^1(\mathcal{M})$ with norm $\langle h, Ah\rangle^{1/2}$: the fields of finite action. It has measure zero, and typical samples are much rougher (Exercise 3.3). In $d = 1$ the GFF is the stationary Ornstein–Uhlenbeck process. In $d = 2$ it lies in $H^{-\kappa}$ for every $\kappa > 0$ but is not a function. In $d = 3$ it lies in $H^{-1/2-\kappa}$. On $\mathbb{R}^d$ the samples live in weighted Sobolev spaces.

3.2 Reflection positivity and the free field

Let $\mathcal{M} = \mathbb{R} \times N$ with $N = \mathbb{R}^{d-1}$ or $\mathbb{T}_L^{d-1}$, points $(\tau, \mathbf{x})$, $\Theta(\tau, \mathbf{x}) = (-\tau, \mathbf{x})$. Then $A = -\partial_\tau^2 + B$ with $B := m^2 - \Delta_N$ on $L^2(N)$, and we set $\omega := B^{1/2} \geq m$.

Proposition 3.4. *(a) As an operator-valued kernel in the time variables,* $C(\tau, \tau') = \mathrm{e}^{-|\tau-\tau'|\omega}(2\omega)^{-1}$. *(b) For* $f \in \mathcal{S}(\mathcal{M})$ *with* $\operatorname{supp} f \subset \{\tau > 0\}$,

$$\langle \Theta f, Cf\rangle = \Big\| (2\omega)^{-1/2} \int_0^\infty \mathrm{e}^{-\tau\omega} f(\tau, \cdot)\, \mathrm{d}\tau \Big\|^2_{L^2(N)} \geq 0. \tag{3.4}$$

(c) μ_m *on* $\mathbb{R}^d$ *satisfies (OS0)–(OS4).*

Proof. (a) follows from the partial Fourier transform in τ, the spectral theorem for B, and the scalar identity $\int_{\mathbb{R}} \mathrm{e}^{\mathrm{i}kt}(k^2 + w^2)^{-1} \frac{\mathrm{d}k}{2\pi} = \mathrm{e}^{-w|t|}/(2w)$, $w > 0$. (b) Since Θf is supported in $\{\tau < 0\}$,

$$\begin{aligned} \langle \Theta f, Cf\rangle &= \iint_{\tau<0<\tau'} \langle f(-\tau), \mathrm{e}^{-(\tau'-\tau)\omega}(2\omega)^{-1} f(\tau')\rangle \, \mathrm{d}\tau \, \mathrm{d}\tau' \\ &= \int_0^\infty \int_0^\infty \langle \mathrm{e}^{-s\omega} f(s), (2\omega)^{-1} \mathrm{e}^{-\tau'\omega} f(\tau')\rangle \, \mathrm{d}s \, \mathrm{d}\tau', \end{aligned}$$

which is (3.4). (c) (OS0)–(OS2) and (OS4) follow from the explicit form of $\hat{\mu}(f)$ and the decay of G_m. For (OS3), C commutes with Θ, so $\hat{\mu}(f_i - \Theta f_j) = a_i a_j \mathrm{e}^{\langle \Theta f_i, Cf_j\rangle}$ with $a_i = \mathrm{e}^{-\frac{1}{2}\langle f_i, Cf_i\rangle}$. The matrix $(\langle \Theta f_i, Cf_j\rangle)_{ij}$ is positive semidefinite by (b), hence so is its entrywise exponential (Schur). □




Figure 2: Reflection positivity (Proposition 3.4). For f supported in $\{\tau > 0\}$ the covariance between f and its mirror image factorises through the time-zero plane: $\langle \Theta f, Cf\rangle = \|(2\omega)^{-1/2}\int_0^\infty \mathrm{e}^{-\tau\omega} f(\tau)\,\mathrm{d}\tau\|^2 \geq 0$.

By the OS reconstruction theorem the GFF yields a Wightman theory, the free field of mass m of Section 1.3. Concretely, the reconstructed Hilbert space is the Fock space over $L^2(N)$ with $H = \mathrm{d}\Gamma(\omega)$ (Nelson's free Markov field [Nel73a, Nel73b]). The time-zero restriction of the GFF is Gaussian with covariance $(2\omega)^{-1}$, the ground state measure of the free Hamiltonian in the Schrödinger representation (Exercise 3.13). The GFF is also the continuum limit of the lattice model (2.12) with $V = 0$ (for $d = 2$; in general d see Exercise 2.3 and Exercise 3.14).

Remark 3.5 (Euclidean Källén–Lehmann representation). The computation in Proposition 3.4(a) says that $\int \mathrm{d}\Omega_M(\mathbf{p})\,\mathrm{e}^{-\omega_M(\mathbf{p})|\tau| + \mathrm{i}\mathbf{p}\cdot\mathbf{x}} = G_M(\tau,\mathbf{x})$: the Wick rotation of the free two-point function $W_2^{(M)}$ is the Green function G_M of $M^2 - \Delta$. Rotating (1.4) term by term, the Schwinger two-point function of any theory satisfying the Wightman (or OS) axioms is

$$S_2(x) = v^2 + \int_{[0,\infty)} \rho(\mathrm{d}M^2)\, G_M(x), \qquad x \neq 0, \qquad \text{i.e.} \qquad \hat{S}_2^T(k) = \int_{[0,\infty)} \frac{\rho(\mathrm{d}M^2)}{|k|^2 + M^2}, \tag{3.5}$$

a Stieltjes transform in $|k|^2$. (The Fourier form holds up to contact terms, i.e. a polynomial in $|k|^2$ invisible at $x \neq 0$, and provided $\int \rho(\mathrm{d}M^2)(1+M^2)^{-1} < \infty$; otherwise subtractions are needed.) The GFF is the case $\rho = \delta_{m^2}$. Three consequences follow.

(i) *Short distances.* $G_M(x)$ is decreasing in M, and for $d \geq 2$ we have $G_M(x)/G_m(x) \to 1$ as $x \to 0$, for every M. Hence, for every K, $S_2^T(x) \geq \rho([0,K^2])\, G_K(x)$, and, if $\rho \neq 0$,

$$\liminf_{x\to 0} \frac{S_2^T(x)}{G_m(x)} \geq \rho([0,\infty)) \in (0,\infty].$$

The two-point function of any nontrivial theory is at least as singular as the free one. In particular a scalar field cannot have scaling dimension below the free value $\frac{d-2}{2}$ (the *unitarity bound*, Exercise 3.2).

(ii) *Long distances.* If $\operatorname{supp}\rho \subset [m_0^2,\infty)$, then $S_2^T(x)$ decays at least like $\mathrm{e}^{-m_0|x|}$, up to subexponential factors, and exactly at this rate if $m_0^2 = \inf\operatorname{supp}\rho$. Conversely, exponential decay at rate m_0 forces $\operatorname{supp}\rho \subset [m_0^2,\infty)$. So a gap in the spectral measure of φ is the same as exponential decay of S_2^T; a mass gap of H corresponds to exponential clustering of all truncated correlations (compare Exercise 2.6).

(iii) *Generalised free fields.* For any positive measure ρ with $\int \rho(\mathrm{d}M^2)(1+M^2)^{-1} < \infty$ (and, so that $\hat{C}_\rho$ is locally integrable at $k = 0$, $\int_{[0,1]} \rho(\mathrm{d}M^2)\log(1/M^2) < \infty$ if $d = 2$ and $\int_{[0,1]} \rho(\mathrm{d}M^2)M^{-1} < \infty$ if $d = 1$), the Gaussian measure on $\mathcal{S}'(\mathbb{R}^d)$ with covariance $C_\rho = \int \rho(\mathrm{d}M^2)(M^2 - \Delta)^{-1}$ satisfies the OS axioms: reflection positivity follows from Proposition 3.4

for each mass $M > 0$ (and, for an atom at $M = 0$ when $d \geq 3$, by letting $M \downarrow 0$) and $\rho \geq 0$. It reconstructs the generalised free field of Exercise 1.6. Its Dyson–Schwinger equation $\mathbb{E}[D_h F] = \mathbb{E}[F\varphi(C_\rho^{-1}h)]$ still characterises it: under mild conditions on ρ (e.g. ρ finite with a mass gap), C_ρ is a bijection of $\mathcal{S}(\mathbb{R}^d)$, which is all the proof of Theorem 3.12 uses. However C_ρ^{-1} is *nonlocal* unless ρ is a single atom, as the following example shows.

Example: two masses. Let $0 < m_1 < m_2$, $c_1, c_2 > 0$, and

$$\rho = c_1^2\,\delta_{m_1^2} + c_2^2\,\delta_{m_2^2}, \qquad C_\rho = c_1^2(m_1^2 - \Delta)^{-1} + c_2^2(m_2^2 - \Delta)^{-1}.$$

Euclidean theory. The Gaussian measure μ_ρ with covariance C_ρ is the law of $c_1\varphi_1 + c_2\varphi_2$, where φ_1, φ_2 are independent GFFs on $\mathbb{R}^d$ of masses m_1, m_2. It satisfies (OS0)–(OS4); reflection positivity follows from Proposition 3.4 applied to each term.

The Minkowski theory. Since a Wightman theory is determined up to unitary equivalence by its Wightman functions, which are in turn determined by the Schwinger functions, μ_ρ reconstructs to exactly the following theory. Let $\mathfrak{h}_m = L^2(\mathbb{R}^s, \mathrm{d}\Omega_m)$ be the one-particle space of mass m and spin 0, carrying the irreducible positive-energy representation u_m of $\mathcal{P}_+^\uparrow$. Then

$$\mathcal{H} = \Gamma(\mathfrak{h}_{m_1} \oplus \mathfrak{h}_{m_2}) \simeq \Gamma(\mathfrak{h}_{m_1}) \otimes \Gamma(\mathfrak{h}_{m_2}), \qquad U = \Gamma(u_{m_1} \oplus u_{m_2}), \qquad \Omega = \Omega_1 \otimes \Omega_2,$$

$$\varphi(x) = c_1\,\varphi_{m_1}(x) \otimes \mathbb{1} + c_2\,\mathbb{1} \otimes \varphi_{m_2}(x),$$

with φ_m the free field (1.2) of mass m. The Wightman functions are given by Wick's theorem with $W_2 = c_1^2 W_2^{(m_1)} + c_2^2 W_2^{(m_2)}$, and all truncated functions of order ≥ 3 vanish. Physically, these are *two species of free scalar particles*, of masses m_1 and m_2, both created from the vacuum by the same field: $\varphi(f)\Omega = c_1 a_1^*(\tilde{f}|_{H_{m_1}^+})\Omega + c_2 a_2^*(\tilde{f}|_{H_{m_2}^+})\Omega$. They do not scatter. The mass spectrum consists of the two shells m_1, m_2 and of the multi-particle continuum from $2m_1$. If $m_2 > 2m_1$, the second shell lies inside the two-particle continuum, but the heavy particle is stable because there is no interaction. This is the generalised free field of Exercise 1.6.

What is lost. (1) There is no field equation of second order: $(\Box + m_1^2)(\Box + m_2^2)\varphi = 0$, but φ solves no Klein–Gordon equation. The time-zero fields $\varphi(0,\cdot)$, $\pi(0,\cdot)$ satisfy $[\varphi(0,\mathbf{x}), \pi(0,\mathbf{y})] = \mathrm{i}(c_1^2 + c_2^2)\delta(\mathbf{x} - \mathbf{y})$, which is canonical iff $c_1^2 + c_2^2 = 1$ (the sum rule of Exercise 1.5). Even when they are canonical, they are not a complete set of Cauchy data: $\varphi(t,\cdot)$ depends on $c_1\varphi_{m_1}(0,\cdot)$ and $c_2\varphi_{m_2}(0,\cdot)$ separately, not only on their sum. (2) In Euclidean terms, the DS equation $\mathbb{E}[D_h F] = \mathbb{E}[F\varphi(C_\rho^{-1}h)]$ involves the nonlocal operator

$$C_\rho^{-1} = PR^{-1}, \qquad P := (m_1^2 - \Delta)(m_2^2 - \Delta), \qquad R := c_1^2(m_2^2 - \Delta) + c_2^2(m_1^2 - \Delta).$$

Replacing h by Rh gives the equivalent local identity $\mathbb{E}[D_{Rh}F] = \mathbb{E}[F\varphi(Ph)]$: a fourth-order Euclidean field equation whose contact term is itself differentiated, the Euclidean form of $(\Box + m_1^2)(\Box + m_2^2)\varphi = 0$. Correspondingly, φ has no Markov property: for stationary Gaussian fields with a rational spectral density, such as $\hat{C}_\rho = \hat{R}/\hat{P}$, the Markov property holds iff the inverse covariance is a differential operator, i.e. iff $1/\hat{C}$ is a polynomial [Roz82], which is not the case here. The pair (φ_1, φ_2) is Markov, but its sum is not: this is the Euclidean counterpart of (1).

Positivity versus locality. Conversely, the local fourth-order action $\frac{1}{2}\langle\varphi, P\varphi\rangle$ defines a Markov Gaussian field with covariance

$$P^{-1} = \frac{(m_1^2 - \Delta)^{-1} - (m_2^2 - \Delta)^{-1}}{m_2^2 - m_1^2}, \qquad \text{i.e.} \qquad \rho = \frac{\delta_{m_1^2} - \delta_{m_2^2}}{m_2^2 - m_1^2},$$

with a *negative* weight at m_2^2. By (3.5) and the uniqueness of the Stieltjes transform it is not reflection positive (Exercise 3.16). It reconstructs at best a theory with an indefinite metric, in which the particle of mass m_2 is a ghost of negative norm [PU50, LW69]. More generally, call the Euclidean field equation *local* if C^{-1} is a differential operator, i.e. $\hat{C} = 1/p(|k|^2)$ with p a polynomial; local field equations give Markov fields, and for rational spectral densities the converse holds [Roz82]. The Fourier transform of a covariance satisfying the OS axioms is $q(|k|^2) + \int \rho(\mathrm{d}M^2)(|k|^2 + M^2)^{-1}$, with $\rho \geq 0$ and q a polynomial coming from contact terms invisible to ρ. If $\deg p \geq 1$, then $q = 0$, since $1/p(s) \to 0$ and the Stieltjes term tends to 0 as $s \to \infty$. Moreover $s \int \rho(\mathrm{d}M^2)(s + M^2)^{-1}$ increases to $\rho([0, \infty))$ as $s \to \infty$, while $s/p(s) \to 0$ if $\deg p \geq 2$; hence $\rho = 0$, a contradiction. If $\deg p = 0$, $\hat{C}$ is constant. So a centred Gaussian measure satisfying the OS axioms has a local Euclidean field equation iff its covariance is $c(M^2 - \Delta)^{-1}$ ($M \geq 0$, with $M > 0$ if $d \leq 2$), or a multiple of the identity (white noise, whose reconstructed Hilbert space is $\mathbb{C}\Omega$). Note that it is the covariance, not ρ, that matters: $a\mathbb{1} + c(m^2 - \Delta)^{-1}$ with $a > 0$ is reflection positive and has $\rho = c\delta_{m^2}$, but its field equation is nonlocal. *Positivity and locality of the Euclidean field equation together single out the free field* (up to the trivial white noise); locality is also behind Nelson's Markov property of the GFF [Nel73a].

Is the nonlocal theory physical? In Minkowski space the two-mass theory is a perfectly good one. It satisfies all the Wightman axioms, including locality (W3): $[\varphi(x), \varphi(y)]$ is a sum of free-field commutators and vanishes at spacelike separation. The energy is positive and the vacuum unique. It is trivial (no interaction) and redundant, but not inconsistent. "Nonlocal" refers only to the Euclidean field equation for φ alone, equivalently (for rational spectral densities, as here) to the absence of the Markov property. On the Minkowski side this means that φ alone is not a complete set of local degrees of freedom. Its time-zero data do not determine its evolution, and there is no Lagrangian or Hamiltonian written in terms of φ only. The cure is to enlarge the field content: the pair (φ_1, φ_2) has a local action and is Markov, and φ is a linear combination of its components.

Generalised free fields become genuinely unphysical when ρ has a continuous part, i.e. (heuristically) infinitely many free species with masses in a bounded interval. With finitely many masses, φ solves a hyperbolic equation of finite order, and the fields in an arbitrarily thin time slice generate all the observables: the *time-slice axiom*, or primitive causality [Haa96, §II.1]. With a continuum of masses there is no dynamical law of finite order, and this property is not expected to hold. More decisively, the theory violates the phase-space requirements that express "finitely many degrees of freedom per unit volume of phase space", namely the compactness criterion of Haag and Swieca [HSw65] and its sharpening, the nuclearity condition of Buchholz and Wichmann [BW86]. For a system of free fields the nuclearity condition holds iff $\sum_i n_i \mathrm{e}^{-\beta m_i} < \infty$ for all $\beta > 0$, where n_i is the multiplicity of the mass m_i [Haa96, §V.5], and this fails when infinitely many species have masses in a bounded interval. Since the nuclearity index of a bounded region plays the role of its partition function, the theory has no sensible thermodynamics. This does not affect interacting theories, whose spectral measure also has a continuous part, the multi-particle continuum. There the continuum is produced by a *local* interaction, and the Euclidean field equation remains local, e.g. $\mathbb{E}[D_h F] = \mathbb{E}[F((m^2 - \Delta)\varphi + 4\lambda{:}\varphi^3{:})(h)]$ for Φ^4_2, although the inverse two-point function is nonlocal. Thus, for a single field: positive and nonlocal means an incomplete description, which is unphysical for continuous ρ; local and not positive means ghosts; local, positive and with a nontrivial continuum in ρ requires interactions.

3.3 Local operators: Wick products

Interactions $\int V(\varphi(x))\mathrm{d}x$ and nonlinear field equations require local functions of $\varphi(x)$, but φ is not a function for $d \geq 2$. We regularise, subtract, and remove the regularisation. The subtraction is encoded in Hermite polynomials with variance $c \geq 0$,

$$\mathrm{e}^{ta-ct^2/2} = \sum_{n\geq 0} \frac{t^n}{n!} H_n(a;c), \qquad H_2 = a^2 - c, \quad H_3 = a^3 - 3ca, \quad H_4 = a^4 - 6ca^2 + 3c^2. \tag{3.6}$$

Lemma 3.6. *If (X, Y) is a centred Gaussian vector with $\mathbb{E}X^2 = a$ and $\mathbb{E}Y^2 = b$, then $\mathbb{E}[H_n(X;a)H_k(Y;b)] = \delta_{nk}\, n!\, (\mathbb{E}[XY])^n$.*

Proof. $\mathbb{E}[\mathrm{e}^{sX - s^2a/2}\mathrm{e}^{tY - t^2b/2}] = \mathrm{e}^{\frac{1}{2}\mathbb{E}(sX+tY)^2 - \frac{1}{2}s^2a - \frac{1}{2}t^2b} = \mathrm{e}^{st\mathbb{E}[XY]}$. Expand both sides in powers of s, t using (3.6) (the exchange of sum and expectation is justified by Gaussian integrability) and compare coefficients. □

Fix an even mollifier $0 \leq \eta \in C_c^\infty(B(0,1))$ with $\int \eta = 1$, $\eta_\varepsilon = \varepsilon^{-d}\eta(\cdot/\varepsilon)$ (periodised on the torus), and let $\varphi_\varepsilon = \eta_\varepsilon * \varphi$, a smooth function. Then $\mathbb{E}[\varphi_\varepsilon(x)\varphi_\delta(y)] = G_{\varepsilon,\delta}(x-y)$ with $G_{\varepsilon,\delta} = \eta_\varepsilon * \eta_\delta * G$. Set $c_\varepsilon = G_{\varepsilon,\varepsilon}(0)$ and

$$:\!\varphi_\varepsilon^n\!:(x) := H_n(\varphi_\varepsilon(x); c_\varepsilon), \qquad :\!\varphi_\varepsilon^n\!:(f) := \int f(x) :\!\varphi_\varepsilon^n\!:(x)\,\mathrm{d}x. \tag{3.7}$$

Theorem 3.7 (Wick powers)**.** *Let $n \geq 1$ with $n(d-2) < d$, i.e. every n if $d \leq 2$, $n \leq 2$ if $d = 3$, $n = 1$ if $d \geq 4$. Then for every $f \in \mathcal{S}(\mathcal{M})$, $:\!\varphi^n\!:(f) := \lim_{\varepsilon\to 0} :\!\varphi_\varepsilon^n\!:(f)$ exists in $L^2(\mu_m)$, is independent of η, and*

$$\mathbb{E}\big[:\!\varphi^n\!:(f) :\!\varphi^k\!:(g)\big] = \delta_{nk}\, n! \iint f(x) G(x-y)^n g(y)\,\mathrm{d}x\,\mathrm{d}y. \tag{3.8}$$

If instead $n(d-2) \geq d$ and $0 \leq f \not\equiv 0$, then $\mathbb{E}[:\!\varphi_\varepsilon^n\!:(f)^2] \to \infty$.

Proof. Step 1: a uniform bound. Let $K(z) = 1 + \log_+(1/|z|)$ if $d = 2$, $K(z) = 1 + |z|^{2-d}$ if $d \geq 3$, $K \equiv 1$ if $d = 1$. Then $0 < G \leq C_0 K$, and we claim $G_{\varepsilon,\delta} \leq CK$ uniformly in $\varepsilon, \delta \in (0,1]$. Let $r = \max(\varepsilon, \delta)$. Then $\operatorname{supp}(\eta_\varepsilon * \eta_\delta) \subset B(0, 2r)$, $\int \eta_\varepsilon * \eta_\delta = 1$ and $\|\eta_\varepsilon * \eta_\delta\|_\infty \leq r^{-d}\|\eta\|_\infty$. If $|z| \geq 4r$, then $G_{\varepsilon,\delta}(z) \leq \sup_{|w|\geq |z|/2} G(w) \leq C_0 K(z/2) \leq CK(z)$. If $|z| < 4r$, then

$$G_{\varepsilon,\delta}(z) \leq \|\eta\|_\infty\, r^{-d} \int_{B(0,6r)} G \leq C \fint_{B(0,6r)} K \leq CK(r) \leq C'K(z).$$

Step 2: convergence. By Lemma 3.6 and Fubini,

$$\mathbb{E}\big[:\!\varphi_\varepsilon^n\!:(f) :\!\varphi_\delta^k\!:(g)\big] = \delta_{nk}\, n! \iint f(x) g(y)\, G_{\varepsilon,\delta}(x-y)^n\,\mathrm{d}x\,\mathrm{d}y.$$

As $\varepsilon, \delta \to 0$, $G_{\varepsilon,\delta}(z) \to G(z)$ for $z \neq 0$. Moreover $K^n \in L^1_{\mathrm{loc}}$ iff $n(d-2) < d$, and K is bounded on $\{|z| \geq 1\}$. Since $f, g \in L^1 \cap L^\infty$, dominated convergence gives $\mathbb{E}[(:\!\varphi_\varepsilon^n\!:(f) - :\!\varphi_\delta^n\!:(f))^2] = n! \iint ff\,(G_{\varepsilon,\varepsilon}^n - 2G_{\varepsilon,\delta}^n + G_{\delta,\delta}^n) \to 0$: the sequence is Cauchy in L^2. The limit covariance is (3.8). Running the same computation with two different mollifiers shows independence of η. *Step 3: divergence.* If $n(d-2) \geq d$, then $G^n \geq c|z|^{-n(d-2)}$ is not locally integrable, and Fatou's lemma gives $\liminf \mathbb{E}[:\!\varphi_\varepsilon^n\!:(f)^2] \geq n! \iint ffG^n = \infty$. □

Remark 3.8. (i) The subtraction of $c_\varepsilon \to \infty$ is the simplest instance of renormalisation: $\varphi_\varepsilon(x)^n$ itself has no limit when $d \geq 2$. (ii) The variables $:\varphi^n:(f)$ belong to the n-th Wiener chaos, and $L^2(\mu_m) \simeq \bigoplus_n \mathcal{H}_n$ is the symmetric Fock space over $H^{-1}(\mathcal{M})$ [Jan97]. By Nelson's hypercontractivity, $\|X\|_{L^p} \leq (p-1)^{n/2}\|X\|_{L^2}$ for $X \in \mathcal{H}_n$, so the convergence holds in every L^p. A Kolmogorov-type argument gives convergence in $C^{-\kappa}$ for $d = 2$. (iii) In $d = 2$ on $\mathbb{T}^2$, the interaction $\int :\varphi^4: \mathrm{d}x$ lies in every $L^p(\mu_m)$, and Nelson's estimate gives $\mathrm{e}^{-\lambda \int :\varphi^4:} \in L^1(\mu_m)$ [Nel66, Sim74, GJ87]. This defines the Φ^4_2 measure on the torus. (iv) Other local operators: derivatives $\partial^\alpha \varphi$, and in $d = 2$ the Wick exponentials $:\mathrm{e}^{\mathrm{i}\beta\varphi}:$ for $\beta^2 < 4\pi$ (in L^2, Exercise 3.6), relevant for the sine-Gordon model, and the real exponentials $:\mathrm{e}^{\alpha\varphi}:$ of the Høegh-Krohn model.

3.4 Gaussian integration by parts and the Dyson–Schwinger equation

Let $\mathcal{F}C_b^\infty$ be the space of cylinder functions $F(\varphi) = g(\varphi(f_1), \dots, \varphi(f_n))$ with $g \in C_b^\infty(\mathbb{R}^n)$ and $f_i \in \mathcal{S}(\mathcal{M})$. For $h \in \mathcal{S}(\mathcal{M})$ define

$$D_h F(\varphi) := \frac{\mathrm{d}}{\mathrm{d}t} F(\varphi + t h)\Big|_{t=0} = \sum_i \partial_i g(\varphi(f_1), \dots, \varphi(f_n)) \langle f_i, h\rangle = \langle DF(\varphi), h\rangle, \tag{3.9}$$

with $DF = \sum_i \partial_i g(\dots) f_i$, formally $DF(x) = \delta F / \delta\varphi(x)$.

Lemma 3.9 (Gaussian integration by parts). *If $(Y, X_1, \dots, X_n)$ is a centred Gaussian vector and $g \in C_b^\infty(\mathbb{R}^n)$, then $\mathbb{E}[Y g(X)] = \sum_i \mathbb{E}[Y X_i]\, \mathbb{E}[\partial_i g(X)]$.*

Proof. For $g = \mathrm{e}^{\mathrm{i}\langle u, \cdot\rangle}$: $\mathbb{E}[Y \mathrm{e}^{\mathrm{i}\langle u, X\rangle}] = -\mathrm{i}\frac{\mathrm{d}}{\mathrm{d}\theta}\mathbb{E}[\mathrm{e}^{\mathrm{i}\theta Y + \mathrm{i}\langle u, X\rangle}]|_{\theta=0} = \mathrm{i}\sum_i u_i \mathbb{E}[Y X_i]\mathbb{E}[\mathrm{e}^{\mathrm{i}\langle u, X\rangle}]$, which is the claim. The general case follows by Fourier inversion for $g \in \mathcal{S}(\mathbb{R}^n)$ and then by approximation. □

Proposition 3.10 (Dyson–Schwinger equation for the GFF). *For all $F \in \mathcal{F}C_b^\infty$ and $h \in \mathcal{S}(\mathcal{M})$,*

$$\mathbb{E}_{\mu_m}[D_h F] = \mathbb{E}_{\mu_m}\big[F\, \varphi(Ah)\big], \qquad \textit{formally} \quad \mathbb{E}\Big[\frac{\delta F}{\delta\varphi(x)}\Big] = \mathbb{E}\big[F\,(m^2 - \Delta)\varphi(x)\big]. \tag{3.10}$$

Proof. Under μ_m, $(\varphi(Ah), \varphi(f_1), \dots, \varphi(f_n))$ is centred Gaussian with $\mathbb{E}[\varphi(Ah)\varphi(f_i)] = \langle Ah, C f_i\rangle = \langle h, f_i\rangle$. Apply Lemma 3.9. □

Remark 3.11 (DS = Euclidean field equation). (i) Choosing $F = \varphi(f_2)\cdots\varphi(f_n)$ (allowed by approximation, since all moments are finite) gives the hierarchy for the Schwinger functions $S_n(x_1, \dots, x_n) = \mathbb{E}[\varphi(x_1)\cdots\varphi(x_n)]$:

$$(m^2 - \Delta_{x_1}) S_n(x_1, \dots, x_n) = \sum_{j=2}^{n} \delta(x_1 - x_j)\, S_{n-2}(x_2, \dots, \widehat{x_j}, \dots, x_n), \tag{3.11}$$

which together with $S_0 = 1$, $S_1 = 0$ determines all moments (Wick's theorem, Exercise 3.7). Compare with (1.5): in imaginary time, $\Box + m^2$ becomes $-\Delta + m^2$, and the contact terms on the right come from the CCR, i.e. from time ordering. (ii) For a formal Gibbs measure $\mu_V \propto \mathrm{e}^{-\int V(\varphi)}\mu_m$ the same computation gives

$$\mathbb{E}_{\mu_V}[D_h F] = \mathbb{E}_{\mu_V}\big[F\,\big((m^2 - \Delta)\varphi + V'(\varphi)\big)(h)\big], \tag{3.12}$$

the Euclidean counterpart of the nonlinear field equation (1.9). Here $V'(\varphi)$ must be a well-defined local operator: for Φ^4_2, $V' = 4\lambda{:}\varphi^3{:}$. For Φ^4_3 the renormalised equation (3.12) is a theorem [GH21].

The DS equation is not only satisfied by the GFF; it characterises it.

Theorem 3.12 (DS characterises the GFF). *Let ν be a probability measure on $\mathcal{S}'(\mathcal{M})$ such that $\varphi(f) \in L^1(\nu)$ for all $f \in \mathcal{S}(\mathcal{M})$ and*

$$\mathbb{E}_\nu[D_h F] = \mathbb{E}_\nu[F\,\varphi(Ah)] \qquad \text{for all } F \in \mathcal{FC}_b^\infty,\ h \in \mathcal{S}(\mathcal{M}). \tag{3.13}$$

Then $\nu = \mu_m$.

Proof. Fix a real $f \in \mathcal{S}(\mathcal{M})$ and let $h = Cf \in \mathcal{S}(\mathcal{M})$, so that $Ah = f$. For $s \in \mathbb{R}$ apply (3.13) to the real and imaginary parts of $F_s(\varphi) = \mathrm{e}^{\mathrm{i}s\varphi(f)}$, which belong to $\mathcal{FC}_b^\infty$, using $D_h F_s = \mathrm{i}s\langle f, Cf\rangle F_s$. With $\chi(s) := \mathbb{E}_\nu[F_s]$ this gives

$$\mathrm{i}s\langle f, Cf\rangle\,\chi(s) = \mathbb{E}_\nu\big[\varphi(f)\,\mathrm{e}^{\mathrm{i}s\varphi(f)}\big] = -\mathrm{i}\chi'(s),$$

where we used $\varphi(f) \in L^1(\nu)$ to differentiate under the expectation. Hence $\chi'(s) = -s\langle f, Cf\rangle\chi(s)$ with $\chi(0) = 1$, so $\chi(s) = \mathrm{e}^{-\frac{s^2}{2}\langle f, Cf\rangle}$. Taking $s = 1$: $\int \mathrm{e}^{\mathrm{i}\varphi(f)}\mathrm{d}\nu = \mathrm{e}^{-\frac{1}{2}\langle f, Cf\rangle}$ for every f. Since $\sum_j u_j\varphi(f_j) = \varphi(\sum_j u_j f_j)$, this determines the characteristic function of every vector $(\varphi(f_1), \dots, \varphi(f_k))$, hence all finite-dimensional distributions, hence ν on the cylinder σ-algebra. So $\nu = \mu_m$. □

Remark 3.13. The integrability assumption is the minimal one needed for (3.13) to make sense, since $\varphi(Ah)$ ranges over all $\varphi(g)$, $g \in \mathcal{S}(\mathcal{M})$. Only test functions $F = \mathrm{e}^{\mathrm{i}s\varphi(f)}$ were used: DS turns into a first-order linear ODE for the characteristic function. For $m = 0$ on the torus no probability measure solves (3.13) (Exercise 3.9). For interacting theories, (3.12) is a singular nonlinear relation, and uniqueness of its solutions is a hard and largely open problem. In finite dimensions it is easy (Exercise 3.8). For a result in this direction for the exponential interaction in $d = 2$ see [DGT22].

3.5 The GFF as stationary marginal of parabolic stochastic quantisation

Langevin dynamics. If $S : \mathbb{R}^N \to \mathbb{R}$ is smooth with $\mathrm{e}^{-S} \in L^1$, the diffusion $\mathrm{d}X_t = -\nabla S(X_t)\mathrm{d}t + \sqrt{2}\,\mathrm{d}B_t$ has generator $\mathcal{L} = \Delta - \nabla S \cdot \nabla$, and $\mathcal{L}^*\mathrm{e}^{-S} = \nabla \cdot (\nabla \mathrm{e}^{-S} + \mathrm{e}^{-S}\nabla S) = 0$: the Gibbs measure $\mathrm{e}^{-S(x)}\mathrm{d}x/Z$ is invariant. Parisi and Wu [PW81] proposed to apply this to the Euclidean action, introducing a fictitious "stochastic time" t:

$$\partial_t \Phi = -(m^2 - \Delta)\Phi - V'(\Phi) + \sqrt{2}\,\xi \qquad \text{on } \mathbb{R}_+ \times \mathcal{M}, \tag{3.14}$$

with ξ space-time white noise. This is *(parabolic) stochastic quantisation*. For $V = 0$ it is the stochastic heat equation, an infinite-dimensional Ornstein–Uhlenbeck process.

Space-time white noise on $\mathbb{R} \times \mathcal{M}$ is a centred Gaussian family $(\xi(u))_{u \in L^2(\mathbb{R}\times\mathcal{M})}$ with $\mathbb{E}[\xi(u)\xi(v)] = \langle u, v\rangle$, formally $\mathbb{E}[\xi(t,x)\xi(s,y)] = \delta(t-s)\delta(x-y)$. Let $W_t(h) = \xi(\mathbb{1}_{[0,t]} \otimes h)$, $t \geq 0$, a cylindrical Brownian motion. For a (weak) solution Φ of (3.14) with $V = 0$ and $F = g(\varphi(f_1), \dots, \varphi(f_n)) \in \mathcal{FC}_b^\infty$, Itô's formula shows that $F(\Phi_t) - \int_0^t \mathcal{L}F(\Phi_r)\mathrm{d}r$ is a martingale, with

$$\mathcal{L}F(\varphi) = \sum_{i,j}\langle f_i, f_j\rangle\,\partial_{ij} g(\dots) - \sum_i \varphi(Af_i)\,\partial_i g(\dots). \tag{3.15}$$

Proposition 3.14 (DS ⇒ invariance). *For $F, G \in \mathcal{FC}_b^\infty$: $\mathbb{E}_{\mu_m}[G\,\mathcal{L}F] = -\mathbb{E}_{\mu_m}[\langle DG, DF\rangle_{L^2(\mathcal{M})}]$. In particular $\mathbb{E}_{\mu_m}[\mathcal{L}F] = 0$.*

Proof. We may assume $F = g(\varphi(f_1), \dots)$ and $G = k(\varphi(f_1), \dots)$ with the same f_i. Apply (3.10) to the cylinder function $k\,\partial_i g$ with $h = f_i$ and sum over i: $\sum_i \mathbb{E}[k\,\partial_i g\,\varphi(Af_i)] = \sum_{i,j}\langle f_i, f_j\rangle\mathbb{E}[\partial_j k\,\partial_i g + k\,\partial_{ij} g]$. Insert this into $\mathbb{E}[G\mathcal{L}F]$. □

So the static characterisation (DS) and the dynamic one (invariance under the Langevin flow) are two faces of the same integration by parts formula.

Theorem 3.15 (Parabolic stochastic quantisation of the GFF)**.** *Let ξ be space-time white noise on $\mathbb{R}\times\mathcal{M}$. For $t\in\mathbb{R}$ and $f\in\mathcal{S}(\mathcal{M})$ define*

$$\Phi_t(f) := \sqrt{2}\,\xi(K_{t,f}), \qquad K_{t,f}(r,x) := \mathbb{1}_{\{r<t\}}\big(\mathrm{e}^{-(t-r)A}f\big)(x). \tag{3.16}$$

(a) Φ is a centred Gaussian process with $\mathbb{E}[\Phi_t(f)\Phi_s(g)] = \langle f, \mathrm{e}^{-|t-s|A}Cg\rangle$. In particular $(\Phi_t)_{t\in\mathbb{R}}$ is stationary and, for every t, Φ_t has a version with values in $\mathcal{S}'(\mathcal{M})$ whose law is μ_m.

(b) Φ is a mild and weak solution of (3.14) with $V=0$: for $s<t$ and $h\in\mathcal{S}(\mathcal{M})$,

$$\Phi_t(h) = \Phi_s(\mathrm{e}^{-(t-s)A}h) + \sqrt{2}\,\xi\big(\mathbb{1}_{(s,t)}(r)\,\mathrm{e}^{-(t-r)A}h\big),$$
$$\Phi_t(h) = \Phi_s(h) - \int_s^t \Phi_r(Ah)\,\mathrm{d}r + \sqrt{2}\,\xi\big(\mathbb{1}_{(s,t)}\otimes h\big).$$

(c) Let Ψ_0 be an $\mathcal{S}'(\mathcal{M})$-valued random variable independent of ξ, and let Ψ be the mild solution started from Ψ_0,

$$\Psi_t(f) = \Psi_0(\mathrm{e}^{-tA}f) + \sqrt{2}\,\xi\big(\mathbb{1}_{(0,t)}(r)\,\mathrm{e}^{-(t-r)A}f\big), \qquad t\geq 0.$$

Then $\mathbb{E}[\mathrm{e}^{\mathrm{i}\Psi_t(f)}] \to \mathrm{e}^{-\frac{1}{2}\langle f,Cf\rangle}$ as $t\to\infty$ for every $f\in\mathcal{S}(\mathcal{M})$. Moreover μ_m is invariant ($\Psi_0\sim\mu_m$ implies $\Psi_t\sim\mu_m$ for all t), and it is the unique invariant probability measure of the dynamics.

Proof. (a) Since A is self-adjoint and commutes with e^{-uA},

$$\langle K_{t,f}, K_{s,g}\rangle = \int_{-\infty}^{t\wedge s}\langle f, \mathrm{e}^{-(t+s-2r)A}g\rangle\mathrm{d}r = \langle f, \mathrm{e}^{-|t-s|A}(2A)^{-1}g\rangle.$$

In particular $K_{t,f}\in L^2(\mathbb{R}\times\mathcal{M})$, and multiplying by 2 gives the covariance. Stationarity follows. For fixed t, $f\mapsto\Phi_t(f)$ is linear and $\mathbb{E}[\Phi_t(f)^2] = \langle f,Cf\rangle \leq m^{-2}\|f\|_{L^2}^2$, so it is continuous from $\mathcal{S}(\mathcal{M})$ to $L^2(\mathbb{P})$. By the Minlos regularisation theorem for random linear functionals on nuclear spaces it has an $\mathcal{S}'(\mathcal{M})$-valued version, which is centred Gaussian with covariance C, i.e. has law μ_m. On the torus one can see this directly: the coordinates $\Phi_t(e_n)$ are independent stationary OU processes $\mathrm{d}X = -\lambda_n X\mathrm{d}t + \sqrt{2}\mathrm{d}B$ with variance λ_n^{-1}, and (3.3) applies.

(b) The mild formula is the identity $K_{t,h} = K_{s,\mathrm{e}^{-(t-s)A}h} + \mathbb{1}_{(s,t)}(r)\mathrm{e}^{-(t-r)A}h$ together with the linearity of ξ. The weak formulation follows from the mild one by a stochastic Fubini argument (Exercise 3.12).

(c) Since Ψ_0 is independent of ξ,

$$\mathbb{E}\big[\mathrm{e}^{\mathrm{i}\Psi_t(f)}\big] = \mathbb{E}\big[\mathrm{e}^{\mathrm{i}\Psi_0(\mathrm{e}^{-tA}f)}\big]\,\mathrm{e}^{-\frac{1}{2}\langle f,(1-\mathrm{e}^{-2tA})Cf\rangle}.$$

For $m>0$, $\mathrm{e}^{-tA}f\to 0$ in $\mathcal{S}(\mathcal{M})$ as $t\to\infty$, so $\Psi_0(\mathrm{e}^{-tA}f)\to 0$ a.s. because $\Psi_0\in\mathcal{S}'(\mathcal{M})$. By dominated convergence the first factor tends to 1, while the second tends to $\mathrm{e}^{-\frac{1}{2}\langle f,Cf\rangle}$. If $\Psi_0\sim\mu_m$, the first factor equals $\mathrm{e}^{-\frac{1}{2}\langle f,\mathrm{e}^{-2tA}Cf\rangle}$ and the product is exactly $\mathrm{e}^{-\frac{1}{2}\langle f,Cf\rangle}$, which proves invariance. If ν is invariant, start from $\Psi_0\sim\nu$: then $\mathrm{Law}(\Psi_t)=\nu$ for all t, so the characteristic functional of ν equals that of μ_m, and $\nu=\mu_m$. □




Figure 3: Parabolic stochastic quantisation of the GFF on a circle of length 8, $m = 1$, started from $\Phi_0 = 0$. Left: a sample of the solution. Right: each Fourier mode is an Ornstein–Uhlenbeck process, whose variance relaxes to λ_k^{-1} at rate $2\lambda_k$ (Theorem 3.15(c)). Dots: Monte Carlo estimates with 4000 samples.

Remark 3.16. As a Gaussian field on $\mathbb{R} \times \mathcal{M}$, the stationary solution has covariance $2(\partial_t + A)^{-1}(-\partial_t + A)^{-1} = 2(-\partial_t^2 + A^2)^{-1}$, and its equal-time slice has covariance $2\int_{\mathbb{R}}(\theta^2 + A^2)^{-1}\frac{\mathrm{d}\theta}{2\pi} = A^{-1}$. The stochastic time t is unrelated to the Euclidean time τ. For interacting theories, (3.14) is a singular SPDE. It was solved by Da Prato and Debussche for Φ^4_2 [DPD03] (see also [JLM85, AR91]), and for Φ^4_3 using regularity structures and paracontrolled calculus [Hai14, GIP15, CC18]. Global-in-time bounds [MW17, GH19] then allow constructing the Euclidean Φ^4_3 theory as the invariant measure of the dynamics [GH21].

3.6 The GFF as one-point marginal of elliptic stochastic quantisation

Parisi and Sourlas [PS79] discovered that the elliptic SPDE in $d+2$ dimensions

$$(m^2 - \Delta_{\mathbb{R}^2\times\mathcal{M}})\Psi + V'(\Psi) = \zeta, \qquad \zeta \text{ white noise on } \mathbb{R}^2 \times \mathcal{M}, \tag{3.17}$$

is related to the d-dimensional Gibbs measure: formally, the law of $\Psi(z,\cdot)$, $z \in \mathbb{R}^2$, is $\propto \exp\big(-4\pi\big[\frac{1}{2}\langle\varphi, A\varphi\rangle + \int V(\varphi)\big]\big)\mathcal{D}\varphi$ (*dimensional reduction*). The underlying mechanism is a hidden supersymmetry [PS79, KLP84]. In [KLP84] the dimensional reduction of the supersymmetric representation is proved rigorously, but not the supersymmetric representation of the solution of the SPDE; the full nonlinear statement was proved for $d = 0$ in [ADG20] and for some two-dimensional models (polynomial interactions on $\mathbb{T}^2$, exponential interactions on $\mathbb{R}^2$) in [ADG21]; elliptic Φ^4 equations in 4 and 5 dimensions are solved in [GH19]. For $V = 0$ the statement is elementary, and it explains the factor 4π. We normalise the noise so that the slice is exactly μ_m.

Theorem 3.17 (Elliptic stochastic quantisation of the GFF)**.** *Let ζ be white noise on $\mathbb{R}^2 \times \mathcal{M}$ (an $\mathcal{S}'(\mathbb{R}^2 \times \mathcal{M})$-valued Gaussian variable with $\mathbb{E}[\zeta(u)\zeta(v)] = \langle u, v\rangle$) and $\mathsf{L} := m^2 - \Delta_{\mathbb{R}^2\times\mathcal{M}} = -\Delta_{\mathbb{R}^2} + A$.*

(a) The equation $\mathsf{L}\Psi = \sqrt{4\pi}\,\zeta$ has a unique $\mathcal{S}'(\mathbb{R}^2 \times \mathcal{M})$-valued solution, $\Psi(u) = \sqrt{4\pi}\,\zeta(\mathsf{L}^{-1}u)$.

(b) *For $z \in \mathbb{R}^2$ and $f \in \mathcal{S}(\mathcal{M})$ the function $K_{z,f} := \mathsf{L}^{-1}(\delta_z \otimes f)$ lies in $L^2(\mathbb{R}^2 \times \mathcal{M})$, and $z \mapsto K_{z,f}$ is continuous. The Wiener integrals $\Psi(z,f) := \sqrt{4\pi}\,\zeta(K_{z,f})$ satisfy $\Psi(v \otimes f) = \int_{\mathbb{R}^2} v(z)\Psi(z,f)\,\mathrm{d}z$ a.s. for $v \in \mathcal{S}(\mathbb{R}^2)$.*

(c) *For all $z \in \mathbb{R}^2$ and $f, g \in \mathcal{S}(\mathcal{M})$, $\mathbb{E}[\Psi(z,f)\Psi(z,g)] = \langle f, Cg\rangle$. Hence $\Psi(z,\cdot)$ has an $\mathcal{S}'(\mathcal{M})$-valued version with law μ_m, and the field $(\Psi(z,\cdot))_{z\in\mathbb{R}^2}$ is stationary and isotropic in z.*

Proof. (a) L is a continuous bijection of $\mathcal{S}(\mathbb{R}^2 \times \mathcal{M})$, so $\Psi(u) := \sqrt{4\pi}\zeta(\mathsf{L}^{-1}u)$ defines an $\mathcal{S}'$-valued variable with $\Psi(\mathsf{L}u) = \sqrt{4\pi}\zeta(u)$. If $D \in \mathcal{S}'$ satisfies $D(\mathsf{L}u) = 0$ for all $u \in \mathcal{S}$, then $D = 0$ since L is onto: the solution is unique.

(b), (c) Take the Fourier transform in z only, with measure $\mathrm{d}q/(2\pi)^2$. The transform of $K_{z,f}$ is the $L^2(\mathcal{M})$-valued function $q \mapsto \mathrm{e}^{-\mathrm{i}q\cdot z}(|q|^2 + A)^{-1}f$. By Plancherel, the spectral theorem for A and Tonelli,

$$\langle K_{z,f}, K_{z,g}\rangle = \int_{\mathbb{R}^2} \langle f, (|q|^2+A)^{-2}g\rangle \frac{\mathrm{d}q}{(2\pi)^2} = \Big\langle f, \Big[\int_{\mathbb{R}^2} \frac{\mathrm{d}q}{(2\pi)^2}(|q|^2+A)^{-2}\Big]g\Big\rangle = \frac{1}{4\pi}\langle f, A^{-1}g\rangle,$$

using the scalar identity

$$\int_{\mathbb{R}^2} \frac{\mathrm{d}q}{(2\pi)^2}\frac{1}{(|q|^2+a)^2} = \frac{1}{2\pi}\int_0^\infty \frac{r\,\mathrm{d}r}{(r^2+a)^2} = \frac{1}{4\pi a}, \qquad a > 0. \tag{3.18}$$

In particular $K_{z,f} \in L^2$, and its continuity in z follows by dominated convergence. Since the transform of $\int v(z)K_{z,f}\mathrm{d}z$ is $\hat{v}(q)(|q|^2+A)^{-1}f$, the Bochner integral $\int v(z)K_{z,f}\mathrm{d}z$ equals $\mathsf{L}^{-1}(v \otimes f)$. The Wiener integral ζ is an isometry $L^2 \to L^2(\mathbb{P})$ and commutes with Bochner integrals, which gives (b). Multiplying by 4π gives the covariance in (c). The $\mathcal{S}'(\mathcal{M})$-valued version is obtained as in Theorem 3.15. Stationarity and isotropy hold because $\mathbb{E}[\Psi(z,f)\Psi(z',g)] = 4\pi\int \mathrm{e}^{\mathrm{i}q\cdot(z-z')}\langle f, (|q|^2+A)^{-2}g\rangle\frac{\mathrm{d}q}{(2\pi)^2}$ depends only on $|z-z'|$. □

Remark 3.18 (Slices of Gaussian fields). Both stochastic quantisations exhibit the GFF as a *slice* of a Gaussian field in higher dimension. For the field on $\mathbb{R}^k \times \mathcal{M}$ with covariance $(-\Delta_{\mathbb{R}^k} + A)^{-p}$, the slice $\{0\} \times \mathcal{M}$ has covariance $c_{k,p}A^{k/2-p}$ with $c_{k,p} = \Gamma(p - \frac{k}{2})/((4\pi)^{k/2}\Gamma(p))$ (Exercise 3.13). The GFF arises iff $p = 1 + k/2$: $(k,p) = (2,2)$ is elliptic SQ, while $(k,p) = (1,1)$ gives instead the time-zero field $(2\omega)^{-1}$ of Proposition 3.4. When $\mathcal{M} = \mathbb{R}^d$, the elliptic field Ψ is Euclidean invariant in $\mathbb{R}^{d+2}$, and its restriction to *every* codimension-two affine subspace is a GFF.

3.7 Outlook

In Lecture 3 we decompose the GFF along a scale parameter: the field at resolution t is a Gaussian martingale, whose terminal value is the GFF. Interactions are then introduced through the Polchinski equation for the effective potential, and through its probabilistic form, a forward–backward stochastic differential equation for the field at increasing resolution. These tools can be used to construct Φ^4_2 and Φ^4_3. On $\mathbb{T}^2$ the Wick-ordered interaction $\lambda\int{:}\varphi^4{:}\mathrm{d}x$ suffices, and the measure satisfies the renormalised DS equation (3.12). In $d = 3$ Wick ordering no longer suffices.

Exercises

Exercise 3.1 (Green functions)**.** (a) Using $G_m = \int_0^\infty \mathrm{e}^{-m^2t}p_t\,\mathrm{d}t$, with p_t the heat kernel, show that $G_m(x) = (2\pi)^{-d/2}(m/|x|)^{d/2-1}K_{d/2-1}(m|x|)$, and derive the asymptotics stated after Definition 3.1. (b) Check $G_m(x) = \mathrm{e}^{-m|x|}/(2m)$ for $d = 1$ and $G_m(x) = \mathrm{e}^{-m|x|}/(4\pi|x|)$ for $d = 3$. (c) Show that on $\mathbb{T}^d_L$ the difference $G - G_m$ is smooth near the diagonal.

Exercise 3.2 (Euclidean Källén–Lehmann and the unitarity bound)**.** (a) Show that

$$\int \mathrm{d}\Omega_M(\mathbf{p})\, \mathrm{e}^{-\omega_M(\mathbf{p})|\tau|+\mathrm{i}\mathbf{p}\cdot\mathbf{x}} = G_M(\tau,\mathbf{x}),$$

and derive (3.5) from (1.4). (b) Show that $G_M(x)$ is decreasing in M and, for $d \geq 2$, that $G_M(x)/G_m(x) \to 1$ as $x \to 0$. Prove the lower bound in Remark 3.5(i). (c) Let $d \geq 3$ and suppose $S_2^T(x) = |x|^{-2\Delta_\varphi}$ (a scale invariant theory). Using $G_M(x) = M^{d-2}G_1(Mx)$, show that $\rho(\mathrm{d}M^2) = c\,(M^2)^{\Delta_\varphi - d/2}\mathrm{d}M^2$ reproduces S_2^T for some $c > 0$, for $\frac{d-2}{2} < \Delta_\varphi < \frac{d}{2}$, and that this is a locally finite measure iff $\Delta_\varphi > \frac{d-2}{2}$ (for $\Delta_\varphi \geq \frac{d}{2}$, S_2^T is not locally integrable). Using uniqueness of the Stieltjes transform, deduce the unitarity bound $\Delta_\varphi \geq \frac{d-2}{2}$, with equality iff $\rho \propto \delta_0$, the massless free field.

Exercise 3.3 (Regularity is sharp)**.** (a) For $a_n \geq 0$ and (g_n) i.i.d. $N(0,1)$, show that $\mathbb{E}\,\mathrm{e}^{-\sum_n a_n g_n^2} = \prod_n (1+2a_n)^{-1/2}$. Deduce that $\sum_n a_n g_n^2 < \infty$ a.s. iff $\sum_n a_n < \infty$. (b) Conclude that $\mu_m(H^{-s}(\mathbb{T}_L^d)) = 1$ iff $s > \frac{d}{2} - 1$. In particular $\mu_m(L^2) = 0$ for $d \geq 2$, and the Cameron–Martin space H^1 is μ_m-null in every dimension. (c*) Using hypercontractivity and a Kolmogorov argument, show that $\varphi \in C^{1-d/2-\kappa}$ a.s. for every $\kappa > 0$.

Exercise 3.4 (Hermite calculus and change of renormalisation)**.** (a) Show that $H_n(a;c) = \sum_{k \leq n/2} \frac{n!}{k!\,(n-2k)!}\big(-\frac{c}{2}\big)^k a^{n-2k}$, $\partial_a H_n = nH_{n-1}$ and $aH_n = H_{n+1} + ncH_{n-1}$. (b) Show that $H_n(a+b;c+c') = \sum_k \binom{n}{k} H_k(a;c)H_{n-k}(b;c')$. (c) Show that $H_n(a;c') = \sum_k \binom{n}{2k}(2k-1)!!\,(c-c')^k H_{n-2k}(a;c)$. Interpret this as the effect of changing the mollifier on $:\varphi^n:$. Wick powers are defined up to lower order Wick powers with *finite* coefficients.

Exercise 3.5 (Wick powers: details)**.** (a) Show that $c_\varepsilon = \frac{1}{2\pi}\log\frac{1}{\varepsilon} + O(1)$ for $d = 2$ and $c_\varepsilon \asymp \varepsilon^{2-d}$ for $d \geq 3$. (b) In $d = 3$, show that $\mathbb{E}[:\varphi_\varepsilon^3:(f)^2] \asymp \log\frac{1}{\varepsilon}$ for $0 \leq f \not\equiv 0$: the cube is borderline divergent. (c) In $d = 4$, show that $\mathbb{E}[:\varphi_\varepsilon^2:(f)^2] \asymp \log\frac{1}{\varepsilon}$ as well.

Exercise 3.6 (Wick exponentials)**.** In $d = 2$ let $:\mathrm{e}^{\mathrm{i}\beta\varphi_\varepsilon}:(x) := \mathrm{e}^{\beta^2 c_\varepsilon/2}\mathrm{e}^{\mathrm{i}\beta\varphi_\varepsilon(x)}$. Show that

$$\mathbb{E}\big[:\mathrm{e}^{\mathrm{i}\beta\varphi_\varepsilon}:(x)\,\overline{:\mathrm{e}^{\mathrm{i}\beta\varphi_\delta}:(y)}\big] = \mathrm{e}^{\beta^2 G_{\varepsilon,\delta}(x-y)},$$

and deduce that $:\mathrm{e}^{\mathrm{i}\beta\varphi_\varepsilon}:(f)$ converges in $L^2(\mu_m)$ for every f if $\beta^2 < 4\pi$, while its second moment diverges for $0 \leq f \not\equiv 0$ if $\beta^2 \geq 4\pi$. What changes for the real exponential $:\mathrm{e}^{\alpha\varphi}:$? (Gaussian multiplicative chaos exists in L^1 for $\alpha^2 < 8\pi$, see [BP25].)

Exercise 3.7 (Wick's theorem from DS)**.** (a) Complete the proof of Lemma 3.9. (b) Derive (3.11) from (3.10), justifying the use of polynomial F. (c) Deduce Isserlis–Wick: $S_{2n+1} = 0$ and $S_{2n}(x_1,\dots,x_{2n}) = \sum_{\text{pairings } P}\prod_{\{i,j\}\in P} G(x_i - x_j)$.

Exercise 3.8 (DS in finite dimensions)**.** Let $S(x) = \frac{1}{2}\langle x, Qx\rangle + V(x)$ on $\mathbb{R}^N$, with $Q > 0$ and V smooth with e^{-V} bounded, and $\mu_V(\mathrm{d}x) = Z^{-1}\mathrm{e}^{-S(x)}\mathrm{d}x$. (a) Show that $\mathbb{E}_{\mu_V}[\partial_h F] = \mathbb{E}_{\mu_V}[F\langle \nabla S, h\rangle]$ for $F \in C_c^\infty$. (b) Let ν be a probability measure with $\nabla S \in L^1(\nu)$ satisfying the same identity. Show that $\nu = \mu_V$. *Hint:* the identity says that the distribution $\mathrm{e}^S\nu$ has zero gradient. (c) Why does this argument fail in infinite dimensions? What replaces it in Theorem 3.12?

Exercise 3.9 (The massless case)**.** (a) Show that for $m = 0$ on $\mathbb{T}^d$ no probability measure on $\mathcal{S}'(\mathbb{T}^d)$ satisfies (3.13) with $A = -\Delta$. *Hint:* take $h = 1$. (b) Show that the massless GFF exists on the quotient $\mathcal{S}'(\mathbb{T}^d)/\mathbb{R}$ (test functions with $\int f = 0$), and characterise it by DS there. (c) On $\mathbb{R}^2$, show that $\langle f, (-\Delta)^{-1}f\rangle < \infty$ iff $\hat f(0) = 0$: the infrared problem of the two-dimensional massless field.

Exercise 3.10 (Infinitesimal invariance characterises μ_m)**.** On $\mathbb{T}_L^d$, let ν be a probability measure on $\mathcal{S}'(\mathbb{T}_L^d)$ with $\varphi(f) \in L^1(\nu)$ and $\mathbb{E}_\nu[\mathcal{L}F] = 0$ for all $F \in \mathcal{F}C_b^\infty$. Show that $\nu = \mu_m$. *Hint:* for $f = \sum_{n \le N} a_n e_n$ and $\psi(a) = \mathbb{E}_\nu e^{\mathrm{i}\varphi(f)}$, show $\sum_n \lambda_n a_n \partial_{a_n} \psi = -|a|^2 \psi$, and differentiate $s \mapsto \psi(e^{sD}a)\exp(\frac{1}{2}\sum_n a_n^2 e^{2\lambda_n s}/\lambda_n)$, where $D = \mathrm{diag}(\lambda_n)$.

Exercise 3.11 (Reversibility)**.** Use Proposition 3.14 to show that $\mathcal{L}$ is symmetric in $L^2(\mu_m)$. Deduce, and check directly from the covariance in Theorem 3.15(a), that the stationary solution is time reversible: $(\Phi_{-t})_{t\in\mathbb{R}}$ and $(\Phi_t)_{t\in\mathbb{R}}$ have the same law.

Exercise 3.12 (Mild implies weak)**.** Prove the weak formulation in Theorem 3.15(b). Show that, as a random distribution on $\mathbb{R}\times\mathcal{M}$, the stationary solution has covariance operator $2(-\partial_t^2 + A^2)^{-1}$.

Exercise 3.13 (Slices of Gaussian fields)**.** Let $k \ge 1$, $p > k/2$, and let Ψ be the centred Gaussian field on $\mathbb{R}^k \times \mathcal{M}$ with covariance $(-\Delta_{\mathbb{R}^k} + A)^{-p}$. (a) Show that $\Psi(0,\cdot)$ is well defined with covariance $c_{k,p} A^{k/2-p}$, $c_{k,p} = \Gamma(p - \frac{k}{2})/((4\pi)^{k/2}\Gamma(p))$. (b) For $(k,p) = (1,1)$ and $\mathcal{M} = N$, show that the time-zero field of the GFF on $\mathbb{R}\times N$ is the ground state measure $\Omega_0(\varphi)^2\mathcal{D}\varphi$ of the free Hamiltonian, i.e. Gaussian with covariance $(2\omega)^{-1}$ (compare Exercise 2.2). (c) Show that the restriction of the elliptic field of Theorem 3.17 to a codimension-*one* slice $\mathbb{R}\times\mathcal{M}$ has covariance proportional to $(-\partial^2 + A)^{-3/2}$, so it is not a GFF. (d) Show that $\mathbb{E}[\Psi(z,f)\Psi(0,g)] = \langle f, |z|A^{-1/2}K_1(|z|A^{1/2})g\rangle$ in Theorem 3.17, and recover $\langle f, Cg\rangle$ as $z \to 0$.

Exercise 3.14 (Lattice approximation)**.** Let μ_ε be the centred Gaussian measure on $\mathbb{R}^{(\varepsilon\mathbb{Z}/L\mathbb{Z})^d}$ with covariance $(m^2 - \Delta_\varepsilon)^{-1}$ (with respect to $\varepsilon^d\sum_u$), embedded in $\mathcal{S}'(\mathbb{T}_L^d)$ via $\varphi(f) = \varepsilon^d\sum_u \varphi_u f(u)$. Show that $\mu_\varepsilon \to \mu_m$ weakly. Relate μ_ε to (2.12) with $V = 0$ (for $d = 2$; in general d see Exercise 2.3).

Exercise 3.15 (Computer experiments)**.** (a) Sample the lattice GFF of Exercise 3.14 on $\mathbb{T}^2$ with $\varepsilon = 1/N$ (i.e. on $(\mathbb{Z}/N\mathbb{Z})^2$ with mass m/N in lattice units) by FFT, for $N = 64, 256, 1024$, and check that the empirical variance of φ_u grows like $\frac{1}{2\pi}\log N$. (b) Implement an explicit Euler scheme for the lattice version of (3.14) with $V = 0$ (mind the stability constraint $\Delta t \lesssim \varepsilon^2$), and check that the variance of the k-th Fourier mode relaxes to λ_k^{-1} at rate $2\lambda_k$, with λ_k the eigenvalues of $m^2 - \Delta_\varepsilon$. (c) Add $V'(\varphi) = 4\lambda(\varphi^3 - 3c_\varepsilon\varphi)$, i.e. $V = \lambda{:}\varphi^4{:}$, and observe that the solutions have a nontrivial limit as $N \to \infty$, while without the counterterm $-12\lambda c_\varepsilon\varphi$ they converge to zero [HRW12].

Exercise 3.16 (Two masses)**.** In the setting of the example after Remark 3.5: (a) Show that μ_ρ satisfies (OS0)–(OS4), and that the theory described there has Schwinger functions equal to the moments of μ_ρ. (b) Show that $[\varphi(0,\mathbf{x}), \partial_t\varphi(0,\mathbf{y})] = \mathrm{i}(c_1^2 + c_2^2)\delta(\mathbf{x}-\mathbf{y})$, and that $\varphi(t,\cdot)$ is not a linear function of $\varphi(0,\cdot)$ and $\partial_t\varphi(0,\cdot)$. (c) Verify $\mathbb{E}[D_{Rh}F] = \mathbb{E}[F\varphi(Ph)]$. (d) For a single spatial Fourier mode $\mathbf{p}$ (on a torus, say, with real test functions $f(\tau,\mathbf{x}) = g(\tau)\cos(\mathbf{p}\cdot\mathbf{x})$), with $\omega_i^2 = |\mathbf{p}|^2 + m_i^2$, show that the time kernel of P^{-1} is $\big(e^{-\omega_1|\tau|}/2\omega_1 - e^{-\omega_2|\tau|}/2\omega_2\big)/(m_2^2 - m_1^2)$, and find f supported in $\{\tau > 0\}$ with $\langle \Theta f, P^{-1}f\rangle < 0$. *Hint:* choose g with $\int_0^\infty e^{-\omega_1\tau}g(\tau)\mathrm{d}\tau = 0$. (e) Let p be a real polynomial of degree ≥ 2, positive on $[0,\infty)$. Show that $1/p(|k|^2)$ is not of the form $\int \rho(\mathrm{d}M^2)(|k|^2 + M^2)^{-1}$ with $\rho \ge 0$. *Hint:* $s\int\rho(\mathrm{d}M^2)(s+M^2)^{-1}$ increases to $\rho([0,\infty))$ as $s \to \infty$ (monotone convergence), while $s/p(s) \to 0$.

References and further reading

Guide. *Gaussian fields and Wick calculus:* [GV64, Bog98, Jan97, She07, LSSW16, BP25]. *Euclidean free field and Markov property:* [Nel73a, Nel73b, GJ87, Sim74]. *Stochastic quantisation:*

parabolic [PW81, JLM85, AR91, DPD03, DPZ14]; singular SPDEs [Hai14, GIP15, CC18, MW17, HRW12]; construction of Φ^4_3 via dynamics and variational methods [GH21, BG20]; integration by parts and DS equations for the exponential model in $d = 2$ [DGT22]; elliptic stochastic quantisation and dimensional reduction [PS79, KLP84, ADG20, ADG21, GH19].

[ADG20] S. Albeverio, F. C. De Vecchi, M. Gubinelli. Elliptic stochastic quantization. *Ann. Probab.* 48 (2020), 1693–1741.

[ADG21] S. Albeverio, F. C. De Vecchi, M. Gubinelli. The elliptic stochastic quantization of some two dimensional Euclidean QFTs. *Ann. Inst. Henri Poincaré Probab. Stat.* 57 (2021), 2372–2414.

[AR91] S. Albeverio, M. Röckner. Stochastic differential equations in infinite dimensions: solutions via Dirichlet forms. *Probab. Theory Related Fields* 89 (1991), 347–386.

[BG20] N. Barashkov, M. Gubinelli. A variational method for Φ^4_3. *Duke Math. J.* 169 (2020), 3339–3415.

[Bog98] V. I. Bogachev. *Gaussian Measures*. Amer. Math. Soc., 1998.

[BP25] N. Berestycki, E. Powell. *Gaussian Free Field and Liouville Quantum Gravity*. Cambridge University Press, 2025.

[BW86] D. Buchholz, E. H. Wichmann. Causal independence and the energy-level density of states in local quantum field theory. *Comm. Math. Phys.* 106 (1986), 321–344.

[CC18] R. Catellier, K. Chouk. Paracontrolled distributions and the 3-dimensional stochastic quantization equation. *Ann. Probab.* 46 (2018), 2621–2679.

[DGT22] F. C. De Vecchi, M. Gubinelli, M. Turra. A singular integration by parts formula for the exponential Euclidean QFT on the plane. Preprint, arXiv:2212.05584.

[DPD03] G. Da Prato, A. Debussche. Strong solutions to the stochastic quantization equations. *Ann. Probab.* 31 (2003), 1900–1916.

[DPZ14] G. Da Prato, J. Zabczyk. *Stochastic Equations in Infinite Dimensions*, 2nd ed. Cambridge University Press, 2014.

[GH19] M. Gubinelli, M. Hofmanová. Global solutions to elliptic and parabolic Φ^4 models in Euclidean space. *Comm. Math. Phys.* 368 (2019), 1201–1266.

[GH21] M. Gubinelli, M. Hofmanová. A PDE construction of the Euclidean Φ^4_3 quantum field theory. *Comm. Math. Phys.* 384 (2021), 1–75.

[GIP15] M. Gubinelli, P. Imkeller, N. Perkowski. Paracontrolled distributions and singular PDEs. *Forum Math. Pi* 3 (2015), e6.

[GJ87] J. Glimm, A. Jaffe. *Quantum Physics: A Functional Integral Point of View*, 2nd ed. Springer, 1987.

[GV64] I. M. Gel'fand, N. Ya. Vilenkin. *Generalized Functions, Vol. 4: Applications of Harmonic Analysis*. Academic Press, 1964.

[Haa96] R. Haag. *Local Quantum Physics*, 2nd ed. Springer, 1996.

[Hai14] M. Hairer. A theory of regularity structures. *Invent. Math.* 198 (2014), 269–504.

[HRW12] M. Hairer, M. D. Ryser, H. Weber. Triviality of the 2D stochastic Allen–Cahn equation. *Electron. J. Probab.* 17 (2012), no. 39.

[HSw65] R. Haag, J. A. Swieca. When does a quantum field theory describe particles? *Comm. Math. Phys.* 1 (1965), 308–320.

[Jan97] S. Janson. *Gaussian Hilbert Spaces*. Cambridge University Press, 1997.

[JLM85] G. Jona-Lasinio, P. K. Mitter. On the stochastic quantization of field theory. *Comm. Math. Phys.* 101 (1985), 409–436.

[KLP84] A. Klein, L. J. Landau, J. F. Perez. Supersymmetry and the Parisi–Sourlas dimensional reduction: a rigorous proof. *Comm. Math. Phys.* 94 (1984), 459–482.

[LSSW16] A. Lodhia, S. Sheffield, X. Sun, S. S. Watson. Fractional Gaussian fields: a survey. *Probab. Surveys* 13 (2016), 1–56.

[LW69] T. D. Lee, G. C. Wick. Negative metric and the unitarity of the S-matrix. *Nuclear Phys. B* 9 (1969), 209–243.

[MW17] J.-C. Mourrat, H. Weber. The dynamic Φ^4_3 model comes down from infinity. *Comm. Math. Phys.* 356 (2017), 673–753.

[Nel66] E. Nelson. A quartic interaction in two dimensions. In *Mathematical Theory of Elementary Particles*, MIT Press, 1966, 69–73.

[Nel73a] E. Nelson. The free Markoff field. *J. Funct. Anal.* 12 (1973), 211–227.

[Nel73b] E. Nelson. Construction of quantum fields from Markoff fields. *J. Funct. Anal.* 12 (1973), 97–112.

[PS79] G. Parisi, N. Sourlas. Random magnetic fields, supersymmetry, and negative dimensions. *Phys. Rev. Lett.* 43 (1979), 744–745.

[PU50] A. Pais, G. E. Uhlenbeck. On field theories with non-localized action. *Phys. Rev.* 79 (1950), 145–165.

[PW81] G. Parisi, Y.-S. Wu. Perturbation theory without gauge fixing. *Sci. Sinica* 24 (1981), 483–496.

[Roz82] Yu. A. Rozanov. *Markov Random Fields*. Springer, 1982.

[She07] S. Sheffield. Gaussian free fields for mathematicians. *Probab. Theory Related Fields* 139 (2007), 521–541.

[Sim74] B. Simon. *The $P(\varphi)_2$ Euclidean (Quantum) Field Theory*. Princeton University Press, 1974.