A Constructive Introduction to Quantum Field Theory

Lecture 4: The Polchinski flow and forward–backward equations

BY M. GUBINELLI

Mathematical Institute, University of Oxford

November 2026 — provisional

Abstract

Building on the scale decomposition and the stochastic observables of Lecture 3, we introduce interactions through the Polchinski equation for the effective potential. Its probabilistic counterpart is a forward–backward stochastic differential equation (FBSDE) for the field seen at increasing resolution, whose effective force is a martingale. Writing the FBSDE in the coordinate given by the conditional mean of the terminal field removes its drift and integrates the transport part of the force flow exactly. In the gradient case the remaining equation for the force is the Wetterich equation for the effective average action, whose propagator is the conditional covariance. We read the FBSDE as the stochastic Pontryagin system of a control problem, bound the conditional covariance from both sides (Brascamp–Lieb and Cramér–Rao), and derive log-Sobolev inequalities along the localisation (multiscale Bakry–Émery). We then discuss the variational structure of the flow (Boué–Dupuis formula and Hamilton–Jacobi–Bellman equation) in finite and infinite volume, where it becomes a variational principle relative to the FBSDE involving only intensive quantities. The additional material covers the algebraic and perturbative structure of the flow (semigroup, flatness, counterterms for Φ^4_3), the Dyson–Schwinger equations at every scale with Gaussian bounds on their noise, and stochastic localisation (why it is useful, a Poincaré inequality, and the comparison with the multiscale Bakry–Émery criterion). The treatment is formal: we compute as if all objects were smooth and integrable, and we indicate where the real difficulties lie.

Table of contents

Introduction . . . 3

Structure . . . 3
What is formal . . . 3
Conventions . . . 3

Main lecture . . . 3

7 Interactions: the Polchinski equation and FBSDEs . . . 3

7.1 Effective potentials . . . 3
7.2 The Polchinski process . . . 4
7.3 Observables of the interacting theory . . . 6
Tangent observables . . . 6
7.4 The force and a Burgers equation . . . 6
Exercises . . . 7

8 The FBSDE in force coordinates: the Wetterich equation . . . 7

8.1 State and costate . . . 7

Pontryagin's maximum principle . . . 7
The costate is a gradient . . . 8
The Hamiltonian of the field . . . 8
With noise . . . 8
8.2 Removing the predictable drift . . . 8
8.3 The FBSDE in the forecast coordinate . . . 9
8.4 Conditional moments . . . 10
Brascamp–Lieb and Cramér–Rao bounds . . . 10
8.5 The gradient sector: Wetterich's equation . . . 11
Phase space and Lagrangian graphs . . . 11
No convexity is needed . . . 11
Probabilistic meaning of Γ . . . 13
8.6 Variational meaning and the tree level . . . 14
Constrained free energies . . . 14
The tree level . . . 14
8.7 Log-Sobolev inequalities along the localisation . . . 14
Entropy as quadratic variation . . . 14
In terms of the conditional covariance . . . 15
Exercises . . . 15

9 Variational structure . . . 16

9.1 Finite volume: the Boué–Dupuis formula . . . 16
Euler–Lagrange equation = FBSDE . . . 17
Entropy . . . 17
Generating functionals . . . 17
9.2 Infinite volume . . . 17
Differences of free energies . . . 18
Exercises . . . 19

Appendices: additional material . . . 19

Appendix A Algebraic structure of the Polchinski flow . . . 19

A.1 A semigroup of renormalisation maps . . . 19
A.2 Perturbation theory and counterterms . . . 20
Exercises . . . 21

Appendix B Dyson–Schwinger equations at every scale . . . 21

B.1 Gaussian bounds for the noise . . . 22
The roughness of the noise . . . 22
Control of the force . . . 23
Consequence for Φ^4_3 . . . 23
Exercises . . . 23

Appendix C Stochastic localisation . . . 24

The localisation as a measurement . . . 24
Why it is useful . . . 24
An application: the KLS conjecture . . . 25
The connection with Bauerschmidt–Bodineau . . . 25
Exercises . . . 25

References . . . 26

Introduction

In Lecture 3 we decomposed the Gaussian free field along a resolution parameter and took as primary objects the stochastic observables: martingales in the filtration of finite-resolution measurements [BCG26]. In this lecture we introduce interactions. The effective potentials solve the Polchinski equation, and the interacting field seen at increasing resolution solves a forward–backward stochastic differential equation whose force is itself an observable. See also [Pol84, BK87, Sal99, BBD24] for the renormalisation group side and [BG20, BG23] for the variational side.

Structure. Sections 7–9 are the core of the lecture: the Polchinski equation and its FBSDE, the FBSDE in force coordinates and the Wetterich equation, and the variational structure. Appendices A, B and C are additional material, referred to from the main text where needed.

What is formal. Rigorous statements require a finite-dimensional setting: a finite lattice, or a Fourier cutoff on the torus, together with a finite terminal resolution T. There, all the identities below are theorems under mild growth assumptions on the interaction, and their proofs are the computations we give. The hard problems are the limits $T\to\infty$ (ultraviolet) and infinite volume (infrared), together with the locality of the limiting objects. We mark with (formal) the statements that we only derive at this level.

Conventions. As in Lectures 1–3: $\mathcal{M}=\mathbb{T}_L^d$ or $\mathbb{R}^d$, $A=m^2-\Delta$, $C=A^{-1}$ with kernel G, μ_m the GFF, $\langle f,g\rangle=\int fg$ for real functions. For a functional F on smooth fields, $DF(\varphi)\in\mathcal{S}'(\mathcal{M})$ and $D^2F(\varphi)$ are its first and second derivatives, $D_hF=\langle DF,h\rangle$. $H_n(a;c)$ are the Hermite polynomials of variance c, and μ^Q is the centred Gaussian measure with covariance Q.

Main lecture

7 Interactions: the Polchinski equation and FBSDEs

7.1 Effective potentials

Fix a finite volume $\mathcal{M}=\mathbb{T}_L^d$ and a terminal resolution $T<\infty$, which plays the role of an ultraviolet cutoff: the reference field $X_T\sim\mu^{Q_T}$ is smooth. Let V_T be an interaction, a functional of smooth fields bounded below. Our main example is

$$V_T(\varphi)=\int_{\mathbb{T}_L^d}[\lambda H_4(\varphi(x);c_T)+\gamma_T H_2(\varphi(x);c_T)]\mathrm{d}x, \tag{7.1}$$

with a mass counterterm γ_T to be fixed in Appendix A.2 ($\gamma_T=0$ for $d=2$). The interacting measure at cutoff T is

$$\nu_T(\mathrm{d}\varphi)=Z_T^{-1}\mathrm{e}^{-V_T(\varphi)}\mu^{Q_T}(\mathrm{d}\varphi). \tag{7.2}$$

Definition 7.1. *The* effective potential *at resolution $t\in[0,T]$ is*

$$\mathrm{e}^{-V_t(\varphi)}:=\int\mathrm{e}^{-V_T(\varphi+\zeta)}\,\mu^{Q_{T,t}}(\mathrm{d}\zeta). \tag{7.3}$$

V_t is the interaction felt by the field at resolution t once the fluctuations between t and T have been integrated out: it is the Wilson effective action. At $t=0$, $V_0(0)=-\log Z_T$ is the free energy. Here is the first key fact.

Proposition 7.2. (Marginals) *Let $\mathbb{P}$ be defined by $\mathrm{d}\mathbb{P}/\mathrm{d}\mathbb{P}_0 = Z_T^{-1}\mathrm{e}^{-V_T(X_T)}$, where $\mathbb{P}_0$ is the law of the Gaussian process X. Then $\mathrm{Law}_{\mathbb{P}}(X_T) = \nu_T$, and for every $t \le T$*

$$\mathrm{Law}_{\mathbb{P}}(X_t) = \nu_t, \qquad \nu_t(\mathrm{d}\varphi) := Z_T^{-1}\mathrm{e}^{-V_t(\varphi)}\mu^{Q_t}(\mathrm{d}\varphi). \tag{7.4}$$

Moreover, the conditional law of X_T given $\mathcal{F}_t$ is

$$\mu_t^{X_t}(\mathrm{d}\varphi), \qquad \mu_t^{\phi}(G) := \mathrm{e}^{V_t(\phi)}\int G(\phi+\zeta)\,\mathrm{e}^{-V_T(\phi+\zeta)}\,\mu^{Q_{T,t}}(\mathrm{d}\zeta). \tag{7.5}$$

Proof. Under $\mathbb{P}_0$, $X_T = X_t + (X_T - X_t)$ with an independent increment of law $\mu^{Q_{T,t}}$. Hence $\mathbb{E}_{\mathbb{P}}[G(X_t)] = Z_T^{-1}\,\mathbb{E}_0[G(X_t)\mathrm{e}^{-V_T(X_T)}] = Z_T^{-1}\,\mathbb{E}_0[G(X_t)\mathrm{e}^{-V_t(X_t)}]$. The conditional law follows in the same way. □

So at every resolution the field is again a Gibbs measure, with reference covariance Q_t and potential V_t. The measure-valued martingale $(\mu_t^{X_t})_t$ goes from ν_T at $t=0$ to δ_{X_T} at $t=T$: it is a *stochastic localisation* of ν_T in the sense of Eldan [Eld23], adapted to the covariance profile [BBD24, BCG26].

Theorem 7.3. (Polchinski equation [Pol84]) (formal) *V_t solves the backward Hamilton–Jacobi equation*

$$\partial_t V_t = \tfrac{1}{2}\langle D V_t, \dot{Q}_t\, D V_t\rangle - \tfrac{1}{2}\mathrm{Tr}(\dot{Q}_t\, D^2 V_t), \qquad t\in[0,T], \tag{7.6}$$

with terminal condition V_T.

Proof. $u_t := \mathrm{e}^{-V_t} = \mathrm{e}^{\frac{1}{2}\Delta_{Q_{T,t}}}u_T$ is a Gaussian convolution, where $\Delta_Q := \mathrm{Tr}(Q\,D^2)$. Since $\partial_t Q_{T,t} = -\dot{Q}_t$, it solves the backward heat equation $\partial_t u_t = -\frac{1}{2}\mathrm{Tr}(\dot{Q}_t D^2 u_t)$. Insert $D^2 u = (D V \otimes D V - D^2 V)\,u$ and divide by $-u$. □

The nonlinear term generates new, generally nonlocal, interactions (trees), and the trace term contracts pairs of fields with the covariance of the shell $[t, t+\mathrm{d}t]$ (loops). The substitution $u = \mathrm{e}^{-V}$, which linearises the equation, is a Cole–Hopf transform: we come back to this in Section 7.4, and in Appendix A.

7.2 The Polchinski process

Theorem 7.4. (FBSDE for the interacting field) (formal) *Under $\mathbb{P}$ the process $\Phi_t := X_t$, the* interacting field *at resolution t, solves*

$$\mathrm{d}\Phi_t = \dot{Q}_t\,F_t\,\mathrm{d}t + \dot{Q}_t^{1/2}\,\mathrm{d}W_t^{\mathbb{P}}, \qquad F_t = -D V_t(\Phi_t) = \mathbb{E}_t^{\mathbb{P}}\left[-D V_T(\Phi_T)\right], \tag{7.7}$$

with $\Phi_0 = 0$ and a $\mathbb{P}$-cylindrical Brownian motion $W^{\mathbb{P}}$. In particular the effective force *$(F_t)_{t\le T}$ is a $\mathbb{P}$-martingale: an observable field of the interacting theory.*

Proof. The density process is $M_t := \mathbb{E}_0\left[\mathrm{d}\mathbb{P}/\mathrm{d}\mathbb{P}_0 | \mathcal{F}_t\right] = Z_T^{-1}\mathrm{e}^{-V_t(X_t)}$, by Proposition 7.2. By Itô's formula and (7.6), its drift vanishes and $\mathrm{d}M_t = -M_t\langle D V_t(X_t), \dot{Q}_t^{1/2}\mathrm{d}W_t\rangle$. By Girsanov's theorem, $W_t^{\mathbb{P}} = W_t + \int_0^t \dot{Q}_s^{1/2} D V_s(X_s)\mathrm{d}s$ is a $\mathbb{P}$-Brownian motion, which is (7.7). Differentiating (7.3) gives $D V_t(\phi) = \mu_t^{\phi}(D V_T)$, which is the second identity. Alternatively, Itô's formula for $D V_t(\Phi_t)$, together with the derivative of (7.6), shows that its drift vanishes. Here one uses that $D^2 V_t$ is symmetric. □

Read as an equation for an unknown pair (Φ, F), (7.7) is a *forward–backward* system [MY99]. The field is driven forward in resolution by a force that is determined backward, as the conditional expectation of a functional of the terminal field:

$$\Phi_t = \int_0^t \dot{Q}_s F_s \, \mathrm{d}s + \int_0^t \dot{Q}_s^{1/2} \mathrm{d}W_s, \qquad F_t = \mathbb{E}_t[-D V_T(\Phi_T)], \qquad t \in [0,T]. \tag{7.8}$$

Here and below we drop the superscript and write W for the Brownian motion driving the interacting field. The Polchinski equation is not needed to state (7.8), and $\nu_T = \mathrm{Law}(\Phi_T)$ is recovered from the solution. This is the starting point of FBSDE constructions of Euclidean fields (Figure 7.1): one solves (7.8) with renormalised V_T, with estimates uniform in T and in the volume, and takes the limit. The field coordinate Φ is *not* an observable, only a semimartingale. Its martingale part is the Gaussian noise, and the observables are built from it and from F.




Figure 7.1. The forward–backward structure of (7.8) on $\mathbb{T}^2$, heat profile, at resolutions $t_1 < t_2 < t_3 < T$, all built from the same noise. Top: the field gains the fluctuation of each shell, $\dot{Q}_t^{1/2}\mathrm{d}W_t$, and reveals finer modes. Bottom: the effective force $F_t = \mathbb{E}_t[-D V_T(\Phi_T)]$ for the Φ^4 interaction, determined backward from the terminal interaction. It is shown to first order in λ, where it is the Wick cube $-4\lambda H_3(\Phi_t; c_t)$, and each panel is in its own units.

Remark 7.5. (Closed versus local) At finite T the force is a closed observable, $F_t = \mathbb{E}_t[F_T]$. For Φ^4, $F_T(x) = -4\lambda H_3(\Phi_T(x); c_T) - 2\gamma_T H_1(\Phi_T(x); c_T)$ is local. As $T \to \infty$ one expects the force to converge as an observable. In $d = 2$ the limit is closed, $F_\infty = -4\lambda :\Phi^3_\infty:$. In $d = 3$ it is expected not to be. If it were, $F_t = \mathbb{E}_t[F_\infty]$ with $F_\infty \in L^1$, the DS equation of Appendix B would pass to the limit ($T \to \infty$, then $t \to \infty$), with F_∞ paired only with functionals of Φ_∞. Conditioning on $\sigma(\Phi_\infty)$ would then give a logarithmic derivative of the Φ^4_3 measure that is an integrable function of the field, $\langle Ah, \Phi_\infty \rangle - \mathbb{E}[F_\infty(h)|\Phi_\infty]$. This is not expected: the regularised logarithmic derivatives do not converge in L^2 [ALZ06]. The limiting force would then be an observable in the proper sense, a compatible family with no terminal representative. Locality of the limiting force is the crux of the construction [BCG26].




Figure 7.2. A zero-dimensional Polchinski process (Exercise 7.2): $Q_t = 1 - \mathrm{e}^{-t}$, $T = 5$, $V_T(x) = \frac{1}{2}x^4 - \frac{3}{2}x^2$. Left: the effective force $F_t = -V_t'$ is almost linear at low resolution and develops the double-well structure as $t \to T$. Centre: paths of the FBSDE (7.7), which split between the two wells (stochastic localisation). Right: histogram of Φ_T ($2 \cdot 10^4$ paths) against the density of ν_T.

7.3 Observables of the interacting theory

For a functional G of the terminal field, the observable $O_t = \mathbb{E}_t^{\mathbb{P}}[G(\Phi_T)]$ is $O_t = G_t(\Phi_t)$, with

$$G_t(\phi) = \mu_t^{\phi}(G) = \mathrm{e}^{V_t(\phi)} \int G(\phi + \zeta)\, \mathrm{e}^{-V_T(\phi+\zeta)} \mu^{Q_{T,t}}(\mathrm{d}\zeta). \tag{7.9}$$

Formally, G_t solves the backward Kolmogorov equation of the interacting dynamics,

$$\partial_t G_t + \tfrac{1}{2}\mathrm{Tr}(\dot{Q}_t D^2 G_t) - \langle \dot{Q}_t D V_t, D G_t \rangle = 0, \qquad G_T = G, \tag{7.10}$$

which is the Doob transform of the Gaussian heat equation by e^{-V_t} (Exercise 7.3). Conversely, any solution of (7.10) yields an observable $G_t(\Phi_t)$; at finite T it is closed, but as $T \to \infty$ it need not be. The force is the case $G = -D V_T$, and is special because it also drives the dynamics.

Tangent observables. Let V_T^{θ} depend on a coupling θ, and let $U_t := \partial_\theta V_t^{\theta}$. Differentiating (7.3) gives $U_t(\Phi_t) = \mathbb{E}_t^{\theta}[\partial_\theta V_T^{\theta}(\Phi_T)]$, an observable, and

$$\partial_\theta \mathbb{E}^{\nu_T^{\theta}}[G] = -\mathrm{Cov}^{\nu_T^{\theta}}(G, \partial_\theta V_T^{\theta}). \tag{7.11}$$

Thus the tangent observables, such as the running $:\varphi^4:$ for $\theta = \lambda$ (together with the derivative of the counterterm $\partial_\lambda \gamma_T \int :\varphi^2:$), describe the response of the theory to a change of coupling. They are the observables conjugate to the coordinates of theory space.

7.4 The force and a Burgers equation

Differentiating (7.6), the effective force $F_t = -D V_t$, now as a function of the field, solves

$$\partial_t F_t = -D F_t[\dot{Q}_t F_t] - \tfrac{1}{2}\mathrm{Tr}(\dot{Q}_t D^2 F_t), \qquad F_T = -D V_T, \tag{7.12}$$

an infinite-dimensional, backward, viscous *Burgers equation*. The Cole–Hopf transform $F = D \log u$ linearises it, which is the statement that $u = \mathrm{e}^{-V}$ solves a heat equation. Two remarks on (7.12):

i. it uses only the force, which is a local quantity (*intensive*), while V_t is *extensive*, of order of the volume. This matters in infinite volume (Section 9.2);

ii. the Burgers flows for different covariances commute, and the curvature vanishes, when DF_t is symmetric, i.e. when F_t is a gradient [BCG26]. For a non-gradient force, as for the general Wilson–Itô diffusions of [BCG26], the flows do not commute.

Exercises

Exercise 7.1. (Quadratic interactions) Let $V_T(\varphi)=\frac{\kappa}{2}\langle\varphi,\varphi\rangle$, $\kappa>0$. (a) Show that $V_t(\varphi)=\frac{1}{2}\langle\varphi,\kappa(1+\kappa Q_{T,t})^{-1}\varphi\rangle+$ const solves (7.6). (b) Show that ν_T is Gaussian with covariance $(Q_T^{-1}+\kappa)^{-1}$: for $Q_T\to(m^2-\Delta)^{-1}$ this is the GFF of mass $(m^2+\kappa)^{1/2}$. (c) Write (7.7) explicitly. It is a linear FBSDE. Solve it, and check that F_t is a martingale. *Moral:* the mass can be put either into the covariance or into the force.

Exercise 7.2. (Zero dimensions) Take $\mathcal{M}$ a point, $Q_t=q(t)$ scalar, increasing, $V_T(x)=\lambda x^4-ax^2$ with $\lambda>0$, $a\geq 0$ (a double well if $a>0$; Figure 7.2 has $\lambda=\frac{1}{2}$, $a=\frac{3}{2}$). (a) Write (7.6) and solve it numerically for $q(t)=1-\mathrm{e}^{-t}$. (b) Simulate (7.7) by first computing V_t' on a grid, and compare the histogram of Φ_T with $\mathrm{e}^{-\lambda x^4+ax^2-x^2/2q(T)}$. (c) Compute V_t to second order in λ.

Exercise 7.3. (Doob transform) Show that G_t in (7.9) solves (7.10), by writing $\mathrm{e}^{-V_t}G_t=\mathrm{e}^{\frac{1}{2}\Delta_{Q_{T,t}}}(\mathrm{e}^{-V_T}G)$. Prove (7.11).

8 The FBSDE in force coordinates: the Wetterich equation

In this section the forward–backward system (7.8) is the primary object. We look for the coordinates in which it is simplest. They remove the predictable part of the forward dynamics, and they integrate exactly the bilinear (tree) part of the force flow (7.12). In the gradient case the resulting equation for the force is the Wetterich equation for the effective average action [Wet93, BTW02]. Throughout,

$$S_t:=Q_T-Q_t=Q_{T,t},\qquad \dot S_t=-\dot Q_t,\qquad S_T=0,$$

is the covariance still unresolved at resolution t, and (Φ,F) solves (7.8) with a martingale force $F_t=F_t(\Phi_t)$, $\mathrm{d}F_t=\Upsilon_t\,\mathrm{d}W_t$.

8.1 State and costate

Pontryagin's maximum principle. Consider first a deterministic problem: minimise the cost $J(u)=g(x_T)+\int_t^T\ell(x_s,u_s)\,\mathrm{d}s$ over controls u on $[t,T]$, where the state solves $\dot x_s=b(x_s,u_s)$, $x_t=x$. With the *Hamiltonian*

$$H(x,p,u):=\langle p,b(x,u)\rangle-\ell(x,u),$$

Pontryagin's principle [PBGM62] says that along an optimal pair (x,u) there is a *costate* p such that

$$\dot x_s=\partial_pH,\qquad \dot p_s=-\partial_xH,\qquad p_T=-Dg(x_T),\qquad u_s\in\arg\max_v H(x_s,p_s,v).$$

The state runs forward from its initial value, the costate backward from its terminal value: a two-point boundary value problem. In the smooth case the principle is the first-order condition. If $\xi=\partial_\varepsilon x$ is the variation of the state under $u\to u+\varepsilon v$, then $\dot\xi=\partial_x b\,\xi+\partial_u b\,v$, $\xi_t=0$, and the costate equation is exactly what makes $\frac{\mathrm{d}}{\mathrm{d}s}\langle p,\xi\rangle=\langle\partial_x\ell,\xi\rangle+\langle p,\partial_u b\,v\rangle$. Integrating over $[t,T]$ and using the terminal condition,

$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}J(u+\varepsilon v)|_{\varepsilon=0}=-\int_t^T\langle\partial_uH(x_s,p_s,u_s),v_s\rangle\,\mathrm{d}s.$$

So the costate is the Lagrange multiplier of the constraint $\dot x=b$, and an optimal control is a critical point of the Hamiltonian.

The costate is a gradient. Let $V(t,x) = \min_u J(u)$ be the value function. Comparing the cost from (t,x) with the cost from $(t+\mathrm{d}t, x+b\,\mathrm{d}t)$ gives the Hamilton–Jacobi–Bellman equation

$$\partial_t V = H_*(x, -DV), \qquad H_*(x,p) := \max_u H(x,p,u), \qquad V(T,\cdot) = g.$$

Its characteristics are Hamilton's equations for H_*, and along them $p_s = -DV(s,x_s)$: the costate is the sensitivity of the optimal cost to a displacement of the state, i.e. the force that the future exerts on the present.

The Hamiltonian of the field. In the problem of Theorem 9.1 the state is the field, $b(x,u) = \dot{Q}_s^{1/2}u$, $\ell = \frac{1}{2}\|u\|^2$ and $g = V_T$. Then $H(p,u) = \langle p, \dot{Q}_s^{1/2}u\rangle - \frac{1}{2}\|u\|^2$ is maximal at $u = \dot{Q}_s^{1/2}p$, and

$$H_*(p) = \frac{1}{2}\langle p, \dot{Q}_s p\rangle$$

is the Hamiltonian of a free particle with mass matrix $\dot{Q}_s^{-1}$: it does not depend on the state. Hence $\dot{p} = 0$ and $\dot{x} = \dot{Q}_s p$: the costate is conserved, and from t to T the state moves along the linear shear

$$\Sigma_t \colon (x,p) \longmapsto (x + S_t p,\ p). \tag{8.1}$$

Its trajectories are the characteristics of the transport part of (7.12). All the interaction sits in the terminal condition $p = -DV_T(x + S_t p)$, an implicit equation for the momentum, solved by a sum of tree graphs (Exercise 8.2). The corresponding Hamilton–Jacobi equation $\partial_t V = \frac{1}{2}\langle DV, \dot{Q}_t DV\rangle$ is the Polchinski equation (7.6) without its trace term.

With noise. For a controlled diffusion $\mathrm{d}X = b(X,u)\,\mathrm{d}s + \sigma\,\mathrm{d}W$, with σ independent of the control, the stochastic maximum principle [Bis73, Pen90, YZ99] keeps the Hamiltonian and the maximisation, but the costate becomes an adapted process which solves a *backward* stochastic differential equation,

$$\mathrm{d}p_s = -\partial_x H\ \mathrm{d}s + \Upsilon_s\,\mathrm{d}W_s, \qquad p_T = -Dg(X_T).$$

The martingale term $\Upsilon\,\mathrm{d}W$ is the price of adaptedness: the costate may not anticipate the noise. For the field $\partial_x H = 0$, so the costate is a martingale, $p_t = \mathbb{E}_t[-DV_T(X_T)]$, and since the value function is the effective potential (Theorem 9.1), $p_t = -DV_t(X_t)$. The noise adds the Itô term to the Hamilton–Jacobi equation, which becomes the Polchinski equation (7.6).

In the notation of Theorem 9.1 the controlled field is $\mathrm{d}X = \dot{Q}_t^{1/2}(u\,\mathrm{d}t + \mathrm{d}W)$ and the optimal control is $u_t = \dot{Q}_t^{1/2}p_t$ (Exercise 8.1). Since p_t is the value F_t of the force, the FBSDE (7.8) is the stochastic Pontryagin system of this problem:

the field Φ is the state, and the force F is the costate.

The function $F_t(\cdot)$ with $F_t = F_t(\Phi_t)$ is the *decoupling field* of the FBSDE, and the Burgers equation (7.12) is its quasilinear PDE; $\Upsilon_t = DF_t(\Phi_t)\,\dot{Q}_t^{1/2}$ is the martingale representation integrand of the costate. The two terms of (7.12) have distinct origins. The transport term $-DF[\dot{Q}F]$ comes from the drift of the state, and generates the tree diagrams. The diffusion term $-\frac{1}{2}\mathrm{Tr}(\dot{Q}D^2F)$ is the Itô correction, and generates the loops.

8.2 Removing the predictable drift

The drift $\dot{Q}_t F_t$ of the state is predictable, and the costate is a martingale. Hence the drift that remains to be accumulated on $[t,T]$ is predicted, at resolution t, by transporting the current state with the current costate along the free flow (8.1).

Proposition 8.1. (The forecast) (formal) *The conditional mean of the terminal field is*

$$\psi_t := \mathbb{E}_t[\Phi_T] = \Phi_t + S_t F_t, \tag{8.2}$$

i.e. $(\psi_t, F_t) = \Sigma_t(\Phi_t, F_t)$. *It is a martingale, with*

$$\mathrm{d}\psi_t = \dot{Q}_t^{1/2}\mathrm{d}W_t + S_t \Upsilon_t \,\mathrm{d}W_t = (1 + S_t D F_t(\Phi_t))\, \dot{Q}_t^{1/2}\mathrm{d}W_t. \tag{8.3}$$

Proof. Integrate the forward equation from t to T and condition on $\mathcal{F}_t$: since F is a martingale, $\mathbb{E}_t \int_t^T \dot{Q}_s F_s \mathrm{d}s = S_t F_t$. For the differential, the drift $\dot{Q}_t F_t$ of Φ cancels against $\dot{S}_t F_t$, and $S_t \,\mathrm{d}F_t = S_t \Upsilon_t \,\mathrm{d}W_t$. □

So ψ is a closed observable, the *field observable* of [BCG26]. The field splits as

$$\Phi_t = \psi_t - S_t F_t: \tag{8.4}$$

a martingale, minus the drift still expected on the unresolved scales, which vanishes at $t = T$. The noise of ψ in (8.3) has two sources: the fresh fluctuation of the shell, $\dot{Q}_t^{1/2}\mathrm{d}W_t$, and the revision of the forecast of the remaining drift, $S_t \,\mathrm{d}F_t$. The coupling between the forward and the backward equations, which was in the drift of Φ, now sits in the volatility of ψ. The coordinate ψ is the *interaction picture* with respect to the noiseless (classical) part of the dynamics.

8.3 The FBSDE in the forecast coordinate

Assume that the map $x \mapsto \psi = x + S_t F_t(x)$ is invertible, and write its inverse as

$$x = \psi + S_t \mathcal{F}_t(\psi), \qquad \mathcal{F}_t(\psi) := -F_t(x), \tag{8.5}$$

so that $F_t = -\mathcal{F}_t(\psi_t)$: the costate is now a function of the forecast. Introduce the *response* and the *regulator*

$$H_t(\psi) := (S_t^{-1} + D\,\mathcal{F}_t(\psi))^{-1}, \qquad \mathcal{R}_t := S_t^{-1} - Q_T^{-1}, \qquad \dot{\mathcal{R}}_t = S_t^{-1}\dot{Q}_t S_t^{-1} \geq 0. \tag{8.6}$$

Differentiating the two maps $x \leftrightarrow \psi$ gives $1 + S_t D F_t(x) = (1 + S_t D\,\mathcal{F}_t(\psi))^{-1} = H_t(\psi)\, S_t^{-1}$.

Theorem 8.2. (FBSDE in force coordinates) (formal) *In the coordinates* $(\psi, \mathcal{F})$, *the system* (7.8) *becomes*

$$\mathrm{d}\psi_t = H_t(\psi_t)\; S_t^{-1}\dot{Q}_t^{1/2}\,\mathrm{d}W_t, \qquad \mathcal{F}_t(\psi_t) = \mathbb{E}_t[D V_T(\psi_T)], \tag{8.7}$$

with $\psi_0 = S_0 F_0 = \mathbb{E}[\Phi_T]$ *and* $\psi_T = \Phi_T$. *The forward equation has no drift, and its covariance is* $C_t := H_t \dot{\mathcal{R}}_t H_t^*$. *The decoupling field* $\mathcal{F}$ *solves the pure diffusion equation*

$$\partial_t \mathcal{F}_t + \frac{1}{2}\mathrm{Tr}(C_t\, D^2 \mathcal{F}_t) = 0, \qquad \mathcal{F}_T = D V_T. \tag{8.8}$$

Proof. The forward equation is (8.3) with $1 + S_t D F_t = H_t S_t^{-1}$, and its covariance is

$$H_t S_t^{-1}\dot{Q}_t S_t^{-1} H_t^* = H_t \dot{\mathcal{R}}_t H_t^*.$$

The backward equation is $F_t = \mathbb{E}_t[F_T]$ with $\psi_T = \Phi_T$. Since $\mathcal{F}_t(\psi_t)$ is a martingale and ψ has no drift, the drift in Itô's formula for $\mathcal{F}_t(\psi_t)$ is the left-hand side of (8.8). □

Compare (8.8) with the Burgers equation (7.12): the transport term has been absorbed exactly by the change of coordinates, and only the Itô term is left. If it is dropped, $\mathcal{F}_t = D V_T$ does not move: all the tree diagrams are contained in the coordinate change (8.5). The loops are generated only by the quadratic variation C_t of the forecast, i.e. by the randomness of the noise and of the force revisions, and the nonlinearity has moved into the field dependence of C_t. None of this uses a gradient structure or a Gaussian convolution: Theorem 8.2 holds for any FBSDE of the form (7.8) with a Markov martingale force, gradient or not, for which the coordinate change is invertible. For a general terminal force $F_T = \mathfrak{f}(\Phi_T)$, replace $D V_T$ by $-\mathfrak{f}$ in (8.7) and (8.8).

8.4 Conditional moments

For the Polchinski process the response has a direct meaning.

Proposition 8.3. (formal) *If $F_t = -D V_t$ then*

$$H_t(\psi_t) = \mathrm{Cov}_t := \mathrm{Cov}^{\mathbb{P}}(\Phi_T \mid \Phi_t) = S_t - S_t D^2 V_t(\Phi_t)\, S_t.$$

In particular $H_t > 0$ is symmetric, $1 + S_t D F_t = \mathrm{Cov}_t S_t^{-1}$ is always invertible, and $H_t \le S_t$ when V_T is convex.

Proof. By (7.5), the law of Φ_T given $\mathcal{F}_t$ depends only on Φ_t, so it is also the law given Φ_t. The conditional law μ_t^{ϕ} is the tilt of $\mu^{S_t}(\cdot - \phi)$ by e^{-V_T}; differentiating twice in ϕ gives $\mathrm{Cov}(\mu_t^{\phi}) = S_t - S_t D^2 V_t(\phi)\, S_t$. On the other hand $H_t S_t^{-1} = 1 + S_t D F_t = 1 - S_t D^2 V_t$. The last claim is the Brascamp–Lieb inequality. □

So the forecast coordinates are the first two conditional moments of the stochastic localisation $\mu_t^{\Phi_t}$ of Proposition 7.2: $(\psi_t, \mathrm{Cov}_t) = (\mathbb{E}[\Phi_T \mid \Phi_t], \mathrm{Cov}(\Phi_T \mid \Phi_t))$ (Figure 8.1). They are tied by Itô's formula: since $\mathbb{E}_t[\Phi_T \otimes \Phi_T] = \psi_t \otimes \psi_t + \mathrm{Cov}_t$ is a martingale,

$$\mathrm{d}\langle \psi, \psi \rangle_t = C_t\, \mathrm{d}t = \mathrm{Cov}_t \dot{\mathcal{R}}_t \mathrm{Cov}_t\, \mathrm{d}t, \qquad \mathrm{dCov}_t = -\mathrm{Cov}_t \dot{\mathcal{R}}_t \mathrm{Cov}_t\, \mathrm{d}t + \mathrm{d}(\text{martingale}). \tag{8.9}$$

Conditional variance is resolved at the rate $\mathrm{Cov}_t \dot{\mathcal{R}}_t \mathrm{Cov}_t$, which for the Gaussian reference is $S_t \dot{\mathcal{R}}_t S_t = \dot{Q}_t$.

Brascamp–Lieb and Cramér–Rao bounds. The conditional law μ_t^{ϕ} has density proportional to e^{-U}, with $U(y) = V_T(y) + \frac{1}{2}\langle y - \phi, S_t^{-1}(y - \phi)\rangle$ and $D^2 U = S_t^{-1} + D^2 V_T$. Two classical inequalities bound its covariance by the Hessian of U. If $D^2 U > 0$, the Brascamp–Lieb inequality [BL76], $\mathrm{Var}_{\mu}(f) \le \mathbb{E}_{\mu}\langle Df, (D^2 U)^{-1} Df\rangle$, applied to linear functions $f = \langle h, \cdot\rangle$, and the Cramér–Rao inequality, which needs no convexity, give respectively the upper and the lower bound in

$$(\mathbb{E}_t[S_t^{-1} + D^2 V_T(\Phi_T)])^{-1} \le \mathrm{Cov}_t \le \mathbb{E}_t[(S_t^{-1} + D^2 V_T(\Phi_T))^{-1}].$$

The lower bound follows from $\mathbb{E}_{\mu}[(y - m) \otimes D U(y)] = 1$ (integration by parts) and the Cauchy–Schwarz inequality, with the Fisher information $\mathbb{E}_{\mu}[D U \otimes D U] = \mathbb{E}_{\mu} D^2 U$ (Exercise 8.10). The two bounds coincide when V_T is quadratic, and their gap is a Jensen gap for the inverse. Through Proposition 8.3 they are two-sided bounds on the Hessian of the effective potential, i.e. on the response H_t, which is the propagator of the Wetterich equation below. Some consequences:

- If V_T is convex, then $\mathrm{Cov}_t \le S_t$: the interaction can only reduce the conditional variance. Equivalently $D^2 V_t \ge 0$: convexity is preserved along the Polchinski flow (as also follows from the Prékopa–Leindler inequality), and $D^2 \Gamma_t \ge 0$ in (8.10).
- If $D^2 V_T \ge \kappa$ with $\kappa > 0$, then $\mathrm{Cov}_t \le (S_t^{-1} + \kappa)^{-1}$ uniformly in the conditioning: a uniform mass in the interaction is a uniform mass in every conditional law.
- For Φ_3^4 convexity is already lost through the Wick ordering: $H_4(\varphi; c_T) = \varphi^4 - 6 c_T \varphi^2 + 3 c_T^2$, so (7.1) has curvature $-12 \lambda c_T + 2 \gamma_T$ at $\varphi = 0$, which diverges as $T \to \infty$ (this happens already for Φ_2^4). The upper bound is lost, and at low resolution the conditional law may spread over several wells (Figure 8.1). This is the large-field problem. In Lecture 5 it is handled by a clipped approximate action, whose global coercivity (Lemma 12.1 there) is a bound of Brascamp–Lieb type on the response.




Figure 8.1. Stochastic localisation for a two-component field: $Q_t = (1-\mathrm{e}^{-t})\,1$, $T=4$, V_T a tilted double well, and the contours show the density of ν_T. One path of (7.7) up to resolution t (navy), with its terminal value Φ_T (open circle). The forecast $\psi_t = \Phi_t + S_t F_t$ (gold) is the mean of the conditional law $\mu_t^{\Phi_t}$: teal samples, and the ellipse is two standard deviations of Cov_t. At low resolution the conditional law covers both wells; as t grows it localises on one of them.

8.5 The gradient sector: Wetterich's equation

Phase space and Lagrangian graphs. Pairs (x,p) of a field and a value of the force form the phase space $\mathbb{R}^n \times \mathbb{R}^n$, with the symplectic form $\omega = \sum_i \mathrm{d}x_i \wedge \mathrm{d}p_i$. A submanifold L of dimension n is *Lagrangian* if ω vanishes on it. The graph $L = \{(x, F(x))\}$ of a force field is Lagrangian if and only if $\sum_{i,j} \partial_j F_i \,\mathrm{d}x_i \wedge \mathrm{d}x_j = 0$, i.e. DF is symmetric, i.e. F is (locally) a gradient: the gradient sector is the sector of Lagrangian graphs. A map that preserves ω (a *symplectic* map) sends Lagrangian submanifolds to Lagrangian submanifolds. The shear (8.1) is symplectic, since $\Sigma_t^* \omega = \omega + \sum_{i,j} (S_t)_{ij} \mathrm{d}p_j \wedge \mathrm{d}p_i = \omega$ for S_t symmetric. So it maps the graph of a gradient to a Lagrangian submanifold; if the image is again a graph over the new base variable, it is the graph of a gradient.

Now let $F_t = -D V_t$. Its graph is Lagrangian, hence so is its image under the shear. The image is a graph over ψ: the map $x \mapsto x + S_t F_t(x)$ is S_t times the gradient of $\frac{1}{2}\langle x, S_t^{-1} x\rangle - V_t(x)$, which is strictly convex since its Hessian is $S_t^{-1}\mathrm{Cov}_t S_t^{-1} > 0$ (Proposition 8.3), so the map is injective. Hence the image is the graph of a gradient:

$$\mathcal{F}_t = D\Gamma_t, \qquad H_t = (S_t^{-1} + D^2\Gamma_t)^{-1}, \qquad \Gamma_T = V_T. \tag{8.10}$$

The generating function of this canonical transformation is the Moreau (infimal convolution) transform

$$V_t(x) = \inf_{\psi} \left\{ \Gamma_t(\psi) + \tfrac{1}{2}\langle x - \psi, S_t^{-1}(x-\psi)\rangle \right\}, \tag{8.11}$$

whose optimality conditions $x = \psi + S_t D\Gamma_t(\psi)$, $D V_t(x) = D\Gamma_t(\psi)$ are (8.5). It is well defined by the same convexity: up to the change of variable $j = S_t^{-1} x$, it is a Legendre transform. We use (8.11) to fix the additive constant in Γ_t. Figure 8.2 shows the construction in zero dimensions.

No convexity is needed. The construction uses no convexity of V_T, V_t or Γ_t. The convex function is $g_t(x) = \frac{1}{2}\langle x, S_t^{-1} x\rangle - V_t(x)$: with $j = S_t^{-1} x$ it is, up to a constant, the log-Laplace transform of the finite measure $\mathrm{e}^{-V_T(y) - \frac{1}{2}\langle y, S_t^{-1} y\rangle}\,\mathrm{d}y$, for every V_T bounded below. Its gradient in j is the conditional mean ψ, and it is strictly convex because $\mathrm{Cov}_t > 0$. In finite dimensions, when V_T is finite, this exponential family is regular with full support, so $j \mapsto \psi$ is a bijection of $\mathbb{R}^n$: the image of the shear is a graph over the whole ψ space, also when V_t is a double well.

For Γ_t one only gets semiconvexity, $S_t^{-1}+D^2\Gamma_t=\mathrm{Cov}_t^{-1}>0$, i.e. $\Gamma_t+\frac{1}{2}\langle\cdot,S_t^{-1}\cdot\rangle$ is strictly convex, which is what (8.11) requires. At $t=0$, where $S_0=Q_T$, this semiconvexity is the convexity of Γ_0^{EAA}. Γ_t itself is convex if and only if $\mathrm{Cov}_t\le S_t$, which holds for convex V_T but may fail for a double well at low resolution. Convexity matters only without noise: the Hopf–Lax potential (8.16) is merely semiconcave, and unless $S_t^{-1}+D^2V_T>0$ the characteristics cross, producing caustics and shocks of the force. The noise removes them.




Figure 8.2. The zero-dimensional model of Figure 7.2 at $t=1.2$. Left: the shear Σ_t (teal arrows) maps the graph of the force $F_t=-V_t'$ (navy) to a graph over the forecast ψ (gold), which is the graph of $-\Gamma_t'$. Right: the bare potential V_T, the effective potential V_t and Γ_t, each shifted to have minimum 0.

Theorem 8.4. (Wetterich equation [Wet93]) (formal) Γ_t *solves*

$$\partial_t\Gamma_t=\tfrac{1}{2}\mathrm{Tr}[\dot{\mathcal{R}}_t((S_t^{-1}+D^2\Gamma_t)^{-1}-S_t)],\qquad \Gamma_T=V_T. \tag{8.12}$$

Equivalently, the effective average action $\Gamma_t^{\mathrm{EAA}}(\psi):=\Gamma_t(\psi)+\frac{1}{2}\langle\psi,Q_T^{-1}\psi\rangle$ *solves, up to a field-independent term,*

$$\partial_t\Gamma_t^{\mathrm{EAA}}=\tfrac{1}{2}\mathrm{Tr}[\dot{\mathcal{R}}_t(D^2\Gamma_t^{\mathrm{EAA}}+\mathcal{R}_t)^{-1}]. \tag{8.13}$$

It interpolates between the bare action $\Gamma_T^{\mathrm{EAA}}=\frac{1}{2}\langle\psi,Q_T^{-1}\psi\rangle+V_T(\psi)$ ($\mathcal{R}_T=\infty$) *and the 1PI effective action* Γ_0^{EAA} *of* ν_T ($\mathcal{R}_0=0$).

Proof. Integrate (8.8) in field space. With $\mathcal{F}=D\Gamma$, the derivative of $H=(S^{-1}+D^2\Gamma)^{-1}$ in a direction v is $-H\,D^3\Gamma[v]\,H$, so by the total symmetry of $D^3\Gamma$,

$$D_v\mathrm{Tr}(\dot{\mathcal{R}}_tH_t)=-\mathrm{Tr}(H_t\dot{\mathcal{R}}_tH_t\,D^3\Gamma_t[v])=-\mathrm{Tr}(C_t\,D^2\mathcal{F}_t)[v].$$

Hence (8.8) reads $D\left(\partial_t\Gamma_t-\frac{1}{2}\mathrm{Tr}(\dot{\mathcal{R}}_tH_t)\right)=0$, and (8.12) holds up to a function of t alone. That this function is $-\frac{1}{2}\mathrm{Tr}(\dot{\mathcal{R}}_tS_t)$ with the normalisation (8.11) follows from the Polchinski equation by the envelope theorem (Exercise 8.5). For (8.13), note that $D^2\Gamma_t^{\mathrm{EAA}}+\mathcal{R}_t=D^2\Gamma_t+S_t^{-1}$. For the endpoints: $S_T=0$; and at $t=0$, $S_0=Q_T$, so (8.11) identifies $\frac{1}{2}\langle x,Q_T^{-1}x\rangle-V_0(x)$, up to a constant, with the cumulant generating function of ν_T at $Q_T^{-1}x$, whose Legendre transform is Γ_0^{EAA} (Exercise 8.7). □

Thus the Wetterich equation is the gradient-integrated form of the force equation (8.8), i.e. of the backward half of the FBSDE in forecast coordinates [Mor94, BCG26]. Every object in it is an observable of the Polchinski process:

Wetterich	FBSDE
field argument ψ	forecast $\psi_t = \mathbb{E}_t[\Phi_T]$, the state transported by Σ_t
$D\Gamma_t(\psi_t)$	minus the costate, $-F_t = \mathbb{E}_t[D V_T(\Phi_T)]$
propagator $(D^2\Gamma_t^{\mathrm{EAA}} + \mathcal{R}_t)^{-1}$	conditional covariance Cov_t
regulator $\mathcal{R}_t = S_t^{-1} - Q_T^{-1}$	precision of the unresolved shell, minus that of the prior
$\partial_t \Gamma_t$	$\frac{1}{2}\mathrm{Tr}[\dot{\mathcal{R}}_t(\mathrm{Cov}_t - S_t)]$

The flow of Γ is driven by the difference between the interacting and the free conditional covariances, paired with the shell $\dot{\mathcal{R}}_t$. It is *one-loop exact*, with the full propagator inside the trace. In the Gaussian case $D^2\Gamma_t$ is constant and only the constant term of Γ flows. For the heat profile with $T = \infty$, $S_t = C(1 - \mathrm{e}^{-A/t^2})$ and

$$\mathcal{R}_t = \frac{A}{\mathrm{e}^{A/t^2} - 1}, \tag{8.14}$$

which behaves like a mass term t^2 on the modes $|k| \ll t$ (for $m \ll t$) and is exponentially small on the modes $|k| \gg t$: the exponential regulator of the functional renormalisation group (Figure 8.3).

Probabilistic meaning of Γ. At $t = 0$ the effective average action is the 1PI effective action of ν_T,

$$\Gamma_0^{\mathrm{EAA}}(\psi) = V_0(0) + \sup_j \{\langle j, \psi\rangle - \Lambda(j)\}, \qquad \Lambda(j) = \log \int \mathrm{e}^{\langle j, \varphi\rangle} \nu_T(\mathrm{d}\varphi),$$

the generating functional of the vertex functions introduced by Jona-Lasinio [JL64]. It is a probabilistic object, as Jona-Lasinio emphasised in his probabilistic reading of the renormalisation group [JL75, JL01]; for the link with large deviations see Section VIII of [JL01]. By Cramér's theorem the empirical mean of N independent samples of ν_T takes the value ψ with probability $\approx \exp\left(-N\left(\Gamma_0^{\mathrm{EAA}}(\psi) - V_0(0)\right)\right)$, so $\Gamma_0^{\mathrm{EAA}} - V_0(0)$ is the rate function of the large deviations of averages of the field. In particular Γ_0^{EAA} is convex even when V_T is not, it is minimal at the mean of ν_T, and its Hessian there is the inverse covariance. Formally the same holds for the field averaged over a large volume, with the volume as the large parameter and the effective potential of quantum field theory (Γ per unit volume at constant ψ, not V_t) as rate function; in that limit flat parts may appear (the Maxwell construction).




Figure 8.3. The regulator $\mathcal{R}_t$ of the heat profile ($T = \infty$, $m = 1$): close to the mass term t^2 (dotted) on the modes $|k| \ll t$, exponentially small on the modes $|k| \gg t$.

8.6 Variational meaning and the tree level

Constrained free energies. (formal) By the Gibbs variational principle applied to the conditional law (7.5), and the identity $\mathrm{Ent}(\nu|\mu^S(\cdot-x)) = \mathrm{Ent}(\nu|\mu^S(\cdot-\psi)) + \frac{1}{2}\langle x-\psi, S^{-1}(x-\psi)\rangle$ for ν of mean ψ,

$$\Gamma_t(\psi) = \inf\Big\{\int V_T\,\mathrm{d}\nu + \mathrm{Ent}(\nu\,|\,\mu^{S_t}(\cdot-\psi)) \colon \textstyle\int \varphi\;\nu(\mathrm{d}\varphi) = \psi\Big\}. \tag{8.15}$$

So Γ_t, the effective average action up to the quadratic term $\frac{1}{2}\langle\psi, Q_T^{-1}\psi\rangle$, is the free energy of the unresolved shell constrained to the forecast ψ, and (8.11) is the minimisation over the forecast. In the language of Theorem 9.1: V_t is the value function of the control problem, and Γ_t is its Moreau transform, i.e. the value function expressed in the forecast coordinate, the interaction picture of the classical part of the dynamics.

The tree level. Dropping the Itô term in (8.8) gives $\Gamma_t = V_T$, and then (8.11) becomes the Hopf–Lax formula

$$V_t^{\mathrm{tree}}(x) = \inf_y\Big\{V_T(y) + \tfrac{1}{2}\langle y-x, S_t^{-1}(y-x)\rangle\Big\}, \tag{8.16}$$

the solution of the noiseless Polchinski equation $\partial_t V = \frac{1}{2}\langle DV, \dot{Q}\,DV\rangle$. It is the saddle point of the Gaussian convolution (7.3). Its minimiser solves the classical field equation $S_t^{-1}(y-x) + DV_T(y) = 0$ in the shell $[t,T]$ with background x, and expanding it in V_T produces exactly the tree diagrams.

8.7 Log-Sobolev inequalities along the localisation

Bauerschmidt and Bodineau [BB21] proved log-Sobolev inequalities by running the Polchinski flow (see the survey [BBD24]). In their language the renormalised measure at a given scale is the law of Φ_t, and the fluctuation measure is the conditional law $\mu_t^{\Phi_t}$ of Proposition 7.2, i.e. the stochastic localisation of ν_T [Eld23]. (Their scale parameter runs opposite to our resolution: their initial scale is our T, where the renormalised measure is ν_T and the fluctuation measure is trivial, and their integrated covariance is our S_t.) In this dictionary their argument is an entropy computation along our filtration; Appendix C develops the comparison.

Entropy as quadratic variation. Let $\rho \geq 0$ with $\mathbb{E}_{\nu_T}\rho = 1$, and let $M_t = \mathbb{E}_t[\rho(\Phi_T)] = G_t(\Phi_t)$, with $G_t(\phi) = \int \rho\,\mathrm{d}\mu_t^{\phi}$. The martingale M runs from $M_0 = 1$ to $M_T = \rho(\Phi_T)$, and its noise is that of the field: $\mathrm{d}M_t = \langle DG_t(\Phi_t), \dot{Q}_t^{1/2}\,\mathrm{d}W_t\rangle$. Itô's formula for $M\log M$ gives

$$\mathrm{Ent}_{\nu_T}(\rho) = \frac{1}{2}\,\mathbb{E}\int_0^T \frac{\mathrm{d}\langle M\rangle_t}{M_t} = 2\,\mathbb{E}\int_0^T \|\dot{Q}_t^{1/2} D\sqrt{G_t}(\Phi_t)\|^2\,\mathrm{d}t,$$

and, stopping at resolution t, the chain rule

$$\mathrm{Ent}_{\nu_T}(\rho) = \mathrm{Ent}(G_t(\Phi_t)) + \mathbb{E}\big[\mathrm{Ent}_{\mu_t^{\Phi_t}}(\rho)\big]\colon$$

the entropy already revealed at resolution t plus the expected entropy still hidden. The increments are controlled by the conditional law: differentiating the density of μ_t^{ϕ} in ϕ, $DG_t(\phi) = S_t^{-1}\mathrm{Cov}_{\mu_t^{\phi}}(\rho, y)$.

Theorem 8.5. (Multiscale Bakry–Émery [BB21, BBD24]) *(formal in infinite dimensions) Suppose that there are real numbers (possibly negative) κ_t such that, for all $t\in[0,T]$ and all ϕ, as quadratic forms,*

$$\dot{Q}_t D^2 V_t(\phi)\,\dot{Q}_t + \frac{1}{2}\ddot{Q}_t \geq \kappa_t \dot{Q}_t.$$

Then ν_T satisfies the log-Sobolev inequality

$$\mathrm{Ent}_{\nu_T}(\rho) \leq \frac{2}{\gamma}\,\mathbb{E}_{\nu_T}[\|\dot{Q}_T^{1/2} D\sqrt{\rho}\|^2], \qquad \frac{1}{\gamma} = \int_0^T \exp\Big(-2\int_t^T \kappa_r\,\mathrm{d}r\Big)\,\mathrm{d}t.$$

The hypothesis gives, at each resolution, the gradient bound

$$\|\dot Q_t^{1/2} D\sqrt{G_t}\|^2 \le \mathrm{e}^{-2\int_t^T \kappa_r \mathrm{d}r} \int \|\dot Q_T^{1/2} D\sqrt{\rho}\|^2 \mathrm{d}\mu_t^\phi ;$$

inserted in the entropy formula it gives the theorem [BBD24]. For $V_T = 0$ and a linear profile $Q_t = t\,Q_T/T$ one may take $\kappa = 0$, so $1/\gamma = T$ and the theorem is the Gaussian log-Sobolev inequality $\mathrm{Ent}(\rho) \le 2\,\mathbb{E}\langle D\sqrt{\rho}, Q_T D\sqrt{\rho}\rangle$. The hypothesis is a convexity condition on the effective potentials at every scale, not on V_T, corrected by the scale dependence $\ddot Q_t$ of the decomposition; κ_t may be negative at some scales: only the weights $\mathrm{e}^{-2\int_t^T \kappa_r \mathrm{d}r}$ enter γ.

In terms of the conditional covariance. By Proposition 8.3, $D^2 V_t = S_t^{-1} - S_t^{-1}\mathrm{Cov}_t S_t^{-1}$, and $S_t^{-1}\dot Q_t S_t^{-1} = \dot{\mathcal{R}}_t$ is the derivative of the regulator (8.6). Since $\dot Q_t S_t^{-1} = S_t\,\dot{\mathcal{R}}_t$, the hypothesis reads

$$S_t\,\dot{\mathcal{R}}_t \mathrm{Cov}_t \dot{\mathcal{R}}_t S_t \le \dot Q_t S_t^{-1} \dot Q_t + \frac{1}{2}\ddot Q_t - \kappa_t \dot Q_t .$$

If $\dot Q_t$ is invertible, multiplying on the left by $S_t \dot Q_t^{-1} = \dot{\mathcal{R}}_t^{-1} S_t^{-1}$ and on the right by its transpose $\dot Q_t^{-1} S_t$ shows that it is equivalent to the explicit bound

$$\mathrm{Cov}_{\mu_t^\phi} \le S_t - S_t \dot Q_t^{-1}\Big(\kappa_t \dot Q_t - \frac{1}{2}\ddot Q_t\Big)\dot Q_t^{-1} S_t \qquad \forall \phi . \tag{8.17}$$

The second derivative $\ddot Q_t$ is not estimated: it is a property of the Gaussian decomposition, and it enters only through the combination $\kappa_t \dot Q_t - \frac{1}{2}\ddot Q_t$. When Q_t and S_t are functions of one operator (Fourier multipliers, say), the right-hand side is $S_t - S_t^2 \dot Q_t^{-2}(\kappa_t \dot Q_t - \frac{1}{2}\ddot Q_t)$ mode by mode. If $\kappa_t \dot Q_t > \frac{1}{2}\ddot Q_t$ (for the linear profile: $\kappa_t > 0$), every conditional law is strictly more concentrated than the unresolved Gaussian fluctuation; if not, Cov_t may exceed S_t, as in a double well at low resolution. A concave profile ($\ddot Q \le 0$) makes the bound smaller but forces κ down by the same amount: for the Gaussian with $Q_t = 1 - \mathrm{e}^{-t}$ one has $\ddot Q = -\dot Q$, the hypothesis holds with $\kappa = -\frac{1}{2}$, and $\mathrm{Cov}_t = S_t$. The hypothesis is thus an upper bound on the conditional covariance, uniform in the conditioning, measured in the metric $\dot{\mathcal{R}}_t$. Closely related is the *entropic stability* of Chen and Eldan [CE22] (see also [BBD24]): if, for all $r \le t$ and all ϕ, $\dot{\mathcal{R}}_r \mathrm{Cov}_{\mu_r^\phi} \dot{\mathcal{R}}_r \le \alpha_r \dot{\mathcal{R}}_r$, then

$$\mathrm{Ent}_{\nu_T}(\rho) \le \exp\Big(\int_0^t \alpha_r \mathrm{d}r\Big)\,\mathbb{E}\big[\mathrm{Ent}_{\mu_t^{\Phi_t}}(\rho)\big] :$$

the total entropy is controlled by the entropy that is still hidden at resolution t. The low-resolution part of the localisation is where the measure may be multimodal, and the condition says that it does not hide too much. With the Brascamp–Lieb bounds of Section 8.4 the conditional covariance is controlled by the Hessian of V_T when this is convex; for Φ_3^4 it is not, and the condition is a statement about the effective potentials, i.e. about the renormalisation group flow itself.

Exercises

Exercise 8.1. (Stochastic maximum principle) (a) For the control problem (9.1), perturb an optimal control $u \to u + \varepsilon v$ and show that $p_t := \mathbb{E}_t[-D V_T(X_T^u)]$ satisfies $u_t = \dot Q_t^{1/2} p_t$. Identify p with the force F of (7.7). (b) Show that $\mathrm{Y}_t = D F_t(\Phi_t)\,\dot Q_t^{1/2}$, and derive (8.3).

Exercise 8.2. (Characteristics) (a) Show that the transport equation $\partial_t F + D F[\dot Q F] = 0$ is solved by $F_t(x) = F_T(y)$, $y = x + S_t F_T(y)$, and that its characteristics are the trajectories of (8.1). (b) Verify that (8.16) solves $\partial_t V = \frac{1}{2}\langle D V, \dot Q D V\rangle$, and that it is the limit $\hbar \to 0$ of $-\hbar \log \int \mathrm{e}^{-V_T(x+\zeta)/\hbar} \mu^{\hbar S_t}(\mathrm{d}\zeta)$. (c) For $V_T(y) = \lambda y^4$ in zero dimensions, solve the equation for y as a power series in λ, and identify the coefficients as tree graphs.

Exercise 8.3. **(Conditional moments)** (a) Prove $\mathrm{Cov}(\mu_t^{\phi}) = S_t - S_t D^2 V_t(\phi)\, S_t$. (b) Prove (8.9).

Exercise 8.4. **(A symplectic shear)** (a) Show that Σ_t preserves $\sum_i \mathrm{d}x_i \wedge \mathrm{d}p_i$ iff S_t is symmetric. (b) Show that the image of the graph of $-DV_t$ under Σ_t is the graph of $-D\Gamma_t$, with Γ_t given by (8.11). (c) For a non-gradient force, show that Theorem 8.2 still holds, but that $\mathcal{F}_t$ need not be a gradient.

Exercise 8.5. **(The constant)** Differentiate (8.11) in t at fixed x (envelope theorem), and use (7.6) to show that the quadratic terms cancel and $\partial_t \Gamma_t = -\frac{1}{2}\mathrm{Tr}(\dot{Q}_t D^2 V_t)$. Deduce (8.12) with the stated constant.

Exercise 8.6. **(Gaussian check)** For $V_T(\varphi) = \frac{\kappa}{2}\langle \varphi, \varphi\rangle$, show that $\Gamma_t(\psi) = \frac{\kappa}{2}\langle \psi, \psi\rangle + \text{const}$, and solve the FBSDE (8.7) explicitly. Compare with Exercise 7.1.

Exercise 8.7. **(1PI effective action)** (a) Let $\mathcal{K}(j) = \log \int \mathrm{e}^{\langle j,\varphi\rangle} \mathrm{e}^{-V_T(\varphi)} \mu^{Q_T}(\mathrm{d}\varphi)$. Show that $\Gamma_0^{\mathrm{EAA}} = \mathcal{K}^*$, and that $D^2\Gamma_0^{\mathrm{EAA}}(\psi)$ is the inverse covariance of the tilted measure with mean ψ. (b) Integrate (8.12) with $D^2\Gamma_t$ replaced by $D^2 V_T$, and obtain the one-loop formula $\Gamma_t \approx V_T + \frac{1}{2}\mathrm{Tr}\log(1 + S_t D^2 V_T)$.

Exercise 8.8. **(Trees versus loops)** Using (8.11), show that $V_t = \Gamma_t - \frac{1}{2}\langle D\Gamma_t, S_t D\Gamma_t\rangle + O(\lambda^3)$ for the Φ^4 interaction (7.1). Deduce from Proposition A.2 that, to order λ^2, Γ_t is obtained from V_t by removing exactly the term $k=1$, the one-particle-reducible tree; the bubble ($k=2$), the sunset ($k=3$) and the vacuum term ($k=4$) are 1PI and remain. Where does the sunset come from in (8.12)?

Exercise 8.9. **(Regulators)** Derive (8.14). Compute $\mathcal{R}_t$ for the sharp Fourier profile, and for $\hat{Q}_t(k) = (|k|^2 + m^2)^{-1} - (|k|^2 + m^2 + t^2)^{-1}$ (whose kernels are not smooth, but the formulas make sense). In the last case show that $\mathcal{R}_t = t^2$ when $T = \infty$: a mass regulator.

Exercise 8.10. **(Brascamp–Lieb and Cramér–Rao)** Let $\mu(\mathrm{d}y) \propto \mathrm{e}^{-U(y)}\,\mathrm{d}y$ on $\mathbb{R}^n$, with mean m. (a) Show that $\mathbb{E}_\mu[(y-m)\otimes DU] = 1$ and $\mathbb{E}_\mu[DU \otimes DU] = \mathbb{E}_\mu D^2 U$, and deduce $\mathrm{Cov}_\mu \geq (\mathbb{E}_\mu D^2 U)^{-1}$. (b) For $D^2 U > 0$, prove the Brascamp–Lieb inequality for linear f by Hörmander's L^2 method: solve $Lg = f - \mathbb{E}_\mu f$ with $L = -\Delta + \langle DU, D\rangle$ and use $\mathbb{E}_\mu[(Lg)^2] \geq \mathbb{E}_\mu\langle Dg, D^2 U\, Dg\rangle$. (c) In zero dimensions, with $S_t = S$ and $V_T(y) = \lambda y^4$, compute both bounds numerically and compare with the conditional variance; repeat for the double well $V_T(y) = \frac{1}{2}y^4 - \frac{3}{2}y^2$.

Exercise 8.11. **(Entropy along the localisation)** (a) Derive the formula for $\mathrm{Ent}_{\nu_T}(\rho)$ as the integrated quadratic variation of M, and the chain rule. (b) Show that $DG_t(\phi) = S_t^{-1}\mathrm{Cov}_{\mu_t^\phi}(\rho, y)$. (c) For $V_T = 0$ and a linear profile, recover the Gaussian log-Sobolev inequality from Theorem 8.5. (d) In zero dimensions, with $Q_t = 1 - \mathrm{e}^{-t}$ and a convex V_T, find κ_t and the resulting constant, and compare with the Bakry–Émery criterion.

9 Variational structure

9.1 Finite volume: the Boué–Dupuis formula

Let $\mathbb{H}_a$ be the set of progressively measurable drifts $u\colon [0,T] \to L^2(M)$ with $\mathbb{E}\int_0^T \|u_s\|^2 \mathrm{d}s < \infty$, and set

$$I_t(u) := \int_0^t \dot{Q}_s^{1/2} u_s\,\mathrm{d}s, \qquad X^u := X + I(u),$$

the field driven by the control u under the Gaussian reference $\mathbb{P}_0$.

Theorem 9.1. **(Boué–Dupuis [BD98]; Polchinski as HJB)** (formal)

$$-\log \mathbb{E}_0[\mathrm{e}^{-V_T(X_T)}] = V_0(0) = \inf_{u\in\mathbb{H}_a} \mathbb{E}_0\Big[V_T(X_T^u) + \frac{1}{2}\int_0^T \|u_s\|^2 \mathrm{d}s\Big]. \tag{9.1}$$

The infimum is attained at the feedback control $u_s^* = -\dot{Q}_s^{1/2} D V_s(X_s^{u^*})$, and $\mathrm{Law}(X_T^{u^*}) = \nu_T$. The effective potential is the value function, and (7.6) is the Hamilton–Jacobi–Bellman equation of this control problem.

Proof. (Verification.) For any u, by Itô's formula and (7.6),

$$\begin{aligned}\mathrm{d}V_t(X_t^u) &= \Big[\tfrac{1}{2}\|\dot{Q}_t^{1/2} D V_t\|^2 + \langle \dot{Q}_t^{1/2} D V_t, u_t\rangle\Big]\mathrm{d}t + \mathrm{d}M_t \\ &= \Big[\tfrac{1}{2}\|u_t + \dot{Q}_t^{1/2} D V_t\|^2 - \tfrac{1}{2}\|u_t\|^2\Big]\mathrm{d}t + \mathrm{d}M_t,\end{aligned}$$

with $DV_t = DV_t(X_t^u)$ and a martingale M. Integrating and taking expectations,

$$\mathbb{E}_0\Big[V_T(X_T^u) + \frac{1}{2}\int_0^T \|u_t\|^2 \mathrm{d}t\Big] = V_0(0) + \frac{1}{2}\,\mathbb{E}_0 \int_0^T \|u_t + \dot{Q}_t^{1/2} DV_t(X_t^u)\|^2 \mathrm{d}t. \tag{9.2}$$

So the cost is $\geq V_0(0)$, with equality iff $u = u^*$. With this control X^{u^*} solves the same SDE as Φ in (7.7). □

Euler–Lagrange equation = FBSDE. Perturbing $u \to u + \varepsilon v$ in (9.1), the first-order condition is $\mathbb{E}\int_0^T \langle u_s + \dot{Q}_s^{1/2}\,\mathbb{E}_s[DV_T(X_T^u)], v_s\rangle \mathrm{d}s = 0$ for all adapted v, i.e.

$$u_s = -\dot{Q}_s^{1/2}\;\mathbb{E}_s[DV_T(X_T^u)] = \dot{Q}_s^{1/2} F_s. \tag{9.3}$$

This is (7.8). The FBSDE is the Euler–Lagrange equation of the variational problem, and the Polchinski equation is its Hamilton–Jacobi equation. These are the three faces of the same object: *the PDE* (Polchinski), *the stochastic dynamics* (FBSDE), and *the variational principle* (Boué–Dupuis).

Entropy. By Girsanov, $\frac{1}{2}\,\mathbb{E}_0 \int \|u\|^2 \geq \mathrm{Ent}(\mathrm{Law}(X^u) \,|\, \mathrm{Law}(X))$, which is at least the entropy of the terminal marginal. So (9.1) is a pathwise lift of the Gibbs variational principle $-\log Z_T = \inf_\nu \{\int V_T \,\mathrm{d}\nu + \mathrm{Ent}(\nu|\mu^{Q_T})\}$. The lift is useful because adapted drifts can be *constructed scale by scale*: one looks for u as a functional of the Gaussian process X, not as an unknown measure.

Generating functionals. Adding a test functional f,

$$\mathcal{W}_T(f) := -\log \int \mathrm{e}^{-f}\mathrm{d}\nu_T = \inf_u\, \mathbb{E}_0\Big[f(X_T^u) + V_T(X_T^u) + \frac{1}{2}\int_0^T \|u_s\|^2 \mathrm{d}s\Big] - V_0(0). \tag{9.4}$$

Bounds on $\mathcal{W}_T(f)$ uniform in T, for f in a rich class, give tightness, moment bounds and the identification of limits. For Φ_3^4 the naive problem does not have uniform bounds: the leading-order optimal drift $u_s \approx -4\lambda\,\dot{Q}_s^{1/2} {:}X_s^3{:}$ has a divergent cost. One renormalises by the change of variables $u = -4\lambda\,\dot{Q}^{1/2} {:}X^3{:} + r$. After it, and after further paracontrolled changes of variables, the singular terms are explicit polynomials in the Wick powers of X, cancelled by *the counterterms*, and the residual functional is coercive. Its control in Besov and paracontrolled norms, uniformly in T, is the construction of Φ_3^4 in [BG20]; see also [BG21].

9.2 Infinite volume

On $\mathbb{T}_L^d$ the free energy $V_0(0)$ and the effective potential V_t are extensive: they grow like L^d, and (9.1) has no limit as $L \to \infty$. Three objects are intensive and survive formally:

i. *the force.* For a local interaction $V_T(\varphi) = \int v_T(\varphi(x))\mathrm{d}x$, $DV_T(\varphi)(x) = v_T'(\varphi(x))$, and (7.8) reads, on $\mathbb{R}^d$,

$$\Phi_t = \int_0^t \dot{Q}_s F_s \,\mathrm{d}s + \int_0^t \dot{Q}_s^{1/2} \mathrm{d}W_s, \qquad F_t(x) = -\mathbb{E}_t[v_T'(\Phi_T(x))]. \tag{9.5}$$

Here every quantity is local, and the equation makes sense without a volume cutoff. Infinite-volume limits of $\nu_T^{(L)}$ are expected to be laws of solutions of (9.5). Uniqueness of such solutions (e.g. for small coupling, or by contraction in weighted spaces) is a uniqueness statement for the infinite-volume theory. Non-uniqueness is expected in the presence of phase transitions, e.g. for Φ_2^4 with a deep double-well potential [GJS75];

ii. *the Burgers flow* (7.12) of the force, for the same reason;

iii. *differences of free energies.* For a local f, $\mathcal{W}_T^{(L)}(f)$ in (9.4) is a difference of two extensive infima, and it stays bounded as $L \to \infty$. Its variational characterisation, through the infinite-volume Euler–Lagrange equation (9.5), is the basis of the infinite-volume analysis of Φ_2^4 and $\exp(\Phi)_2$ in [BG23].

Differences of free energies. Fix the volume, and let u^* be the optimal drift of (9.1), so that $X^{u^*} = \Phi$ is the Polchinski process (Theorem 9.1). In (9.4) the infimum may be taken over perturbations $u = u^* + w$ of the optimal drift. The drift enters linearly, so $X^{u^*+w} = \Phi + I(w)$ with $I_t(w) = \int_0^t \dot{Q}_s^{1/2} w_s \, \mathrm{d}s$, and by the Euler–Lagrange equation (9.3) the cross term of the energy is $\mathbb{E}\int_0^T \langle u_s^*, w_s \rangle \, \mathrm{d}s = -\mathbb{E}\langle D V_T(\Phi_T), I_T(w) \rangle$. The two extensive infima then cancel exactly, and what is left is a variational problem for the perturbation alone.

Proposition 9.2. (Variational principle relative to the FBSDE) *(formal) Let Φ be the Polchinski process, i.e. the solution of the FBSDE (7.8) driven by the Brownian motion W, and ν_T the law of Φ_T. For every f bounded from below,*

$$-\log \int \mathrm{e}^{-f} \, \mathrm{d}\nu_T = \inf_w \mathbb{E}\Big[f(\Phi_T + I_T(w)) + B_{V_T}(\Phi_T, I_T(w)) + \frac{1}{2}\int_0^T \|w_s\|^2 \, \mathrm{d}s \Big], \tag{9.6}$$

where the infimum runs over adapted w with $\mathbb{E}\int_0^T \|w_s\|^2 \, \mathrm{d}s < \infty$, and

$$B_V(\phi, h) := V(\phi + h) - V(\phi) - \langle D V(\phi), h \rangle$$

is the remainder of the first-order Taylor expansion of V. The minimiser satisfies $w_s = -\dot{Q}_s^{1/2} \, \mathbb{E}_s[D f(\Phi_T^f) + D V_T(\Phi_T^f) - D V_T(\Phi_T)]$, with $\Phi^f = \Phi + I(w)$: the perturbed field solves the FBSDE with terminal potential $V_T + f$, and its force is that of Φ corrected by $F_s^f - F_s = -\mathbb{E}_s[D f(\Phi_T^f) + D V_T(\Phi_T^f) - D V_T(\Phi_T)]$.

Proof. By (9.4), the left-hand side is $\inf_u J_f(u) - J_0(u^*)$, with

$$J_f(u) = \mathbb{E}\Big[f(X_T^u) + V_T(X_T^u) + \frac{1}{2}\int_0^T \|u_s\|^2 \, \mathrm{d}s \Big].$$

Writing $u = u^* + w$ and expanding $\frac{1}{2}\|u^* + w\|^2$, the terms $V_T(\Phi_T)$ and $\frac{1}{2}\int \|u^*\|^2$ cancel against $J_0(u^*)$, and the cross term is computed above. The first-order condition in w is obtained as in (9.3). □

For a local interaction $V_T(\varphi) = \int v_T(\varphi(x)) \, \mathrm{d}x$,

$$B_{V_T}(\phi, h) = \int b_T(\phi(x), h(x)) \, \mathrm{d}x, \qquad b_T(a, h) = v_T(a + h) - v_T(a) - v_T'(a) \, h,$$

and $b_T(a, 0) = 0$. So every term of (9.6) is local: there is no free energy and no extensive quantity, only the field Φ and the perturbation that f induces on it. For f supported in a bounded region one expects the optimal w, and with it $I(w)$, to decay away from the support, at the rate of the decay of correlations of the measure. Both sides of (9.6) then make sense in infinite volume, with Φ a solution of the infinite-volume FBSDE (9.5). This suggests the following formal description of the infinite-volume measures (at a fixed ultraviolet cutoff T):

an infinite-volume measure is the law of Φ_T for a solution Φ of (9.5) which satisfies (9.6), with $B = \int b_T \, \mathrm{d}x$, for all local f.

The Laplace functionals in (9.6) determine the measure, and the condition involves only intensive quantities. When the FBSDE has several solutions, as is expected in the presence of phase transitions, each solution gives a measure with its own variational principle; the perturbations w of finite energy only probe a given solution locally. For Φ_3^4 this remainder term contains the divergent counterterms, and, as for (9.4), the change of variables of [BG20] is needed before (9.6) has bounds uniform in T; in two dimensions, compare the infinite-volume analysis of [BG23].

The coercive estimates must then be *local*: one uses weighted norms and needs bounds whose constants do not depend on the volume. This infrared problem is the analogue, for the scale dynamics, of the "coming down from infinity" estimates for the parabolic dynamics.

Exercises

Exercise 9.1. (Optimal drift) (a) Derive (9.3) from (9.1), and check that the optimal u is a martingale up to the factor $\dot{Q}^{1/2}$. (b) For the quadratic interaction of Exercise 7.1, compute the optimal drift and the value (9.1) explicitly.

Exercise 9.2. (Cost of the leading drift) For $V_T = \lambda \int {:}\varphi^4{:}_T$ on $\mathbb{T}^3_L$, show that

$$\frac{1}{2}\,\mathbb{E}\int_0^T \|4\lambda\,\dot{Q}_s^{1/2}{:}X_s^3{:}\|^2 \mathrm{d}s = 12\lambda^2 L^3 \int Q_T(z)^4\,\mathrm{d}z,$$

which diverges linearly in T. Relate this to the $k=4$ term of (A.6).

Exercise 9.3. (The relative variational principle) (a) Derive (9.6) and the equation for the minimiser. (b) For a Gaussian ν_T (quadratic V_T) and a linear $f = \langle g, \cdot\rangle$, compute the remainder term, solve (9.6), and recover $-\log \int \mathrm{e}^{-\langle g,\cdot\rangle}\,\mathrm{d}\nu_T = \langle g, m\rangle - \frac{1}{2}\langle g, \Sigma\, g\rangle$, with m and Σ the mean and covariance. (c) For a local interaction and f supported in a ball, show that B_{V_T} only sees the region where $I(w) \neq 0$, and discuss why (9.6) is bounded uniformly in the volume.

Appendices: additional material

The following appendices complement the main lecture. They develop the algebraic and perturbative structure of the Polchinski flow, the Dyson–Schwinger equations at every scale, and stochastic localisation.

Appendix A Algebraic structure of the Polchinski flow

A.1 A semigroup of renormalisation maps

For a positive covariance S define the renormalisation map

$$\mathcal{R}_S(V) := -\log\,(\mu^S * \mathrm{e}^{-V}), \qquad (\mu^S * u)(\phi) = \int u\,(\phi+\zeta)\,\mu^S(\mathrm{d}\zeta). \tag{A.1}$$

Since $\mu^S * \mu^{S'} = \mu^{S+S'}$,

$$\mathcal{R}_S \circ \mathcal{R}_{S'} = \mathcal{R}_{S+S'} = \mathcal{R}_{S'} \circ \mathcal{R}_S, \qquad V_t = \mathcal{R}_{Q_{T,t}}(V_T). \tag{A.2}$$

So the effective potentials are an orbit of an *abelian* semigroup, indexed by the cone of positive covariances, acting nonlinearly on interactions and linearly on Boltzmann factors e^{-V}. Its infinitesimal generators are the vector fields

$$\mathcal{D}_B(V) := \tfrac{1}{2}\mathrm{Tr}(B\,D^2 V) - \tfrac{1}{2}\langle D\,V, B\,D\,V\rangle, \qquad \frac{\mathrm{d}}{\mathrm{d}s}\mathcal{R}_{sB}(V)\Big|_{s=0} = \mathcal{D}_B(V), \tag{A.3}$$

and (7.6) reads $\partial_t V_t = -\mathcal{D}_{\dot{Q}_t}(V_t)$.

Proposition A.1. (Flatness) (formal) *(a)* $[\mathcal{D}_B, \mathcal{D}_{B'}] = 0$ *for all positive* B, B'. *(b)* V_t *depends on the profile only through* $Q_{T,t}$. *In particular the free energy* $V_0(0)$ *and the law* ν_T *depend only on* Q_T, *not on the path* $t \mapsto Q_t$.

Proof. (a) Differentiate $\mathcal{R}_{sB}\,\mathcal{R}_{rB'} = \mathcal{R}_{rB'}\,\mathcal{R}_{sB}$ at $s = r = 0$, or compute directly (Exercise A.1). (b) is (A.2). □

Different profiles with the same endpoints are different *coordinate systems* on the same renormalisation group trajectory. They change the process Φ and the observables at intermediate resolution, but not the terminal theory. The flow has no holonomy [BCG26].

A.2 Perturbation theory and counterterms

Expanding $V_t = \mathcal{R}_{Q_{T,t}}(V_T)$ in cumulants of the Gaussian increment $\zeta \sim \mu^{Q_{T,t}}$ gives

$$V_t(\phi) = \mathbb{E}_\zeta\,[V_T(\phi+\zeta)] - \tfrac{1}{2}\mathrm{Var}_\zeta(V_T(\phi+\zeta)) + \dots \tag{A.4}$$

For (7.1), Exercise 6.1 of Lecture 3 shows that Gaussian convolution simply *lowers the Wick ordering*: since $Q_{T,t}(x,x) = c_T - c_t$,

$$V_t(\phi) = \int [\lambda H_4(\phi(x); c_t) + \gamma_T H_2(\phi(x); c_t)]\,\mathrm{d}x + O(\lambda^2). \tag{A.5}$$

Here $\gamma_T = O(\lambda^2)$ is kept for later use. At first order the effective potential is the Wick-ordered potential at the current resolution. This is the dual statement of Proposition 6.1 of Lecture 3: $\int H_4(X_t; c_t)$ is an observable.

Proposition A.2. (Second order) (formal) *Write $S = Q_{T,t}$ and $:\phi^p:_t = H_p(\phi; c_t)$. Then*

$$V_t(\phi) = \int [\lambda{:}\phi^4{:}_t + \gamma_T{:}\phi^2{:}_t] - \frac{\lambda^2}{2}\sum_{k=1}^{4}\binom{4}{k}^2 k! \iint S(x,y)^k\, {:}\phi^{4-k}{:}_t(x)\, {:}\phi^{4-k}{:}_t(y)\,\mathrm{d}x\,\mathrm{d}y, \tag{A.6}$$

up to $O(\lambda^3)$.

Proof. By the binomial formula for Hermite polynomials,

$$H_4(\phi+\zeta; c_t+s) = \sum_{k=0}^{4}\binom{4}{k} H_k(\zeta; s)\, H_{4-k}(\phi; c_t), \qquad s = S(x,x).$$

Then compute the variance in (A.4) with Lemma 3.8 of Lecture 2. □

Now let $T \to \infty$ at fixed t, so that $S \to C - Q_t$, which near the diagonal is as singular as G. In $d = 2$ every S^k is locally integrable, so with $\gamma_T = 0$ every term has a limit: Wick ordering suffices. In $d = 3$, $G(z) \sim (4\pi|z|)^{-1}$:

- $k = 1, 2$: S and S^2 are locally integrable. These terms are finite and nonlocal.
- $k = 3$ (the *sunset*): $S^3 \sim |z|^{-3}$ is not integrable. Writing $\phi(x)\,\phi(y) = \frac{1}{2}(\phi(x)^2 + \phi(y)^2) - \frac{1}{2}(\phi(x) - \phi(y))^2$, up to a field-independent constant the divergent part is local:

  $$-48\lambda^2\Big(\int S^3\Big)\int {:}\phi^2{:}_t + \text{(finite)},$$

  with $\int S^3 \sim \frac{1}{16\pi^2}\log T$. It is cancelled by

  $$\gamma_T := 48\lambda^2 \int_{\mathbb{T}_L^3} Q_T(z)^3\,\mathrm{d}z = \frac{3\lambda^2}{\pi^2}\log T + O(1), \tag{A.7}$$

  since $\gamma_T - 48\lambda^2\int S^3$ converges (Exercise A.2).
- $k = 4$: a divergent constant $-12\lambda^2 L^3 \int S^4 \sim T$, the vacuum energy. It cancels in ν_T but not in the free energy.

Power counting (Exercise A.3) shows that no further field-dependent counterterms are needed in $d = 3$: Φ^4_3 is superrenormalisable. Thus renormalisation appears as a property of the flow: one chooses the terminal data V_T so that the effective potentials V_t at *fixed* resolution have limits. The divergences at second order are also related to the singularity of Φ^4_3 with respect to the GFF [BG21]: the measures ν_t at fixed t are expected to stay absolutely continuous with respect to μ^{Q_t}, while the terminal one is not.

Exercises

Exercise A.1. (**Commuting flows**) Show by direct computation that $D\,\mathcal{P}_B(V)[\mathcal{P}_{B'}(V)] = D\,\mathcal{P}_{B'}(V)[\mathcal{P}_B(V)]$. Which property of D^2V and D^3V is used?

Exercise A.2. (**The sunset**) (a) Show that $\int_{|z|\le 1} G(z)^3\,\mathrm{d}z = \infty$ in $d=3$, and that $\int Q_T^3 = \frac{1}{16\pi^2}\log T + O(1)$ for the heat profile. (b) Show that $Q_T^3 - (Q_T - Q_t)^3$ is integrable uniformly in T, and deduce that the counterterm (A.7) makes the $:\phi^2:$ coefficient in (A.6) converge. (c) Show that the remaining part $\iint S^3(x,y)\,(\phi(x)-\phi(y))^2$ converges for smooth ϕ.

Exercise A.3. (**Power counting**) Show that the superficial degree of divergence of a connected graph of Φ_d^4 with n vertices and E external legs is $\omega = d - \frac{d-2}{2}E + (d-4)\,n$. Deduce that in $d=3$ the only divergent graphs with loops are the two-point graphs with $n \le 2$ (the tadpole and double tadpole, removed by Wick ordering, and the sunset) and vacuum graphs with $n \le 3$.

Appendix B Dyson–Schwinger equations at every scale

Assume now that all the operators involved are functions of A, hence commute, and let

$$K := Q_T^{-1}, \qquad J_t := K Q_t, \qquad J_0 = 0, \quad J_T = 1.$$

For the heat profile, $K = A\mathrm{e}^{A/T^2}$ and $J_t = \mathrm{e}^{-A(t^{-2}-T^{-2})}$ is a heat semigroup, which tends to e^{-A/t^2} as $T \to \infty$. F_t is the force observable of Theorem 7.4.

Theorem B.1. (**DS equations at resolution t**) (formal) *The* noise observable

$$\eta_t := K\,\Phi_t - J_t F_t \tag{B.1}$$

is a centred $\mathbb{P}$-martingale, with $\mathrm{d}\eta_t = K\,\dot{Q}_t^{1/2}\mathrm{d}W_t^{\mathbb{P}} - J_t\,\mathrm{d}F_t$. For every $t \le T$, every h and every functional G,

$$\mathbb{E}^{\mathbb{P}}\,[D_{J_t h}G(\Phi_t)] = \mathbb{E}^{\mathbb{P}}[G(\Phi_t)\,\langle h, \eta_t\rangle] = \mathbb{E}^{\mathbb{P}}[G(\Phi_t)\,\langle h, K\,\Phi_t - J_t F_t\rangle]. \tag{B.2}$$

At $t = T$ this is the Dyson–Schwinger equation of ν_T,

$$\mathbb{E}^{\nu_T}[D_h G] = \mathbb{E}^{\nu_T}[G\,\langle h, K\,\varphi + D\,V_T(\varphi)\rangle]. \tag{B.3}$$

Proof. From (7.7) and $\dot{J}_t = K\,\dot{Q}_t$, $\mathrm{d}\eta_t = K\,(\dot{Q}_t F_t \mathrm{d}t + \dot{Q}_t^{1/2}\mathrm{d}W_t^{\mathbb{P}}) - K\,\dot{Q}_t F_t \mathrm{d}t - J_t \mathrm{d}F_t$. The drift cancels, F is a martingale and $\eta_0 = 0$. For (B.2) use Proposition 7.2: $\Phi_t \sim \nu_t \propto \mathrm{e}^{-V_t}\mu^{Q_t}$. Gaussian integration by parts for μ^{Q_t} in the direction $Q_t k$ gives $\mathbb{E}^{\nu_t}[D_{Q_t k}G] = \mathbb{E}^{\nu_t}[G\,(\langle k, \varphi\rangle + \langle Q_t k, D\,V_t(\varphi)\rangle)]$. Take $k = Kh$, so that $Q_t k = J_t h$, and recall $F_t = -D\,V_t(\Phi_t)$. □

Formally letting $T \to \infty$, $K \to A$, and (B.1) becomes an exact identity between observables at every resolution:

$$A\,\Phi_t = A\,Q_t\,F_t + \eta_t, \tag{B.4}$$

with $A\,Q_t = \mathrm{e}^{-A/t^2}$ for the heat profile. This is the Euclidean field equation $A\varphi = -V'(\varphi) + \text{noise}$ of Lectures 1–2, seen at resolution t. It needs no products of distributions: the nonlinearity enters only through the force *observable* F_t, smeared at scale $1/t$. For the GFF ($F = 0$), (B.2) is Gaussian integration by parts for μ^{Q_t}. For Φ_3^4, where F_t plays the role of the renormalised cube, the limit $t \to \infty$ of (B.4) corresponds to the renormalised DS equation of [GH21].

Remark B.2. **(Comparison with stochastic quantisation)** In parabolic or elliptic stochastic quantisation (Lecture 3, §4) the field equation is imposed in an auxiliary time, and the field at fixed time has the right law. Here the field equation (B.4) holds at each resolution, and the auxiliary parameter is the resolution itself. Taking conditional expectations of an SPDE solution with respect to a scale filtration gives identities of the same shape, but with a different noise covariance: the martingale $J_t\,\mathrm{d}F_t$ in $\mathrm{d}\eta_t$ is essential [BCG26].

B.1 Gaussian bounds for the noise

The noise observable η_t is not Gaussian: it depends on the unknown law of the interacting field. Its moments are nevertheless bounded by those of an explicit Gaussian, uniformly in the interaction. Work at finite cutoff, where Φ_t has a Lebesgue density p_t, and let $0<t<T$. By Proposition 7.2, $-D\log p_t(a)=Q_t^{-1}a-F_t(a)$, and since $J_tQ_t^{-1}=K$,

$$\eta_t=-J_t\,D\log p_t(\Phi_t). \tag{B.5}$$

The noise is the score of the law of the field at resolution t, seen through J_t. Given its terminal value, the interacting field is still the Gaussian bridge: the density of $\mathbb{P}$ with respect to $\mathbb{P}_0$ depends on X_T only. Hence

$$\Phi_t=J_t\Phi_T+\zeta_t,\qquad \zeta_t\sim N(0,Q_t(1-J_t)),\quad \text{independent of } \Phi_T,$$

and only deterministic covariance operators, all functions of A, are commuted below. The variance that appears is the regulator (8.6) of the Wetterich equation: since $S_t=Q_T(1-J_t)$,

$$J_t^2\,(Q_t(1-J_t))^{-1}=KJ_t(1-J_t)^{-1}=S_t^{-1}-Q_T^{-1}=\mathcal{R}_t. \tag{B.6}$$

Proposition B.3. **(Sub-Gaussian noise)** *Let $0<t<T$. Then $\eta_t=\mathbb{E}[J_t\,(Q_t(1-J_t))^{-1}\zeta_t|\Phi_t]$, and η_t is dominated in the convex order by $N(0,\mathcal{R}_t)$: $\mathbb{E}\,\Psi(\eta_t)\le\mathbb{E}\,\Psi(\Xi)$, $\Xi\sim N(0,\mathcal{R}_t)$, for every convex Ψ for which both sides are defined. In particular, for every h and $u\in\mathbb{R}$,*

$$\mathbb{E}\,\mathrm{e}^{u\langle h,\eta_t\rangle}\le\exp\left(\frac{u^2}{2}\langle h,\mathcal{R}_t h\rangle\right): \tag{B.7}$$

every smeared noise observable is centred and sub-Gaussian, with the regulator as variance and with no convexity assumption on the interaction.

Proof. Write $B=Q_t(1-J_t)$ and g for the density of $N(0,B)$. Then $p_t(a)=\mathbb{E}\,g(a-J_t\Phi_T)$ and $Dg(z)=-B^{-1}z\,g(z)$, so $-D\log p_t(\Phi_t)=\mathbb{E}[B^{-1}\zeta_t|\Phi_t]$; apply J_t and use (B.5). By conditional Jensen, $\Psi(\eta_t)\le\mathbb{E}[\Psi(J_tB^{-1}\zeta_t)|\Phi_t]$, and $J_tB^{-1}\zeta_t$ is centred Gaussian with covariance $J_tB^{-1}J_t=\mathcal{R}_t$. □

For the heat profile (compare (8.14)), $K=A\,\mathrm{e}^{A/T^2}$ and $J_t=\mathrm{e}^{-A(t^{-2}-T^{-2})}$ give

$$\mathcal{R}_t=\frac{A\,\mathrm{e}^{A/T^2}}{\mathrm{e}^{A(t^{-2}-T^{-2})}-1}\le\frac{4}{3}t^2\quad(T\ge 2t),\qquad \mathcal{R}_t=\frac{A}{\mathrm{e}^{A/t^2}-1}\le t^2\quad(T=\infty)$$

(Exercise B.3). At a finite spatial cutoff all of this holds with terminal resolution $T=\infty$, before any continuum measure is constructed.

The roughness of the noise. Compare spectral densities for the heat profile with $T=\infty$ and $A=|k|^2+m^2$: white noise ξ has density 1, the GFF has A^{-1}, the free noise $\eta_t=A\,X_t$ has $A\,\mathrm{e}^{-A/t^2}$, and the bound of Proposition B.3 is

$$\mathcal{R}_t(k)=\frac{A}{\mathrm{e}^{A/t^2}-1}\approx\begin{cases}t^2, & |k|\ll t,\\ A\,\mathrm{e}^{-A/t^2}, & |k|\gg t.\end{cases}$$

Since $x/(\mathrm{e}^x-1)\le \mathrm{e}^{-x/2}$, $\mathscr{R}_t\le t^2\mathrm{e}^{-A/2t^2}$: at resolution t the noise is no larger than t times a white noise smoothed at scale $1/t$. Pointwise, $\mathrm{Var}\,\eta_t(x)\le\mathscr{R}_t(x,x)\lesssim t^{d+2}$, i.e. $\eta_t(x)=O(t^{1+d/2})$. This is the size at scale $1/t$ of a distribution of regularity $-1-d/2$ ($-5/2$ in $d=3$): seen at its own resolution, the noise is one derivative rougher than white noise ($-d/2$) and two rougher than the GFF ($1-d/2$), the same as the free noise $A\varphi=A^{1/2}\xi$ of the Dyson–Schwinger equation, whatever the interaction. It is not a bound on a limit distribution: η_t is a martingale in t, not the smoothing of a fixed distribution, and on a fixed test function (B.7) only gives $\langle h,\eta_t\rangle=O(t)$, while the free noise stays bounded. For Φ^4_3 no L^2 limit of the logarithmic derivative is expected [ALZ06]. Compared with the free noise, the bound is sharp at and above the resolution, where $\mathscr{R}_t/(A\mathrm{e}^{-A/t^2})=(1-\mathrm{e}^{-A/t^2})^{-1}\to 1$, and larger by the factor t^2/A on the resolved modes. Uniformly in the interaction it cannot be improved: for a pinning potential $V_T=\frac{\kappa}{2}\|\varphi\|^2$, the covariance of η_t tends to $\mathscr{R}_t$ as $\kappa\to\infty$ (Exercise B.3).

Remark B.4. (Pathwise bounds) (sketch) Work at finite spatial cutoff, with $T=\infty$ or $t\le T/2$, so that $\mathscr{R}_t$ obeys the bounds below uniformly in the cutoff. The norm $\|f\|_\varepsilon=\sup_x(1+|x|)^{-\varepsilon}|f(x)|$ is convex, so Proposition B.3 dominates $\|\eta_t\|_\varepsilon$ by $\|\Xi\|_\varepsilon$, with $\Xi\sim N(0,\mathscr{R}_t)$. For this Gaussian field, $\mathscr{R}_t(x,x)\lesssim t^{d+2}$ and $\langle\delta_x-\delta_y,\mathscr{R}_t(\delta_x-\delta_y)\rangle\lesssim t^{d+4}|x-y|^2$, so Dudley's bound and Gaussian concentration give Gaussian tails for $\|\Xi\|_\varepsilon$ at the level $t^{1+d/2}\sqrt{\log(2+t)}$, uniformly in L. Since η is a martingale, $\|\eta_s\|_\varepsilon$ is a submartingale, and Doob's inequality extends the tails to $\sup_{s\le t}$. Summing over dyadic t, almost surely

$$\|\eta_t\|_\varepsilon\le C(\omega)\,t^{1+d/2}\sqrt{\log(2+t)}\qquad\text{for all } t\ge 1\quad(1\le t\le T/2 \text{ if } T<\infty),$$

with C having Gaussian tails uniformly in the interaction and in the spatial cutoff. This is a pathwise bound at each resolution, in the sense above.

Control of the force. Write $\mathrm{d}F_s=\Upsilon_s\,\mathrm{d}W_s$ as in Section 8. By Theorem B.1, $\mathrm{d}\eta_s=(K\dot Q_s^{1/2}-J_s\Upsilon_s)\,\mathrm{d}W_s$, so $\int_0^t J_s\Upsilon_s\,\mathrm{d}W_s=K\int_0^t\dot Q_s^{1/2}\,\mathrm{d}W_s-\eta_t$. Itô's isometry, $(a+b)^2\le 2a^2+2b^2$ and (B.7) give

$$\mathbb{E}\int_0^t\|\Upsilon_s^*J_sh\|^2\,\mathrm{d}s\le 2\langle h,\mathscr{R}_th\rangle+2\langle h,KQ_tKh\rangle. \tag{B.8}$$

For the heat profile with $T=\infty$, $KQ_tK=A\,\mathrm{e}^{-A/t^2}\le t^2/\mathrm{e}$, so the right-hand side is at most $2(1+\mathrm{e}^{-1})\,t^2\|h\|_{L^2}^2$. This controls a smoothed derivative of the force, $\Upsilon_s=DF_s(\Phi_s)\,\dot Q_s^{1/2}$, without any estimate on the terminal force.

Consequence for Φ^4_3. In the field equation (B.4) the noise is therefore controlled a priori at each fixed resolution, uniformly in the ultraviolet cutoff. A bound on Φ_t uniform in the ultraviolet cutoff then follows from a coercivity estimate for the renormalised effective force F_t, which converts the control of $A\Phi_t-AQ_tF_t$ into control of Φ_t. This is the remaining analytic input, and it is where the large-field problem of Lecture 5 enters.

Exercises

Exercise B.1. (DS from the dynamics) Prove (B.2) without using Proposition 7.2, for G of the form $G_t(\Phi_t)$ with G_t solving (7.10). Show that the drift of $D_{J_th}G_t(\Phi_t)$ equals the covariation of $G_t(\Phi_t)$ with $\langle h,\eta_t\rangle$. Where is the symmetry of DF_t used?

Exercise B.2. (Gaussian check) For $V_T=0$ show that (B.2) reduces to $\mathbb{E}[D_{Q_tKh}G(X_t)]=\mathbb{E}[G(X_t)\langle Kh,X_t\rangle]$, and compare with the DS characterisation of the GFF in Lecture 2.

Exercise B.3. (Sub-Gaussian noise) (a) Check (B.5) and the bridge representation, and show that the Girsanov density depends only on X_T. (b) Compute $\mathscr{R}_t$ for the heat profile, and show that $\mathscr{R}_t\le\frac43t^2$ for $T\ge 2t$ and $\mathscr{R}_t\le t^2$ for $T=\infty$. (c) Compute the law of η_t for $V_T=0$ and for $V_T=\frac{\kappa}{2}\|\varphi\|^2$, and show that its covariance tends to $\mathscr{R}_t$ as $\kappa\to\infty$. In what sense is (B.7) sharp? (d) Show that $\mathscr{R}_t(x,x)\asymp t^{d+2}$ for $t\gg m$, and complete the argument of Remark B.4.

Appendix C Stochastic localisation

Stochastic localisation was introduced by Eldan [Eld13] to study measures in high dimension (see [Eld23] for a survey). The idea is to write a measure as a mixture of simpler ones, through a measure-valued martingale driven by noise. Proposition 7.2 says that the Polchinski process does exactly this for ν_T.

The localisation as a measurement. From (7.5) and $S_t^{-1} - Q_T^{-1} = \mathcal{R}_t$,

$$\mu_t^{\phi}(\mathrm{d}y) \propto \exp\Big(\langle j, y\rangle - \frac{1}{2}\langle y, \mathcal{R}_t y\rangle\Big)\,\nu_T(\mathrm{d}y), \qquad j = S_t^{-1}\phi.$$

The conditional law is ν_T tilted by a Gaussian factor of precision $\mathcal{R}_t$. Moreover, by the bridge representation of Appendix B.1 and $S_t^{-1} J_t = \mathcal{R}_t$,

$$j_t := S_t^{-1}\Phi_t = \mathcal{R}_t\Phi_T + N_t, \qquad N \text{ Gaussian, independent of } \Phi_T, \quad \mathbb{E}[N_s \otimes N_t] = \mathcal{R}_{s\wedge t}.$$

So j_t is a noisy measurement of the terminal field with signal-to-noise ratio $\mathcal{R}_t$, and $\mu_t^{\Phi_t}$ is the Bayesian posterior given the measurements up to t. The regulator of the Wetterich equation is the information gained. The forecast ψ_t and Cov_t are the posterior mean and covariance, and (8.9) describes how the uncertainty is resolved. The basic form of Eldan's scheme, with identity control, is the profile $S_t = (Q_T^{-1} + t)^{-1}$, $t \in [0,\infty)$, where $\mathcal{R}_t = t$ and $j_t = t\,\Phi_T + N_t$ with N a Brownian motion; Eldan's original construction [Eld13] uses an adapted control matrix.

Why it is useful. Three features make this decomposition a tool:

- *It is a martingale.* $\nu_T = \mathbb{E}\,\mu_t^{\Phi_t}$, and the expectation of any test function under $\mu_t^{\Phi_t}$ is a martingale. Itô calculus then splits variance and entropy exactly into the part already revealed and the part still hidden.
- *The pieces are simpler.* The Gaussian factor adds $\mathcal{R}_t$ to the Hessian of the potential. If ν_T is log-concave, μ_t^{ϕ} is $\mathcal{R}_t$-uniformly log-concave, so Bakry–Émery and Brascamp–Lieb apply to it even when their constants are useless for ν_T itself.
- *One quantity controls the loss.* What is revealed per unit resolution is governed by Cov_t, whose evolution (8.9) is explicit. An inequality for ν_T reduces to showing that the localisation does not happen too fast.

The process can also be run as an algorithm: with an approximate force it is a sampler, as in the algorithmic stochastic localisation of [EMS22], and denoising diffusions are of the same kind [Mon23]. By (B.5), the noise of the field equation is, up to J_t, the score that such algorithms estimate.

Proposition C.1. (Poincaré from the localisation) *Suppose that, for all $r \le t$ and all ϕ, $\dot{\mathcal{R}}_r \mathrm{Cov}_{\mu_r^{\phi}} \dot{\mathcal{R}}_r \le \alpha_r \dot{\mathcal{R}}_r$. Then, for every f,*

$$\mathrm{Var}_{\nu_T}(f) \le \exp\Big(\int_0^t \alpha_r\,\mathrm{d}r\Big)\,\mathbb{E}\,\mathrm{Var}_{\mu_t^{\Phi_t}}(f).$$

If moreover ν_T is log-concave, then

$$\mathrm{Var}_{\nu_T}(f) \le \exp\Big(\int_0^t \alpha_r\,\mathrm{d}r\Big)\,\mathbb{E}_{\nu_T}\langle Df, \mathcal{R}_t^{-1} Df\rangle.$$

Proof. Let $M_r = \mathbb{E}_r f(\Phi_T)$ and $\chi_r = \mathrm{Cov}_{\mu_r^{\Phi_r}}(f, y)$. As in Section 8.7, $\mathrm{d}\langle M\rangle_r = \langle \chi_r, \dot{\mathcal{R}}_r \chi_r\rangle\,\mathrm{d}r$, so $v(r) := \mathbb{E}\,\mathrm{Var}_{\mu_r^{\Phi_r}}(f)$ has $v' = -\mathbb{E}\langle \chi_r, \dot{\mathcal{R}}_r \chi_r\rangle$ and $v(0) = \mathrm{Var}_{\nu_T}(f)$. By Cauchy–Schwarz, with $h = \dot{\mathcal{R}}_r \chi_r$, $\langle \chi_r, h\rangle^2 \le \mathrm{Var}_{\mu_r^{\Phi_r}}(f)\,\langle h, \mathrm{Cov}_r h\rangle \le \alpha_r\,\mathrm{Var}_{\mu_r^{\Phi_r}}(f)\,\langle \chi_r, \dot{\mathcal{R}}_r \chi_r\rangle$. Hence $v' \ge -\alpha_r v$, which gives the first bound. For the second, the Hessian of the potential of μ_t^{ϕ} is at least $\mathcal{R}_t$: apply Brascamp–Lieb to μ_t^{ϕ} and take the expectation. □

For $V_T = 0$ one has $\mathrm{Cov}_r = S_r$. In one mode, or for the linear profile, where $\dot{Q}_r S_r^{-1}$ is a number, $\alpha_r = \dot{Q}_r S_r^{-1}$ and $\exp \int_0^t \alpha_r \, \mathrm{d}r = Q_T S_t^{-1}$: the first bound is an equality for linear f, and the second tends to the Gaussian Poincaré inequality as $t \to T$ (Exercise C.1).

An application: the KLS conjecture. Take an isotropic log-concave measure on $\mathbb{R}^n$ (mean zero, covariance 1) and the identity control, $\mathscr{R}_t = t$. The second bound reads $\mathrm{Var}(f) \le t^{-1} \, \mathrm{e}^{\int_0^t \alpha} \, \mathbb{E}|Df|^2$, so a Poincaré inequality follows if Cov_r stays bounded in operator norm up to some time t. The Kannan–Lovász–Simonovits conjecture asks for a Poincaré constant independent of n, and hence for a time t independent of n. Along the path, the bound holds only with high probability, not uniformly in ϕ, so the argument has to be adapted, and proving that the covariance stays bounded is the heart of the matter. Following Eldan [Eld13], this has given the bounds of Chen [Che21] and Klartag–Lehec [KL22], and then Klartag's bound [Kla23]: the KLS isoperimetric constant is at most $C\sqrt{\log n}$, i.e. the Poincaré constant is $O(\log n)$. The related slicing conjecture has since been proved [KL24].

The connection with Bauerschmidt–Bodineau. The multiscale Bakry–Émery criterion of Theorem 8.5 and the localisation study the same measure-valued martingale from two sides. Bauerschmidt and Bodineau [BB21] (see [BB19] for an earlier decomposition argument, and [BBD24] for a survey) start from the renormalisation group: integrate out the fine scales and follow the potential. Eldan's approach follows, along one path, the conditional law. The dictionary is:

this lecture	Bauerschmidt–Bodineau	stochastic localisation
resolution t increasing	their scale, reversed	time t increasing
law ν_t of Φ_t	renormalised measure	law of the tilt
μ_t^ϕ	fluctuation measure	tilted measure
S_t, $\mathscr{R}_t$	integrated covariance, —	—, t
$G_t = \int \rho \, \mathrm{d}\mu_t^\phi$	Polchinski semigroup on ρ	martingale $\int \rho \, \mathrm{d}\mu_t$
$D^2 V_t = S_t^{-1} - S_t^{-1} \mathrm{Cov}_t S_t^{-1}$	Hessian of the potential	covariance of the tilted measure

Proposition 8.3 is the bridge: convexity of the renormalised potentials is an upper bound on the conditional covariance, namely (8.17). The two methods then use this information differently.

- *Bauerschmidt–Bodineau.* The Kolmogorov equation (7.10) transports G_t by the drift $\dot{Q}_t F_t = -\dot{Q}_t D V_t$. Its Jacobian plays the role of the curvature in the Bakry–Émery computation, and the scale dependence of the metric $\dot{Q}_t$ adds $\frac{1}{2}\ddot{Q}_t$. This yields the gradient bound for G_t at each scale, which is integrated in the entropy formula. The output is a log-Sobolev inequality with an explicit constant, directly at the terminal scale, also far from log-concavity. The hypothesis is uniform in ϕ, which suits field theories where the renormalisation group controls V_t for all fields, as for sine-Gordon at $\beta < 6\pi$ [BB21].
- *Localisation.* One bounds what is lost on $[0, t]$ (the proposition above, or entropic stability [CE22]) and uses a known inequality for μ_t^ϕ. It works directly with moments of the conditional law, and can use bounds that hold only with high probability along the path, as in the KLS problem.

Exercises

Exercise C.1. (Poincaré from the localisation) (a) Check that $N_t = S_t^{-1} \zeta_t$ has covariance $\mathscr{R}_{s \wedge t}$. (b) For $V_T = 0$ and one mode, compute α_r and verify the Gaussian check above. (c) In zero dimensions, with $\mathscr{R}_r = r$ and $\nu = \frac{1}{2} N(a, \sigma^2) + \frac{1}{2} N(-a, \sigma^2)$, compute $\mathrm{Cov}_{\mu_r^\phi}$ and its supremum over ϕ. Using the first bound as $t \to \infty$, deduce the Poincaré constant $\le \sigma^2 \mathrm{e}^{a^2/\sigma^2}$. Compare with the true order $\mathrm{e}^{a^2/2\sigma^2}$: why is the exponent off by a factor 2?

References

[ALZ06] S. Albeverio, S. Liang, B. Zegarlinski. Remark on the integration by parts formula for the φ_3^4-quantum field model. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 9 (2006), 149–154.

[BB19] R. Bauerschmidt, T. Bodineau. A very simple proof of the LSI for high temperature spin systems. *J. Funct. Anal.* 276 (2019), 2582–2588.

[BB21] R. Bauerschmidt, T. Bodineau. Log-Sobolev inequality for the continuum sine-Gordon model. *Comm. Pure Appl. Math.* 74 (2021), 2064–2113.

[BBD24] R. Bauerschmidt, T. Bodineau, B. Dagallier. Stochastic dynamics and the Polchinski equation: an introduction. *Probab. Surveys* 21 (2024), 200–290.

[BCG26] I. Bailleul, I. Chevyrev, M. Gubinelli. Stochastic observables and Wilson–Itô diffusions. In preparation.

[BD98] M. Boué, P. Dupuis. A variational representation for certain functionals of Brownian motion. *Ann. Probab.* 26 (1998), 1641–1659.

[BG20] N. Barashkov, M. Gubinelli. A variational method for Φ_3^4. *Duke Math. J.* 169 (2020), 3339–3415.

[BG21] N. Barashkov, M. Gubinelli. The Φ_3^4 measure via Girsanov's theorem. *Electron. J. Probab.* 26 (2021), 1–29.

[BG23] N. Barashkov, M. Gubinelli. On the variational method for Euclidean quantum fields in infinite volume. *Probab. Math. Phys.* 4 (2023), 761–801.

[Bis73] J.-M. Bismut. Conjugate convex functions in optimal stochastic control. *J. Math. Anal. Appl.* 44 (1973), 384–404.

[BK87] D. C. Brydges, T. Kennedy. Mayer expansions and the Hamilton–Jacobi equation. *J. Stat. Phys.* 48 (1987), 19–49.

[BL76] H. J. Brascamp, E. H. Lieb. On extensions of the Brunn–Minkowski and Prékopa–Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *J. Funct. Anal.* 22 (1976), 366–389.

[BTW02] J. Berges, N. Tetradis, C. Wetterich. Non-perturbative renormalization flow in quantum field theory and statistical physics. *Phys. Rep.* 363 (2002), 223–386.

[CE22] Y. Chen, R. Eldan. Localization schemes: a framework for proving mixing bounds for Markov chains. In *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*, 2022, 110–122.

[Che21] Y. Chen. An almost constant lower bound of the isoperimetric coefficient in the KLS conjecture. *Geom. Funct. Anal.* 31 (2021), 34–61.

[Eld13] R. Eldan. Thin shell implies spectral gap up to polylog via a stochastic localization scheme. *Geom. Funct. Anal.* 23 (2013), 532–569.

[Eld23] R. Eldan. Analysis of high-dimensional distributions using pathwise methods. In *Proceedings of the ICM 2022*, EMS Press, 2023, 4246–4270.

[EMS22] A. El Alaoui, A. Montanari, M. Sellke. Sampling from the Sherrington–Kirkpatrick Gibbs measure via algorithmic stochastic localization. In *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)*, 2022.

[GH21] M. Gubinelli, M. Hofmanová. A PDE construction of the Euclidean Φ_3^4 quantum field theory. *Comm. Math. Phys.* 384 (2021), 1–75.

[GJS75] J. Glimm, A. Jaffe, T. Spencer. Phase transitions for φ_2^4 quantum fields. *Comm. Math. Phys.* 45 (1975), 203–216.

[JL01] G. Jona-Lasinio. Renormalization group and probability theory. *Phys. Rep.* 352 (2001), 439–458.

[JL64] G. Jona-Lasinio. Relativistic field theories with symmetry-breaking solutions. *Nuovo Cimento* 34 (1964), 1790–1795.

[JL75] G. Jona-Lasinio. The renormalization group: a probabilistic view. *Nuovo Cimento B* 26 (1975), 99–119.

[**KL22**] B. Klartag, J. Lehec. Bourgain's slicing problem and KLS isoperimetry up to polylog. *Geom. Funct. Anal.* 32 (2022), 1134–1159.

[**KL24**] B. Klartag, J. Lehec. Affirmative resolution of Bourgain's slicing problem using Guan's bound. Preprint, arXiv:2412.15044, 2024.

[**Kla23**] B. Klartag. Logarithmic bounds for isoperimetry and slices of convex sets. *Ars Inven. Anal.* (2023), Paper No. 4.

[**Mon23**] A. Montanari. Sampling, diffusions, and stochastic localization. Preprint, arXiv:2305.10690, 2023.

[**Mor94**] T. R. Morris. The exact renormalization group and approximate solutions. *Int. J. Mod. Phys. A* 9 (1994), 2411–2449.

[**MY99**] J. Ma, J. Yong. *Forward-Backward Stochastic Differential Equations and their Applications*. Lecture Notes in Math. 1702, Springer, 1999.

[**PBGM62**] L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze, E. F. Mishchenko. *The Mathematical Theory of Optimal Processes*. Interscience, 1962.

[**Pen90**] S. Peng. A general stochastic maximum principle for optimal control problems. *SIAM J. Control Optim.* 28 (1990), 966–979.

[**Pol84**] J. Polchinski. Renormalization and effective Lagrangians. *Nuclear Phys. B* 231 (1984), 269–295.

[**Sal99**] M. Salmhofer. *Renormalization: An Introduction*. Springer, 1999.

[**Wet93**] C. Wetterich. Exact evolution equation for the effective potential. *Phys. Lett. B* 301 (1993), 90–94.

[**YZ99**] J. Yong, X. Y. Zhou. *Stochastic Controls: Hamiltonian Systems and HJB Equations*. Springer, 1999.