

### **Epilepsy Awareness**

This course aims to equip healthcare professionals, caregivers, and individuals affected by epilepsy with the knowledge and skills necessary to understand the condition, provide appropriate care, and contribute to improving the quality of life for people with epilepsy in the UK.

### **Introduction to Epilepsy**

Epilepsy is a complex neurological condition characterized by recurrent seizures. In this module, we will provide an overview of epilepsy, including its prevalence, common causes, types of seizures, and diagnostic criteria.

- The Epilepsy is a chronic disorder of the brain that affects the normal electrical activity, leading to recurrent seizures.
- The latest transfer of transfer of the latest transfer of the latest
- Seizures in epilepsy can range from brief moments of unconsciousness or staring spells to convulsions and loss of control over bodily functions.

# Prevalence and Incidence of Epilepsy in the UK

- The Epilepsy is one of the most common neurological conditions in the UK, with a significant impact on individuals and their families.
- According to estimates, around 600,000 people in the UK have epilepsy, representing approximately 1 in 100 individuals.
- The incidence of epilepsy varies across age groups, with a higher prevalence observed in children and older adults.

#### **Common Causes and Risk Factors**

- TA Epilepsy can have various causes, and in about half of the cases, the underlying cause remains unknown (idiopathic epilepsy).
- The Common causes of epilepsy include brain injuries, such as head trauma, strokes, infections, developmental disorders, genetic factors, and brain tumours.
- The Certain risk factors, such as a family history of epilepsy, prenatal exposure to drugs or alcohol, and certain neurodevelopmental disorders, may increase the likelihood of developing epilepsy.

## Types of Seizures and Their Classification:

- Seizures in epilepsy can be classified into two main categories: focal seizures (previously known as partial seizures) and generalized seizures.
- Tocal seizures originate in a specific part of the brain and can result in various symptoms, including altered consciousness, abnormal movements, sensory disturbances, or emotional changes.
- Generalized seizures involve both hemispheres of the brain and can cause loss of consciousness, convulsions, muscle rigidity, or brief staring spells.

#### Diagnostic Criteria and Epilepsy Syndromes:

- The diagnosis of epilepsy is based on a comprehensive evaluation of the individual's medical history, seizure description, physical examination, and diagnostic tests.
- The International League Against Epilepsy (ILAE) provides guidelines and criteria for diagnosing epilepsy.
- TA Epilepsy syndromes are specific patterns of seizures and associated features that help classify and guide treatment decisions. Examples include childhood absence epilepsy, juvenile myoclonic epilepsy, and temporal lobe epilepsy.

Understanding the fundamental aspects of epilepsy, including its prevalence, causes, types of seizures, and diagnostic criteria, lays the groundwork for further exploration of epilepsy management, treatment options, and support systems.

## Epilepsy in the UK: Statistics and Epidemiology

Epilepsy is a significant neurological condition affecting a considerable number of individuals in the United Kingdom. Understanding the statistics and epidemiology of epilepsy is crucial for identifying the magnitude of the problem and developing targeted interventions.

#### **Prevalence and Incidence**

- According to the latest statistics, it is estimated that around 600,000 people in the UK have epilepsy, making it one of the most common neurological disorders.
- The prevalence of epilepsy varies across different age groups, with a higher incidence observed in children and older adults.
- Research suggests that epilepsy prevalence in the UK is slightly higher compared to other European countries.

## Demographic Patterns and Population Groups Affected

- The Epilepsy affects individuals of all ages, genders, and ethnicities. However, certain demographic patterns have been identified:
  - Thildren: Epilepsy is most commonly diagnosed in childhood, with a peak incidence between 1 to 5 years of age.
  - The highest incidence among individuals aged 75 years and older.
  - prevalence of epilepsy among individuals from lower socioeconomic backgrounds, suggesting a potential link between social determinants of health and epilepsy.

### Impact on Quality of Life

- TA Epilepsy can significantly impact various aspects of an individual's life, including education, employment, social relationships, and overall quality of life.
- Studies have shown that individuals with epilepsy may face educational challenges, such as learning difficulties and school absenteeism, leading to lower academic achievements.
- The compared to the general population, partly due to misconceptions, stigma, and concerns about seizure-related safety risks in the workplace.
- TA Epilepsy can also impact social relationships, leading to social isolation, anxiety, and depression.

## Health Disparities and Inequality in Epilepsy Care

- Research has highlighted disparities in epilepsy care and outcomes among different population groups within the UK, suggesting the presence of health inequalities.
- Certain ethnic minority groups, such as Black and South Asian communities, have been found to have a higher incidence of epilepsy, along with increased challenges in accessing appropriate healthcare services.
- There is a need to address these disparities through targeted interventions, culturally sensitive approaches, and improved access to epilepsy care for underserved populations.

#### **Diagnosis and Assessment**

Accurately diagnosing epilepsy is essential for providing appropriate treatment and support to individuals with the condition. We will explore the steps involved in diagnosing epilepsy, the significance of history-taking and observation, diagnostic tests and imaging techniques used in epilepsy, as well as the process of differential diagnosis to distinguish seizures from other conditions.

# Steps Involved in Diagnosing Epilepsy

- The diagnosis of epilepsy is typically made by a healthcare professional, such as a neurologist or epileptologist, through a systematic evaluation of the individual's medical history, symptoms, and diagnostic tests.
- The initial step involves taking a detailed history, which includes gathering information about the frequency, duration, and characteristics of seizures, as well as potential triggers or precipitating factors.
- A thorough physical examination is conducted to identify any neurological abnormalities or signs that may provide clues about the underlying cause of seizures.

### Importance of Accurate History-Taking and Observation

Accurate history-taking is crucial for the diagnosis of epilepsy as it helps healthcare professionals understand the pattern of seizures and identify any specific features or triggers.

- Individuals are encouraged to maintain a seizure diary, noting the date, time, duration, and description of each seizure episode, as well as any associated symptoms or warning signs.
- Observations by witnesses or caregivers can provide valuable insights into the seizure presentation and aid in determining the type of seizure.

# Diagnostic Tests and Imaging Techniques Used in Epilepsy

- Several diagnostic tests and imaging techniques are employed to support the diagnosis of epilepsy and assess the underlying cause of seizures.
- The brain and is a fundamental test for diagnosing and classifying seizures. It helps identify abnormal brain wave patterns associated with epilepsy.
- video-EEG Monitoring: This combines continuous EEG recording with video monitoring, allowing simultaneous documentation of seizures and their associated electrographic patterns.
- Neuroimaging: Magnetic resonance imaging (MRI) is commonly used to evaluate the brain structure and identify any structural abnormalities, such as tumours, vascular malformations, or scarring, that may be contributing to seizures.

### Differential Diagnosis and Distinguishing Seizures from Other Conditions

- Differential diagnosis involves distinguishing epilepsy from other conditions that may cause similar symptoms or seizure-like events.
- Some conditions that can mimic seizures include syncope (fainting), psychogenic nonepileptic seizures (PNES), sleep disorders, and movement disorders.
- Additional diagnostic tests, such as blood tests, genetic testing, and neuropsychological assessments, may be necessary to rule out other possible causes and confirm the diagnosis of epilepsy.

Accurate diagnosis and differentiation of seizures from other conditions enable healthcare professionals to develop an appropriate management and treatment plan tailored to the individual's specific needs. It also ensures that individuals with epilepsy receive timely and effective interventions, reducing the risk of misdiagnosis and unnecessary treatments.

#### **Antiepileptic Drugs (AEDs)**

Antiepileptic drugs (AEDs) play a crucial role in the management of epilepsy by reducing the frequency and severity of seizures. This session will provide an overview of common AEDs used in the UK, their mechanisms of action, potential side effects, individualizing AED therapy, and the importance of adherence to medication and therapeutic level monitoring.

#### Overview of Common AEDs Used in the UK

- There are several AEDs available in the UK, each with its own efficacy and safety profile.
- Commonly prescribed AEDs include carbamazepine, valproate, lamotrigine, levetiracetam, and oxcarbazepine, among others.
- The choice of AED depends on factors such as the type of seizures, epilepsy syndrome, age, gender, comorbidities, and individual response to specific medications.

## Mechanisms of Action and Side Effects

- AEDs act through various mechanisms to suppress abnormal neuronal activity and prevent the occurrence of seizures.
- The For example, sodium channel blockers (e.g., carbamazepine) reduce abnormal electrical discharges, while GABA-enhancing drugs (e.g., valproate) increase inhibitory neurotransmission.
- AEDs can have potential side effects, which vary depending on the medication. Common side effects include drowsiness, dizziness, cognitive impairment, gastrointestinal disturbances, and skin rashes.
- Some AEDs may also have specific risks, such as the potential for teratogenic effects (e.g., valproate) or cognitive side effects (e.g., topiramate).

## Individualizing AED Therapy and Treatment Options

- AED therapy should be individualized based on several factors, including seizure type, epilepsy syndrome, comorbidities, and the individual's lifestyle and preferences.
- Treatment goals include achieving seizure control, minimizing side effects, improving quality of life, and considering long-term management.
- In some cases, combination therapy (using multiple AEDs) may be required to achieve optimal seizure control.
- Newer AEDs with different mechanisms of action continue to be developed, expanding the treatment options available to individuals with epilepsy.

## Adherence to Medication and Monitoring Therapeutic Levels

- Adherence to prescribed AEDs is crucial for achieving effective seizure control and preventing breakthrough seizures.
- Healthcare professionals should educate individuals about the importance of taking medication as prescribed and provide strategies to improve adherence.
- Regular monitoring of therapeutic drug levels (when applicable) helps ensure that the individual is receiving an adequate dose for optimal seizure control.
- Therapeutic drug monitoring is particularly important for AEDs with narrow therapeutic indices, such as phenytoin or phenobarbital.

By understanding the common AEDs used in the UK, their mechanisms of action, potential side effects, and the importance of individualising therapy, healthcare professionals can work closely with individuals with epilepsy to develop personalized treatment plans. Promoting medication adherence and monitoring therapeutic drug levels contribute to optimizing seizure management

#### **Managing Epilepsy Effectively**

Managing epilepsy effectively involves a comprehensive approach that includes lifestyle modifications, self-management strategies, identification and avoidance of seizure triggers, developing seizure action plans, and implementing safety measures.

## Lifestyle Modifications and Self-Management Strategies

- Encouraging a healthy lifestyle can help individuals with epilepsy better manage their condition. This includes:
  - Adequate sleep: Ensuring a regular sleep schedule and addressing sleep-related issues can reduce seizure frequency.
  - The Stress management: Employing stress-reducing techniques, such as relaxation exercises or mindfulness, can be beneficial.
  - The Balanced diet: Maintaining a well-balanced diet and avoiding triggers like excessive caffeine or alcohol.
  - Regular exercise: Engaging in regular physical activity, based on individual capabilities, can contribute to overall well-being.

# Seizure Triggers and Their Avoidance

- and Identifying individual seizure triggers is crucial for managing epilepsy effectively. Common triggers may include:
  - ¬Lack of sleep or irregular sleep patterns
  - TAEmotional stress and anxiety
  - ¬

    ∠Hormonal changes (in women)
  - ¬¬Flickering lights or specific visual patterns (photosensitivity)
  - TACertain medications or drug interactions
- Once triggers are identified, individuals can take proactive steps to avoid or minimize exposure to them, thereby reducing the likelihood of seizures.

#### Developing Seizure Action Plans and Emergency Response Protocols

- Seizure action plans outline specific steps to be taken during and after a seizure, providing guidance to individuals, caregivers, and healthcare professionals.
- ¬¬ Action plans typically include:
  - ¬¬Description of seizure types and typical duration
  - namedication details and instructions on administering rescue medication, if prescribed
  - TAEmergency contact information and instructions for seeking medical assistance
  - Post-seizure care and recovery guidelines
- It is important to regularly review, and update seizure action plans based on individual needs and any changes in seizure patterns or treatment.

# Safety Measures and Seizure Precautions

- Implementing safety measures and seizure precautions can help minimize the risk of injury during seizures. These may include:
  - ¬¬Creating a safe environment: Removing sharp objects or potential hazards, securing furniture, and ensuring a clear space during activities.
  - Jacubing assistive devices: Wearing helmets or protective gear during activities that pose a higher risk of injury.
  - TAEducating family, friends, and caregivers: Providing information on seizure first aid, including proper positioning, protecting the head, and not restraining the person during a seizure.
  - Raising awareness: Informing teachers, employers, and other relevant individuals about the person's condition and necessary accommodations, if required.

By incorporating lifestyle modifications, identifying and avoiding seizure triggers, developing seizure action plans, and implementing safety measures, individuals with epilepsy can take an active role in managing their condition and reducing the impact of seizures on their daily lives.

Collaboration with healthcare professionals and education of family members, friends, and caregivers are also crucial for effective epilepsy management.

### Module 6: Seizure first aid and appropriate emergency management

Seizure first aid and appropriate emergency management are vital skills for individuals, caregivers, and bystanders to ensure the safety and well-being of someone experiencing a seizure. In this module, we will explore different types of seizures, recognizing and responding to seizures appropriately, administering first aid during seizures, and understanding when to seek emergency medical assistance.

#### 1. <u>Understanding Different Types of Seizures:</u>

- Seizures can manifest in various ways, and it is important to recognize the different types to provide appropriate aid. Common types of seizures include:
  - Generalized tonic-clonic seizures: Involves loss of consciousness, convulsions, and stiffening of the body.
  - Absence seizures: Brief staring spells with a temporary loss of awareness.
  - The Focal seizures: May involve altered consciousness, abnormal movements, sensory disturbances, or emotional changes, depending on the part of the brain affected.

### Seizure first aid and appropriate emergency management

Seizure first aid and appropriate emergency management are vital skills for individuals, caregivers, and bystanders to ensure the safety and well-being of someone experiencing a seizure. We will explore different types of seizures, recognizing and responding to seizures appropriately, administering first aid during seizures, and understanding when to seek emergency medical assistance.

### **Understanding Different Types of Seizures:**

Seizures can manifest in various ways, and it is important to recognize the different types to provide appropriate aid. Common types of seizures include:

- Generalized tonic-clonic seizures: Involves loss of consciousness, convulsions, and stiffening of the body.
- Absence seizures: Brief staring spells with a temporary loss of awareness.
- The Focal seizures: May involve altered consciousness, abnormal movements, sensory disturbances, or emotional changes, depending on the part of the brain affected.

## Recognizing and Responding to Seizures Appropriately

- Remain calm and stay with the person experiencing the seizure. Do not panic or restrain them.
- Protect the person from injury by moving away hazardous objects and creating a safe space.
- Time the duration of the seizure, as this information can help medical professionals later.
- Do not attempt to insert anything into the person's mouth during the seizure. Contrary to popular belief, it is not necessary and can cause harm.

# Administering First Aid During Seizures

- The Ensure the person's safety by gently guiding them to a lying or supported position, if possible.
- The Cushion their head with a soft object or clothing to prevent injury.
- Loosen tight clothing around their neck to facilitate breathing.
- Turn the person onto their side (recovery position) after convulsions stop to prevent choking on saliva or vomit.
- 3 Stay with them until they regain full consciousness and are fully alert.

### When to Seek Emergency Medical Assistance

- Most seizures are self-limiting and stop on their own. However, emergency medical assistance should be sought in the following situations:
  - The seizure lasts longer than five minutes (status epilepticus).
  - The person has difficulty breathing or turns blue.
  - AMultiple seizures occur without regaining consciousness in between.
  - <sub>¬¬</sub>Injury occurs during the seizure.
  - The person is pregnant, diabetic, or has a known heart condition.

It is crucial to educate individuals, caregivers, and bystanders about seizure first aid and emergency management.

By understanding the different types of seizures, recognizing and responding appropriately, administering first aid, and knowing when to seek emergency medical assistance, individuals can contribute to ensuring the safety and well-being of those experiencing seizures

#### Support Systems and Resources in the UK

Accessing appropriate support systems and resources is essential for individuals with epilepsy to manage their condition effectively. In this module, we will explore the epilepsy support organizations and networks available in the UK, accessing specialist epilepsy services and centers, understanding the role of healthcare professionals such as epilepsy nurses, neurologists, and epilepsy specialist nurses, as well as the legal rights and protections for individuals with epilepsy.

# Overview of Epilepsy Support Organizations and Networks in the UK

- The UK has several epilepsy support organizations and networks dedicated to providing information, resources, and support to individuals with epilepsy and their families.
- TA Examples include Epilepsy Action, Epilepsy Society, Young Epilepsy, and The Daisy Garland, among others.
- These organizations offer helplines, online communities, educational materials, and advocacy initiatives to raise awareness and promote the well-being of individuals with epilepsy.

### Accessing Specialist Epilepsy Services

- Specialist epilepsy services and centre's play a crucial role in the comprehensive management of epilepsy. They offer specialized care, diagnostic evaluations, treatment options, and ongoing support.
- ndividuals can access these services through referral from their primary care physician or neurologist. Some services may require a referral from a specialist epilepsy nurse or other healthcare professionals.

### Understanding the Role of Epilepsy Nurses, Neurologists, and Epilepsy Specialist Nurses

- The professionals who provide education, support, and guidance to individuals with epilepsy and their families. They assist in managing medication, monitoring seizure activity, and addressing concerns related to epilepsy.
- Neurologists: Neurologists are medical doctors who specialize in the diagnosis and treatment of neurological disorders, including epilepsy. They provide expertise in the management of epilepsy, conduct diagnostic evaluations, and prescribe appropriate medications or treatment options.
- The Epilepsy Specialist Nurses: Epilepsy specialist nurses are registered nurses with specialized training in epilepsy care. They work alongside neurologists to provide comprehensive care, education, and ongoing support to individuals with epilepsy.

# Legal Rights and Protections for Individuals with Epilepsy

- In the UK, individuals with epilepsy are protected by various legal rights and provisions to ensure equal opportunities and prevent discrimination.
- The Equality Act 2010 prohibits discrimination on the grounds of disability, including epilepsy, in various areas such as employment, education, and provision of goods and services.
- The Disability Discrimination Act (DDA) provides additional protections, requiring reasonable adjustments to be made to accommodate individuals with disabilities, including epilepsy.
- rights, seek legal advice if necessary, and advocate for their needs when accessing education, employment, healthcare, and other services.

- Healthcare professionals play a critical role in advocating for the legal rights and protections of individuals with epilepsy. By raising awareness, providing necessary documentation, and supporting individuals in navigating legal frameworks, healthcare professionals ensure equal opportunities and protection against discrimination.
- The management of epilepsy by healthcare professionals encompasses a comprehensive approach that includes accurate diagnosis, personalized treatment plans, ongoing support, and advocacy for the rights of individuals with epilepsy. By working collaboratively with individuals, families, and support organizations, healthcare professionals contribute to enhancing the well-being and quality of life of individuals living with epilepsy.



#### CONTACT INFO



+44 207 692 8709



admin@belmatt.co.uk info@belmatt.co.uk



www.belmatt.co.uk



Suite 570, 405 Kings Road Chelsea SW10 0BB

