



Prevalence of Diabetes

- •>4 million people with diabetes
- Prevalence rising with obesity
- Half a million undiagnosed
- •10% annual NHS budget spent on diabetes
- £8million per year on treatment of complications

1. DUK Facts and Statistics, 2016, https://www.diabetes.org.uk/Documents/Position%20statements/DiabetesUK Facts Stats Oct16.pdf, accessed May 2017

3

Types of Diabetes

| Type 1 | Type 2 | |
|---|--|--|
| Absence of insulin- auto-immune disease | Insufficient insulin/insulin resistance | |
| Age of onset- 6/12- 40+ years GAD positive c-peptide negative <80 | Generally >40yrs GAD negative c-peptide positive >200 | |
| Prevalence=10% | Prevalence=85% | |
| Quick onset- acute symptoms | Slow onset-unrecognised symptoms | |
| Treatment- dependant on insulin | Diet/Exercise/oral and injectable therapy | |
| Complications- Retinopathy/CKD | Complications- CVD | |
| | | |

www.belmatt.co.uk

What is diabetes?


- Diabetes mellitus is a condition in which the amount of glucose (sugar) in the blood is too high because the body cannot use it properly.¹
- Cardiovascular disease
- Inflammatory response


1. Diabetes UK, https://www.diabetes.org.uk/Diabetes-the-basic:

www.belmatt.co.u

5


Diabetes





Inflammatory Response

- Visceral Obesity
- Pro-Inflammatory cytokines
- Free Fatty Acids-Gycerol
- Insulin Resistance



www.belmatt.co.uk

_


Risk Factors for Type 2 Diabetes

Risk Factors

- Family History
- Lifestyle/obesity/smoking
- Ethnicity
- Age
- High blood pressure/cholesterol
- Dementia
- Gestational diabetes

Other causes

- · Drug related
- (steroid / psychotics/diuretics)
- Pancreatic disease/damage
- · Endocrine disease
- (cushing's disease)



Diagnosis

| Normal | Non-diabetic hyperglycaemia (NDH) | Diabetes |
|---|--------------------------------------|--------------|
| Random blood glucose level (2hours post meal) < 11mmols | | >11mmols |
| Fasting blood glucose level <7mmols | >5.5mmols | >7mmols |
| HbA1c <42mmols/mol | 42-48 mmols/mol | >48mmols/mol |
| | | |





Glycaemia Control

- HbA1c
- Individualised targets
- Blood Glucose Monitoring
- Relationship with Diet/activity
- Medication

15

CVD Prevention

• Diabetes Kidney Disease:

Bloods: u&e's- serum creatinine & -eGFR


Urine: albumin creatinine ratio (ACR)


• Hypertension:

BP

• Lipid Profile: Statins

www.belmatt.co.uk







Diabetic Kidney Disease

- Indicator of CVD Risk
- Importance of screening and identification
- Addressing lifestyle
- Optimising glycaemia and CVD risk factors
- Role of therapies and de-prescribing
- Safety- AKI/ Hypoglycaemia

www.belmatt.co.uk

19









Healthy eating/portion control

Mediterranean

Low carbohydrate/high fat diet

Very low calorie diet (VLCD)- e.g Fast 800

5:2 diet

Intermittent fasting- time restricted eating (TRE)



/ww.belmatt.co.uk

23


The T-Plate Model Www.belmatt.co.uk

Benefits of Exercise

- · 20% reduction in total mortality
- · 30% reduction in diabetes related deaths
- 50% reduction in fasting glucose
- 40% reduction in obesity related cancer deaths
- 15% reduction in LDL
- 18% increase in ADL
- · 30% reduction in triglycerides
- 10 mm Hg in systolic blood pressure
- · 20 mm Hg reduction in diastolic blood pressure

www.belmatt.co.u

25



Dave

- Age 54yrs
- T2D for 4 years
- Father had MI age 58yrs
- HbA1c 82mmols
- Weight 132kgs BMI 40
- BP 134/82
- Good kidney function
- Medication-only metforminmax dose
- Gliclazide 80mgs BD
- Travels a lot for work

ww.belmatt.co.uk

27

27

Dave

- Age <mark>54yrs</mark>
- T2D for 4 years
- Father had MI age <mark>58</mark>yrs
- HbA1c 82mmols
- Weight 132kgs BMI 40
- BP <mark>134/82</mark>
- Good kidney function
- Medication-only metformin- max do
- Gliclazide 80mgs BD
- Travels a lot for work

vww.belmatt.co.uk



28

