



— B E L M A T T —
HEALTHCARE TRAINING

Anaphylaxis, BLS , AED and Choking



TABLE OF CONTENTS

Page No.

		Page No.
1	ANAPHYLAXIS	3-14
2	ANAPHYLAXIS AND VACCINATION SETTINGS	16
3	BASIC LIFE SUPPORT	21-34
4	AED	35-36
5	RECOVERY POSITION	37-38
6	CHOKING	39-45

ANAPHYLAXIS



— BELMATT —
HEALTHCARE TRAINING

INTRODUCTION

Anaphylaxis is a severe, life-threatening, generalised or systemic hypersensitivity reaction. It is characterised by rapidly developing, life threatening problems involving: the airway and/or breathing and/or circulation. In most cases, there are associated skin and mucosal changes. The UK incidence of anaphylactic reactions is increasing, there are approximately 20 anaphylaxis deaths reported each year in the UK, although this may be a substantial under-estimate.

Aims & Objectives

To produce a comprehensive set of guidelines regarding the treatment of anaphylaxis.
To provide a framework which facilitates early recognition and diagnosis of anaphylaxis.
To promote consistency in the immediate emergency treatment of anaphylaxis.
To determine the roles and responsibilities of clinical staff.

Scope

This particularly involves those who carry out clinical procedures such as vaccination, steroid injections, acupuncture and local anaesthesia. It does not need to be carried by those practitioners who are not involved in any of these clinical activities.

Competencies

All staff administering Adrenaline (Epinephrine) must be registered and working within the guidelines and Code of Conduct of their professional body.

All clinical staff will complete life support skills as part of the annual mandatory clinical updating programme

All staff undertaking treatment of anaphylaxis should have competency in basic life support, having attended induction training and annual updating.

All individuals have a professional responsibility to attend mandatory training.

Competencies

Anaphylaxis is a severe, life-threatening, generalised or systemic hypersensitivity reaction. It is characterised by rapidly developing, life threatening problems involving: the airway and/or breathing and/or circulation. In most cases, there are associated skin and mucosal changes. The UK incidence of anaphylactic reactions is increasing, there are approximately 20 anaphylaxis deaths reported each year in the UK, although this may be a substantial under-estimate.

Patients to whom this procedure refers

This procedure relates to all patients treated by, or in contact with clinicians in their working environment, who exhibit symptoms or signs of anaphylaxis or a severe allergic reaction following administration of medication, local anaesthetic, vaccine, injection, acupuncture or history of exposure to other antigen.

Recognition & Diagnosis

A reaction may occur following exposure to a variety of different agents:-

- Drugs including:
 - Vaccines and Immunisations
 - Antibiotics
 - Aspirin
 - Non-Steroidal anti – inflammatory drugs (NSAIDs) or Heparin
 - Blood and blood products or Anaesthetic drugs
 - Local anaesthetics – particularly Mepivacaine (brand name Scandonest),
 - Lignocaine, Benzocaine, Procaine and Tetracaine.
 - Contrast media.
- Some foods such as shellfish, bananas, eggs.
- Peanuts and Tree Nuts
- Insect stings
- Latex products
- Cosmetic dyes

This is not an exhaustive list and any substance may be implicated as a causative allergen.

Anaphylactic reactions vary in severity and progress; they may be rapid or slow in onset. In rare events manifestations may be delayed by a few hours (adding to diagnostic difficulty), or persist for more than 24 hours.

The patient may show one or several of the following signs:-

- Hypotension
- Dyspnoea and wheezing
- Laryngeal oedema
- Classic angio-oedema (facial swelling)
- Urticarial rash and skin or mucosal changes (These alone are not a sign of anaphylaxis and may not be present in 20% of cases)

Other symptoms may include rhinitis, conjunctivitis, abdominal pain, vomiting, diarrhoea, and a commonly reported “a sense of impending doom”.

Cardiovascular collapse is a common manifestation, especially in response to intravenous drugs or stings, and is caused by vasodilatation and loss of plasma from the blood compartment

Recognition & Diagnosis

There is no single set of criteria or symptoms to identify an anaphylactic reaction. However this range of signs and symptoms, which in certain combinations, make a diagnosis more likely and become the assessed elements of recognition strategy.

Sudden onset and rapid progression of these symptoms, problems with airway, breathing or circulations and skin and mucosal changes are the triggers for treatment actions.

Immediate Action

Initial treatments should not be delayed by the lack of an incomplete history or definite diagnosis.

The initial treatment algorithm (reproduced in Appendix 1) involves:-

Calling for help and dialling 999, mentioning suspected anaphylaxis

Patient assessment based on ABCDE approach.

Remove suspected allergen where possible.

Consideration of patient position.

Administration of adrenaline.

1. Call for help

Making 999 call mentioning anaphylaxis informed dispatcher of need for adrenaline
clarify wording

2. ABCDE

Undertake examination, removing clothes considering dignity, are there adverse signs?

3. Allergen removal

If possible limit or remove exposure to potential allergen, stop any drug suspected and remove bee sting (Early attempts at removal are more important than method!)
DO NOT encourage vomiting if ingested allergen suspected.

Immediate Action

4. Patient position

NEVER STAND THE PATIENT UP! This may potentiate cardiovascular collapse. All victims should adopt a position of comfort, lying flat with or without leg elevation may be helpful for hypotension but unhelpful for breathing difficulties. Pregnant patients are best laying on their left side to reduce caval compression.

5. Adrenaline administration

Adrenaline is generally regarded as the most important drug for any severe anaphylactic reaction.

It should be administered intramuscularly for the immediate treatment of anaphylactic reaction and can be given in an emergency situation without the necessity of a Patient Group Direction (PGD) or prescription.

Adrenaline should be administered intramuscularly (preferably in the midpoint of the thigh, anterolateral aspect) to all patients with clinical signs of anaphylaxis.

Doses of Adrenaline

Adults

A dose of 500 micrograms adrenaline 1: 1000 solution (0.5 ml) should be administered intramuscularly, and repeated after 5 minutes in the absence of clinical improvement or if deterioration continues after the initial treatment, especially if consciousness becomes, or remains impaired. In some cases several doses may be required.

Children

The dose of adrenaline administered in children is determined by age. The recommended doses are based on what is considered to be safe and practical to draw up and inject in an emergency (Resuscitation Council UK (2008, 2012)).

> 12 years:	500 micrograms IM (0.5 mL) i.e. same as adult dose 300 micrograms IM (0.3 mL) if child is small or prepubertal
> 6 – 12 years:	300 micrograms IM (0.3 mL)
> 6 months – 6 years:	150 micrograms IM (0.15 mL)
< 6 months:	150 micrograms IM (0.15 mL)

Immediate Action

Additional Responses

Should staff with additional skills and competence be available to treat the patient reference can be made to the additional elements in the full anaphylaxis treatment algorithm shown in Appendix 2.

It must however be remembered the administration of drugs additional to adrenaline and any IV fluids requires prescription and confirmable competences that may not be found in all areas of LCHS.

IF IN DOUBT STAFF SHOULD RESTRICT ACTIONS TO THOSE IN THE INITIAL TREATMENT PROCESS.

Adrenaline for the treatment of anaphylaxis should be by intramuscular injection ONLY. IV adrenaline use only applies to those experienced in the use and titration of vasopressors in their normal clinical practice (e.g., anaesthetists, emergency physicians, intensive care doctors). It is therefore NOT appropriate to consider this route within LCHS.

Equipment

Patients Own Medication

If patient carries their own prescribed pre-filled autoinjector device, eg. Epipen, Jext, Emerade, anyone can assist the patient to take his or her medication.

These are single use devices and should be sent with the patient to secondary care.

Use of the device should follow manufacturer's instructions, familiarity can be improved using instructional videos.

Immediate Action

Community Teams

Packs containing adrenaline (epinephrine) are provided by the Community Team Lead / Service Leads (Appendix 3). Packs are provided to all new members of staff following completion of internal training and recalled and replaced when expired.

Addition of syringes and needles to the adrenaline (epinephrine) will be the responsibility of each individual practitioner from within their own service stock. Adrenaline will be ordered centrally and supplied to the Community Team Lead / Service Leads.

All packs are sealed and marked with an expiry date and a central register of the issue of packs to staff will be held by each Team Lead.

It is the responsibility of all practitioners to be aware of the expiry date of their own pack and contact team leads for replacement. The Community Team Lead /Service Leads must be advised of any changes to role or location. This will enable the tracking system for packs to be maintained effectively.

Adrenaline (epinephrine) should be kept in the original packaging. It should not be stored in hot places such as cars for prolonged periods or where it could be accessed by unauthorised persons.

Clinical Areas

Adrenaline for anaphylaxis is supplied via the usual route of medicines supply for each individual clinical area i.e. usual stock order form.

Checking of emergency drugs should be included in local processes to check expiry date, maintenance of seal and appropriate storage.

Further information refer to the Safe and Secure Handling of Medicines Policy for the processes for all medicines (LCHS P-CIG-20).

Record Keeping

Document the acute clinical features of the suspected anaphylactic reaction and record the time of the onset of reaction. This is the time that symptoms are first noticed.

Record the circumstances immediately before the onset of symptoms to help identify any potential trigger.

A full record must be kept of adrenaline administered paying particular attention to timings. Should the circumstances involve any drug, whether prescribed or otherwise, herbal or homeopathic treatment the reaction should be reported to the MHRA (Medicines and Healthcare products Regulatory Agency) using the Yellow Card scheme. Details of the Yellow Card reporting process can be found on the specific website; -

<https://yellowcard.mhra.gov.uk/>

Following any incident staff must report and record the episode in accordance with the Trusts Incident Reporting Policy using DATIX.

References

Resuscitation Council UK (2008, 2012) Emergency treatment of anaphylactic reactions. Guidelines for healthcare providers.

<https://www.resus.org.uk/resuscitation-guidelines/> Last accessed 20.02.2020

NICE (2011) Clinical Guideline 134 Anaphylaxis: assessment to confirm an anaphylactic episode and the decision to refer after emergency treatment for a suspected anaphylactic episode

<https://www.nice.org.uk/guidance/cg134/evidence/anaphylaxis-full-guidelinepdf-184946941>


Last accessed 20.02.2020

Appendix 1



Resuscitation Council (UK)

Anaphylactic reactions – Initial treatment



¹ Life-threatening problems:

Airway: swelling, hoarseness, stridor

Breathing: rapid breathing, wheeze, fatigue, cyanosis, SpO₂ < 92%, confusion

Circulation: pale, clammy, low blood pressure, faintness, drowsy/coma

² Intramuscular Adrenaline

IM doses of 1:1000 adrenaline (repeat after 5 min if no better)

- Adult 500 micrograms IM (0.5 mL)
- Child more than 12 years: 500 micrograms IM (0.5 mL)
- Child 6 -12 years: 300 micrograms IM (0.3 mL)
- Child less than 6 years: 150 micrograms IM (0.15 mL)

March 2008

Appendix 2



Resuscitation Council (UK)

Anaphylaxis algorithm

Anaphylactic reaction?

Airway, Breathing, Circulation, Disability, Exposure

Diagnosis - look for:

- Acute onset of illness
- Life-threatening Airway and/or Breathing and/or Circulation problems¹
- And usually skin changes

- Call for help
- Lie patient flat
- Raise patient's legs

Adrenaline²

When skills and equipment available:

- Establish airway
- High flow oxygen
- IV fluid challenge³
- Chlorphenamine⁴
- Hydrocortisone⁵

Monitor:

- Pulse oximetry
- ECG
- Blood pressure

1 Life-threatening problems:

- Airway: swelling, hoarseness, stridor
Breathing: rapid breathing, wheeze, fatigue, cyanosis, SpO₂ < 92%, confusion
Circulation: pale, clammy, low blood pressure, faintness, drowsy/coma

2 Adrenaline (give IM unless experienced with IV adrenaline)

IM doses of 1:1000 adrenaline (repeat after 5 min if no better)

- Adult 500 micrograms IM (0.5 mL)
- Child more than 12 years: 500 micrograms IM (0.5 mL)
- Child 6 -12 years: 300 micrograms IM (0.3 mL)
- Child less than 6 years: 150 micrograms IM (0.15 mL)

Adrenaline IV to be given only by experienced specialists

Titrate: Adults 50 micrograms; Children 1 microgram/kg

3 IV fluid challenge:

Adult - 500 – 1000 mL
Child - crystalloid 20 mL/kg

Stop IV colloid
if this might be the cause
of anaphylaxis

4 Chlorphenamine
(IM or slow IV)

- | | |
|-----------------------------------|-------------------|
| Adult or child more than 12 years | 10 mg |
| Child 6 - 12 years | 5 mg |
| Child 6 months to 6 years | 2.5 mg |
| Child less than 6 months | 250 micrograms/kg |

5 Hydrocortisone
(IM or slow IV)

- | |
|--------|
| 200 mg |
| 100 mg |
| 50 mg |
| 25 mg |

March
2008

5th Floor, Tavistock House North, Tavistock Square, London WC1H 9HR
Telephone (020) 7388-4678 • Fax (020) 7383-0773 • Email enquiries@resus.org.uk
www.resus.org.uk • Registered Charity No. 286360

Appendix 3 Adrenalin (epinephrine) Packs in Community Services

All packs to contain:

Item	Amount	NHS Supply Chain
Blunt Needles	X2	FTR1827
Blue Needles	X4	
Green Needles	X4	
1ml syringes	X 4	FWC429



All packs to contain laminated dosage chart:

Age Dose	Dose	
<6 months	150 micrograms IM	0.15 ml 1:1000 solution
6 months – 6 years	150 micrograms IM	0.15 ml 1:1000 solution
6 – 12 years	300 micrograms IM	0.3 ml 1:1000 solution
>12 years	500 micrograms IM	0.5 ml 1:1000 solution
Small or pre-pubertal child	300 micrograms IM	0.3 ml 1:1000 solution



Adrenaline (epinephrine) should be;

- Kept in the original packaging and should not be split
- Not be stored in hot places such as cars for prolonged periods of time
- Not be stored in cars over night or where it could be accessed by unauthorised persons.



All packs should be sealed and marked with an expiry date and a central register of the location of packs will be held by each Team Lead. If not sealed these boxes need opening and checking daily

DS MEDICAL Clear View Zip Pouch – Red Product
Code: DS02315 £9.95 (EX. VAT)

Security tag; numbered and record of checks made
Scissors to be ready available




An appropriate sized plastic container
which will hold contents as above

Seal robust to show if pack has been

All packs must have an outside Label

	Batch Number	Expiry Date
Adrenaline (epinephrine) 1:1000 1mg/ 1ml		
Date Sealed	Signature	

Management of Anaphylaxis In The Vaccination Setting



MANAGEMENT OF ANAPHYLAXIS IN THE VACCINATION SETTING

Vaccines stimulate an immune response to cause protection. Most adverse events after vaccination are due to this immune response and are not allergic. Allergic reactions can happen after vaccination, but these are uncommon. Anaphylaxis following vaccination is rare, occurring at less than 1 per million doses for vaccines in the UK. "COVID-19 vaccines" are new, so it is uncertain if the risk of anaphylaxis is higher compared to other vaccines.

Anaphylaxis is a serious allergic reaction that is usually rapid in onset and may cause death. Severe anaphylaxis is characterised by life-threatening compromise in airway, breathing and/or the circulation, but **may occur without typical skin features or circulatory shock being present**.

Anaphylaxis is likely when all of the following 3 criteria are met:

- Sudden onset and rapid progression of symptoms
- Airway and/or Breathing and/or Circulation problems
- Skin and/or mucosal changes (itching, flushing, urticaria, angioedema)

Remember:

- Skin or mucosal changes alone are not a sign of anaphylaxis
- Skin and mucosal changes can be subtle or absent in 10-20% of reactions (e.g. some patients can present with only bronchospasm or hypotension)

Airway problems	Breathing problems	Circulation problems
<ul style="list-style-type: none">▪ Airway swelling e.g. throat and tongue swelling causing difficulty in breathing and/or swallowing. Patients may feel their throat is closing.▪ Hoarse voice▪ Stridor (a high-pitched inspiratory noise caused by upper airway obstruction)	<ul style="list-style-type: none">▪ Shortness of breath – increased respiratory rate▪ Wheeze (bronchospasm) and/ or persistent cough▪ Patient becoming tired with the effort of breathing▪ Confusion due to hypoxia▪ Cyanosis (a late sign)▪ Respiratory arrest	<ul style="list-style-type: none">▪ Signs of shock: pale, clammy▪ Significant tachycardia▪ Arrhythmia▪ Hypotension – feeling faint▪ (dizziness), collapse▪ Decreased conscious level or loss of consciousness▪ Cardiac arrest

Fainting

Fainting is relatively common following vaccination. Symptoms should resolve rapidly on lying flat. **If rapid recovery does not happen, treat as anaphylaxis and give IM adrenaline.**

	Faint	Anaphylaxis
Onset	Before, during or within minutes of vaccination	Usually within 15 minutes, but can occur later
Features	<ul style="list-style-type: none">Generalised pallor, cold clammy skinNormal respiration – may be shallow, but not labouredBradycardia, but with strong central pulse, Hypotension usually transient and responds to lying the patient downSense of light-headedness/dizziness, improves lying down. Transient jerking of the limbs and eye-rolling which may be confused with seizure.	<ul style="list-style-type: none">Itchy skin rash / urticaria (wheals) or swelling (angioedema) e.g. lips, faceAirway/Breathing symptoms (see above)Tachycardia, with weak/absent central pulse; hypotension – sustainedSense of “impending doom”; loss of consciousness – with no improvement once supine or head down position

Treatment of anaphylaxis

Treatment of anaphylaxis is based on general life support principles:

Call for help immediately.

Use the **Airway, Breathing, Circulation** approach to recognize and treat problems.

Intramuscular adrenaline is the first-line treatment for anaphylaxis (even if intravenous access is available). IV adrenaline

bolus outside the context of cardiac arrest is dangerous.

- Give intramuscular (IM) adrenaline to treat Airway/Breathing/Circulation problems.
- Do not delay initial treatment if the diagnosis is unclear: a single dose of IM adrenaline is well-tolerated and poses minimal risk.
- Repeat IM adrenaline every 5 minutes if features of anaphylaxis do not resolve.

Further treatment of anaphylaxis depends on:

1. Location

Treating a patient with anaphylaxis in the community will not be the same as in an acute hospital. Dial 999 urgently for ambulance support and clearly state “ANAPHYLAXIS”.

2. Training of rescuers

All clinical staff should be able to recognise anaphylaxis, call for help and start treatment. Staff who give immunisations should have annual updates in anaphylaxis management.

3. Number of responders

A single responder must always ensure that help is coming. If there are several rescuers, several actions can be undertaken simultaneously.

4. Equipment and drugs available

An anaphylaxis pack normally contains at least:

- Two ampoules of adrenaline (epinephrine) 1mg/mL (1:1000)
- Four 23G needles and four graduated 1mL syringes
- Oxygen supply, with face masks suitable for children and adults and tubing.

A pack should be immediately available in each location where vaccines are being given, and should not be stored in a locked cupboard or trolley. Check packs regularly to ensure the contents are within their expiry dates.

Antihistamines and steroids are no longer recommended for the immediate management of anaphylaxis, and do not need to be included in the emergency anaphylaxis pack.

Treatment of anaphylaxis

In all healthcare settings, giving adrenaline by ampoule/needle/syringe is preferred, since auto-injectors will only deliver a maximum of 300micrograms adrenaline while **the appropriate dose in older children and adults is 500micrograms**. Some settings may prefer to use an auto-injector for the first dose used to treat anaphylaxis, for speed and ease. If further doses of adrenaline are needed, give these by ampoule/ needle/syringe.

Patient positioning

Death can occur within minutes if a patient stands, walks or sits up suddenly. Patients must NOT walk or stand during acute reactions. Use caution when transferring patients who have been stabilised.

- Patients with Airway and Breathing problems may prefer to sit up.
- Lying flat with or without leg elevation is helpful for patients with a low blood pressure (Circulation problem).
- Patients who are breathing normally and unconscious should be placed on their side (recovery position). Monitor breathing and intervene if needed.
- Pregnant patients should lie on their left side to prevent aortocaval compression.

Other supportive measures:

- Give further repeat doses of IM adrenaline every 5 minutes if symptoms do not resolve.
- Give oxygen and apply pulse oximetry (if available), to achieve oxygen saturations of 94-98%, but do not delay giving oxygen while waiting for a pulse oximeter.

Severe upper airways obstruction is uncommon.

- Seek urgent expert help if there is airway obstruction.
- Nebulised adrenaline (5mL of 1mg/mL adrenaline) can be used to treat upper airways obstruction but **must not be prioritised over further IM adrenaline every 5 minutes**.

- Bronchospasm - consider further inhaled bronchodilator therapy with salbutamol and/or ipratropium, but must not be prioritised over IM adrenaline.
- A reduction in blood flow is common in anaphylaxis, even in the absence of obvious circulatory compromise. Give a fluid bolus if there is an inadequate response to initial IM adrenaline, to support tissue perfusion and drug delivery.

Possible cross-reactivity between vaccines and drugs

Antihistamines and steroids are no longer recommended for

management of anaphylaxis. Some oral medicines used to treat more mild reactions contain the same substances found in some vaccines (and which can cause allergic reactions). Caution is recommended following allergic reactions to vaccines. In the UK, use an oral liquid antihistamine (e.g. liquid cetirizine). Chlorphenamine tablets can also be used, but these can cause drowsiness and therefore mimic symptoms of anaphylaxis.

Treatment of anaphylaxis

CARDIAC ARREST:

Recognise cardiac arrest has occurred if the person becomes unresponsive or unconscious, and breathing is absent or abnormal.

- Start chest compressions as soon as cardiac arrest is suspected.
- Ensure expert help (resuscitation team or ambulance) has been called.
- Follow standard cardiac arrest guidelines (use IV/IO adrenaline in preference to IM route, as per protocol), available at www.resus.org.uk.

BLOODS:

The specific test to help confirm a diagnosis of anaphylaxis is measurement of mast cell tryptase. Ideally, three timed samples (serum or plasma, e.g. yellow top bottle) are needed:

1. An initial sample as soon as feasible, but do not delay starting resuscitation.
2. A second sample at 1-2 hours (but no later than 4 hours) from the onset of symptoms.

If possible, take 5-10mL extra for serum store to facilitate further investigations.

A convalescent sample should be obtained at least 24 hours after complete resolution (e.g. at follow-up allergy clinic), to provide a baseline tryptase value.

SURVEILLANCE:

- Report all suspected vaccine-induced adverse drug reactions (ADRs) via the MHRA's Yellow Card scheme: <https://coronavirus-yellowcard.mhra.gov.uk>
- All patients with suspected anaphylaxis should be referred to an allergy clinic – see bsaci.org for details of clinics.

BASIC LIFE SUPPORT



— BELMATT —
HEALTHCARE TRAINING

Introduction

The community response to cardiac arrest is critical to saving lives. Each year, UK ambulance services respond to approximately 60,000 cases of suspected cardiac arrest. Resuscitation is attempted by ambulance services in less than half of these cases (approximately 28,000).³ The main reasons are that either the victim has been dead for several hours or has not received bystander CPR so by the time the emergency services arrive the person has died. Even when resuscitation is attempted, less than one in ten victims survive to go home from hospital. Strengthening the community response to cardiac arrest by training and empowering more bystanders to perform CPR and by increasing the use of automated external defibrillators (AEDs) at least doubles the chances of survival and could save thousands of lives each year.^{4,5}

This guideline is based on the International Liaison Committee on Resuscitation (ILCOR) 2015 Consensus on Science and Treatment Recommendations (CoSTR) for Basic Life Support and Automated External Defibrillation and the European Resuscitation Council Guidelines for Resuscitation 2015 Section 2 Adult basic life support and automated external defibrillation.^{2,6} These contain all the reference material for this section.

Chain of Survival

The Chain of Survival (Figure 1) describes four key, inter-related steps, which if delivered effectively and in sequence, optimise survival from out-of-hospital cardiac arrest.

1: Early recognition and call for help

If untreated, cardiac arrest occurs in a quarter to a third of patients with myocardial ischaemia within the first hour after onset of chest pain. Once cardiac arrest has occurred, early recognition is critical to enable rapid activation of the ambulance service and prompt initiation of bystander CPR.

2: Early bystander CPR

The immediate initiation of bystander CPR can double or quadruple survival from out-of-hospital cardiac arrest. Despite this compelling evidence, only 40% of victims receive bystander CPR in the UK.¹

Chain of Survival

3: Early defibrillation

Defibrillation within 3–5 min of collapse can produce survival rates as high as 50–70%.¹⁵ This can be achieved through public access defibrillation, when a bystander uses a nearby AED to deliver the first shock.^{4,15–17} Each minute of delay to defibrillation reduces the probability of survival to hospital discharge by 10%. In the UK, fewer than 2% of victims have an AED deployed before the ambulance arrives.

4: Early advanced life support and standardised postresuscitation care

Advanced life support with airway management, drugs and the correction of causal factors may be needed if initial attempts at resuscitation are unsuccessful.

The quality of treatment during the post-resuscitation phase affects outcome and is addressed in the Adult advanced life support and Post-resuscitation care sections.

Improving survival from out-of-hospital cardiac arrest

Resuscitation Council UK recommends that to improve survival from cardiac arrest:

1. All school children are taught CPR and how to use an AED.
2. Everyone who is able to should learn CPR.
3. Defibrillators are available in places where there are large numbers of people (e.g. airports, railway stations, shopping centres, sports stadiums), increased risk of cardiac arrest (e.g. gyms, sports facilities) or where access to emergency services can be delayed (e.g. aircraft and other remote locations).
4. Owners of defibrillators should register the location and availability of devices with their local ambulance services.
5. Systems are implemented to enable ambulance services to identify and deploy the nearest available defibrillator to the scene of a suspected cardiac arrest.
6. All out-of-hospital cardiac arrest resuscitation attempts are reported to the National Out-of-Hospital Cardiac Arrest Audit.

Resuscitation Council UK's BLS/AED guidelines

The remainder of this section contains guidance on the initial resuscitation of an adult cardiac arrest victim where the cardiac arrest occurs outside a hospital.

This includes basic life support (BLS: airway, breathing and circulation support without the use of equipment other than a protective barrier device) and the use of an automated external defibrillator (AED). Simple techniques used in the management of choking (i.e. foreign body airway obstruction) are also included.

Guidelines for the use of manual defibrillators and starting in-hospital resuscitation are found in Advanced life support guidelines section.

The guidelines are based on the ILCOR 2015 Consensus on Science and Treatment Recommendations (CoSTR) for BLS/AED and European Resuscitation Council Guidelines for BLS/AED.

Key messages from Guidelines 2015

- Ensure it is safe to approach the victim.
- Promptly assess the unresponsive victim to determine if they are breathing normally.
- Be suspicious of cardiac arrest in any patient presenting with seizures and carefully assess whether the victim is breathing normally.
- For the victim who is unresponsive and not breathing normally:
 - Dial 999 and ask for an ambulance. If possible stay with the victim and get someone else to make the emergency call.
 - Start CPR and send for an AED as soon as possible.
 - If trained and able, combine chest compressions and rescue breaths, otherwise provide compression-only CPR.
 - If an AED arrives, switch it on and follow the instructions.
 - Minimise interruptions to CPR when attaching the AED pads to the victim.
- Do not stop CPR unless you are certain the victim has recovered and is breathing normally or a health professional tells you to stop
- Treat the victim who is choking by encouraging them to cough. If the victim deteriorates, give up to 5 back slaps followed by up to 5 abdominal thrusts. If the victim becomes unconscious – start CPR.
- The same steps can be followed for resuscitation of children by those who are not specifically trained in resuscitation for children – it is far better to use the adult BLS sequence for resuscitation of a child than to do nothing.

Adult BLS Sequence

The sequence of steps for the initial assessment and treatment of the unresponsive victim are summarised in Figure 2. Further technical information on each of the steps is presented in Table 1 and below.

The sequence of steps takes the reader through recognition of cardiac arrest, calling an ambulance, starting CPR and using an AED. The number of steps has been reduced to focus on the key actions. The intent of the revised algorithm is to present the steps in a logical and concise manner that is easy for all types of rescuers to learn, remember and perform CPR and use an AED.

Figure 2. Adult basic life support algorithm

Table 1. BLS/AED detailed sequence of steps

Initial Assessment

For clarity, the algorithm is presented as a linear sequence of steps. It is recognised that the early steps of ensuring the scene is safe, checking for a response, opening the airway, checking for breathing and calling the ambulance may be accomplished simultaneously or in rapid succession.

Airway

Open the airway using the head tilt and chin lift technique whilst assessing whether the person is breathing normally. Do not delay assessment by checking for obstructions in the airway. The jaw thrust and finger sweep are not recommended for the lay provider.

Dial 999

Early contact with the ambulance service will facilitate dispatcher assistance in the recognition of cardiac arrest, telephone instruction on how to perform CPR and locating and dispatching the nearest AED.

If possible, stay with the victim while calling the ambulance. If the phone has a speaker facility, switch it to speaker mode as this will facilitate continuous dialogue with the dispatcher including (if required) CPR instructions. It seems reasonable that CPR training should include how to activate the speaker phone.

Additional bystanders may be used to call the ambulance service.

Circulation

In adults needing CPR, there is a high probability of a primary cardiac cause for their cardiac arrest. When blood flow stops after cardiac arrest, the blood in the lungs and arterial system remains oxygenated for some minutes. To emphasise the priority of chest compressions, start CPR with chest compressions rather than initial ventilations.

Deliver compressions ‘in the centre of the chest’

Experimental studies show better haemodynamic responses when chest compressions are performed on the lower half of the sternum. Teach this location simply, such as, “place the heel of your hand in the centre of the chest with the other hand on top”. Accompany this instruction by a demonstration of placing the hands on the lower half of the sternum.

Chest compressions are most easily delivered by a single CPR provider kneeling by the side of the victim, as this facilitates movement between compressions and ventilations with minimal interruptions. Over-the-head CPR for single CPR providers and straddle-CPR for two CPR providers may be considered when it is not possible to perform compressions from the side, for example when the victim is in a confined space.

Compress the chest to a depth of 5–6 cm

Fear of doing harm, fatigue and limited muscle strength frequently result in CPR providers compressing the chest less deeply than recommended. Four observational studies, published after the 2010 Guidelines, suggest that a compression depth range of 4.5–5.5 cm in adults leads to better outcomes than all other compression depths during manual CPR. Resuscitation Council UK endorses the ILCOR recommendation that it is reasonable to aim for a chest compression depth of approximately 5 cm but not more than 6 cm in the average sized adult. In making this recommendation, Resuscitation Council UK recognises that it can be difficult to estimate chest compression depth and that compressions that are too shallow are more harmful than compressions that are too deep. Training should continue to prioritise achieving adequate compression depth.

Compress the chest at a rate of 100–120 per minute (min⁻¹)

Two studies, with a total of 13,469 patients, found higher survival among patients who received chest compressions at a rate of 100–120 min⁻¹.⁶ Very high chest compression rates were associated with declining chest compression depths.^{39,40} Resuscitation Council UK therefore recommends that chest compressions are performed at a rate of 100–120 min⁻¹.

Circulation

Minimise pauses in chest compressions

Delivery of rescue breaths, defibrillation shocks, ventilations and rhythm analysis lead to pauses in chest compressions. Pre- and post-shock pauses of less than 10 seconds, and minimising interruptions in chest compressions are associated with improved outcomes.⁴¹⁻⁴⁵ Pauses in chest compressions should be minimised and training should emphasise the importance of close co-operation between CPR providers to achieve this.

Chest recoil

Leaning on the chest preventing full chest wall recoil is common during CPR.

Allowing complete recoil of the chest after each compression results in better venous return to the chest and may improve the effectiveness of CPR. CPR providers should, therefore, take care to avoid leaning forward after each chest compression.

Duty cycle

The proportion of a chest compression spent in compression compared to relaxation is referred to as the duty cycle. There is very little evidence to recommend any specific duty cycle and, therefore, insufficient new evidence to prompt a change from the currently recommended ratio of 50%.

Feedback on compression technique

CPR feedback and prompt devices (e.g. voice prompts, metronomes, visual dials, numerical displays, waveforms, verbal prompts, and visual alarms) should be used when possible during CPR training. Their use during clinical practice should be integrated with comprehensive CPR quality improvement initiatives rather than as an isolated intervention.

CPR provider fatigue

Chest compression depth can decrease as soon as two minutes after starting chest compressions. If there are sufficiently trained CPR providers, they should change over approximately every two minutes to prevent a decrease in compression quality. Changing CPR providers should not interrupt chest compressions.

Rescue breaths

CPR providers should give rescue breaths with an inflation duration of 1 second and provide sufficient volume to make the victim's chest rise. Avoid rapid or forceful breaths. The maximum interruption in chest compression to give two breaths should not exceed 10 seconds.

Mouth-to-nose ventilation

Mouth-to-nose ventilation is an acceptable alternative to mouth-to-mouth ventilation. It may be considered if the victim's mouth is seriously injured or cannot be opened, the CPR provider is assisting a victim in the water, or a mouth-to-mouth seal is difficult to achieve.

Mouth-to-tracheostomy ventilation

Mouth-to-tracheostomy ventilation may be used for a victim with a tracheostomy tube or tracheal stoma who requires rescue breathing.

Barrier devices for use with rescue breaths

Barrier devices decrease transmission of bacteria during rescue breathing in controlled laboratory settings. Their effectiveness in clinical practice is unknown.

If a barrier device is used, care should be taken to avoid unnecessary interruptions in CPR. Manikin studies indicate that the quality of CPR is improved when a pocket mask is used, compared to a bag-mask or simple face shield during basic life support.

Compression-only CPR

CPR providers trained and able to perform rescue breaths should perform chest compressions and rescue breaths as this may provide additional benefit for children and those who sustain an asphyxial cardiac arrest or where the EMS response interval is prolonged.⁵⁴⁻⁵⁷ Only if rescuers are unable to give rescue breaths should they do compression-only CPR.

Resuscitation Council UK has carefully considered the balance between potential benefit and harm from compression-only CPR compared to standard CPR that includes ventilation. Our confidence in the equivalence between compression-only and standard CPR is not sufficient to change current practice.

Resuscitation Council UK, therefore, endorses the ILCOR and ERC recommendations that CPR providers should perform chest compressions for all patients in cardiac arrest. CPR providers trained and able to perform rescue breaths should perform chest compressions and rescue breaths as this may provide additional benefit for children and those who sustain an asphyxia cardiac arrest or where the EMS response interval is prolonged.

When an untrained bystander dials 999, the ambulance dispatcher should instruct them to give chest-compression-only CPR while awaiting the arrival of trained help. Further guidance on dispatcher-assisted CPR is given in the Prehospital resuscitation guidelines.

Use of an automated external defibrillator

AEDs are safe and effective when used by laypeople, including if they have had minimal or no training.⁵⁸ AEDs may make it possible to defibrillate many minutes before professional help arrives. CPR providers should continue CPR with minimal interruption to chest compressions both while attaching an AED and during its use. CPR providers should concentrate on following the voice prompts, particularly when instructed to resume CPR, and minimising interruptions in chest compression.

Compression-only CPR

Public access AED programmes should be actively implemented in public places with a high density and movement of people such as airports, railway stations, bus terminals, sport facilities, shopping malls, stadiums, centres, offices, and casinos – where cardiac arrests are frequently witnessed and trained CPR providers can quickly be on scene. AEDs should also be provided in remote locations where an emergency ambulance response would be likely to be delayed (e.g. aircraft, ferries and off-shore locations). The potential benefits of AEDs being placed in schools as a method to raise awareness and familiarity with this lifesaving equipment is highlighted in the Education and implementation of resuscitation section.

Registration of defibrillators with the local ambulance services is highly desirable so that dispatchers can direct CPR providers to the nearest AED.

Registration of defibrillators with the local ambulance services is highly desirable so that dispatchers can direct CPR providers to the nearest AED.

When implementing an AED programme, community and programme leaders should consider factors such as the development of a team with responsibility for monitoring and maintaining the devices, training and retraining individuals who are likely to use the AED, and identification of a group of volunteer individuals who are committed to using the AED in victims of cardiac arrest. Funds must be allocated on a permanent basis to maintain the programme.

Resuscitation Council UK and British Heart Foundation have produced information endorsed by the National Ambulance Service Medical Directors

Group about AEDs and how they can be deployed in the community – A guide to Automated External Defibrillators.

Risks to recipients of CPR

It is extremely rare for bystander CPR to cause serious harm in victims who are eventually found not to be in cardiac arrest. Those who are in cardiac arrest and exposed to longer durations of CPR are likely to sustain rib and sternal fractures.

Damage to internal organs can occur but is rare. The balance of benefits from bystander CPR far outweighs the risks. CPR providers should not, therefore, be reluctant to start CPR because of the concern of causing harm.


Risks to the CPR provider

CPR training and actual performance is safe in most circumstances. Although rare occurrences of muscle strain, back symptoms, shortness of breath, hyperventilation, pneumothorax, chest pain, myocardial infarction and nerve injury have been described in rescuers, the incidence of these events is extremely low. Individuals undertaking CPR training should be advised of the nature and extent of the physical activity required during the training programme. Learners and CPR providers who develop significant symptoms (e.g. chest pain or severe shortness of breath) during CPR training should be advised to stop and seek medical attention.

Although injury to the CPR provider from a defibrillator shock is extremely rare, standard surgical or clinical gloves do not provide adequate electrical protection. CPR providers, therefore, should not continue manual chest compressions during shock delivery. Avoid direct contact between the CPR provider and the victim when defibrillation is performed. Implantable cardioverter defibrillators (ICDs) can discharge without warning during CPR and rescuers may therefore be in contact with the patient when this occurs. However the current reaching the rescuer from the ICD is minimal and harm to the rescuer is unlikely.

Adverse psychological effects after performing CPR are relatively rare. If symptoms do occur the CPR provider should consult their general practitioner.

Adult BLS



SEQUENCE	PRACTICAL PROCESS *** Please read below then watch the video
SAFETY	Make sure you, the victim and any bystanders are safe
RESPONSE	<p>Check the victim for a response Gently shake their shoulders and ask loudly: "Are you all right?" If they respond, leave them in the position in which you find them, provided there is no further danger; try to find out what is wrong with the person and get help if needed; reassess them regularly</p>
AIRWAY	<p>Open the airway Turn the victim onto their back Place your hand on their forehead and gently tilt their head back; with your fingertips under the point of the victim's chin, lift the chin to open the airway</p>
BREATHING	<p>Look, listen and feel for normal breathing for no more than 10 seconds In the first few minutes after cardiac arrest, a victim may be barely breathing, or taking infrequent, slow and noisy gasps. Do not confuse this with normal breathing. If you have any doubt whether breathing is normal, act as if they are not breathing normally and prepare to start CPR</p>
DIAL	<p>If you are in General Practice Assess patient, if they are unresponsive, ensure that you call for help before initiating CPR. There is usually a green button that you click on your system which alerts staff of which room and where in the practice you are based. It is imperative that someone knows you need help with the patient. When calling 999, state where you are in the surgery. In the event that the paramedics turn up, they should not be wasting time trying to find you.</p>
SEND FOR AED	<p>Send someone to get an AED if available If you are on your own, do not leave the victim, start CPR</p>
CIRCULATION	<p>Start chest compressions Kneel by the side of the victim Place the heel of one hand in the centre of the victim's chest (which is the lower half of the victim's breastbone (sternum)) Place the heel of your other hand on top of the first hand Interlock the fingers of your hands and ensure that pressure is not applied over the victim's ribs Keep your arms straight Do not apply any pressure over the upper abdomen or the bottom end of the bony sternum (breastbone) Position your shoulders vertically above the victim's chest and press down on the sternum to a depth of 5–6 cm After each compression, release all the pressure on the chest without losing contact between your hands and the sternum; Repeat at a rate of 100–120 min-1</p>

GIVE RESCUE BREATHS	<p>After 30 compressions open the airway again using head tilt and chin lift and give 2 rescue breaths</p> <p>Pinch the soft part of the nose closed, using the index finger and thumb of your hand on the forehead</p> <p>Allow the mouth to open, but maintain chin lift</p> <p>Take a normal breath and place your lips around their mouth, making sure that you have a good seal</p> <p>Blow steadily into the mouth while watching for the chest to rise, taking about 1 second as in normal breathing; this is an effective rescue breath</p> <p>Maintaining head tilt and chin lift, take your mouth away from the victim and watch for the chest to fall as air comes out</p> <p>Take another normal breath and blow into the victim's mouth once more to achieve a total of two effective rescue breaths. Do not interrupt compressions by more than 10 seconds to deliver two breaths. Then return your hands without delay to the correct position on the sternum and give a further 30 chest compressions</p> <p>Continue with chest compressions and rescue breaths in a ratio of 30:2</p> <p>If you are untrained or unable to do rescue breaths, give chest compression only CPR (i.e. continuous compressions at a rate of at least 100–120 min⁻¹)</p>
----------------------------	--



AED



— BELMATT —
HEALTHCARE TRAINING



Use of AED

USE OF AED	<p>Switch on the AED</p> <p>Attach the electrode pads on the victim's bare chest</p> <p>If more than one rescuer is present, CPR should be continued while electrode pads are being attached to the chest</p> <p>Follow the spoken/visual directions</p> <p>Ensure that nobody is touching the victim while the AED is analysing the rhythm</p> <p>If a shock is indicated, deliver shock</p> <p>Ensure that nobody is touching the victim</p> <p>Push shock button as directed (fully automatic AEDs will deliver the shock automatically)</p> <p>Immediately restart CPR at a ratio of 30:2</p> <p>Continue as directed by the voice/visual prompts</p> <p>If no shock is indicated, continue CPR</p> <p>Immediately resume CPR</p> <p>Continue as directed by the voice/visual prompts</p>
CONTINUE CPR	<p>Do not interrupt resuscitation until:</p> <p>A health professional tells you to stop</p> <p>You become exhausted</p> <p>The victim is definitely waking up, moving, opening eyes and breathing normally</p> <p>It is rare for CPR alone to restart the heart. Unless you are certain the person has recovered, continue CPR</p>




RECOVERY POSITION



— BELMATT —
HEALTHCARE TRAINING

The Recovery Position

THE RECOVERY POSITION	<p>If you are certain the victim is breathing normally but is still unresponsive, place in the recovery position</p> <p>Remove the victim's glasses, if worn</p> <p>Kneel beside the victim and make sure that both their legs are straight</p> <p>Place the arm nearest to you out at right angles to his body, elbow bent with the hand palm-up</p> <p>Bring the far arm across the chest, and hold the back of the hand against the victim's cheek nearest to you</p> <p>With your other hand, grasp the far leg just above the knee and pull it up, keeping the foot on the ground</p> <p>Keeping their hand pressed against his cheek, pull on the far leg to roll the victim towards you on to their side</p> <p>Adjust the upper leg so that both the hip and knee are bent at right angles</p> <p>Tilt the head back to make sure that the airway remains open</p> <p>If necessary, adjust the hand under the cheek to keep the head tilted and facing downwards to allow liquid material to drain from the mouth</p> <p>Check breathing regularly</p> <p>**** Please see recovery position video</p> <p>Be prepared to restart CPR immediately if the victim deteriorates or stops breathing normally</p>
------------------------------	--



CHOKING



— BELMATT —
HEALTHCARE TRAINING

Choking

Choking is an uncommon but potentially treatable cause of accidental death. As most choking events are associated with eating, they are commonly witnessed.


As victims are initially conscious and responsive, early interventions can be lifesaving.

Recognition

Recognition of airway obstruction is the key to successful outcome, so do not confuse this emergency with fainting, myocardial infarction, seizure or other conditions that may cause sudden respiratory distress, cyanosis or loss of consciousness. Choking usually occurs while the victim is eating or drinking.

People at increased risk of choking include those with reduced consciousness, drug and/or alcohol intoxication, neurological impairment with reduced swallowing and cough reflexes (e.g. stroke, Parkinson's disease), respiratory disease, mental impairment, dementia, poor dentition and older age.

Table 2 and Figure 3 present the treatment for the adult with choking. Foreign bodies may cause either mild or severe airway obstruction. It is important to ask the conscious victim "Are you choking?" The victim that is able to speak, cough and breathe has mild obstruction. The victim that is unable to speak, has a weakening cough, is struggling or unable to breathe, has severe airway obstruction.






Table 2: Sequence of steps for managing the adult victim who is choking

SEQUENCE	PRACTICAL PROCESS
SUSPECT CHOKING	Be alert to choking particularly if victim is eating
ENCOURAGE TO COUGH	Instruct victim to cough
GIVE BACK BLOWS	If cough becomes ineffective give up to 5 back blows <ul style="list-style-type: none"> • Stand to the side and slightly behind the victim • Support the chest with one hand and lean the victim well forwards so that when the obstructing object is dislodged it comes out of the mouth rather than goes further down the airway • Give five sharp blows between the shoulder blades with the heel of your other hand
GIVE ABDOMINAL THRUSTS	If back blows are ineffective give up to 5 abdominal thrusts <ul style="list-style-type: none"> • Stand behind the victim and put both arms round the upper part of the abdomen • Lean the victim forwards • Clench your fist and place it between the umbilicus (navel) and the ribcage • Grasp this hand with your other hand and pull sharply inwards and upwards • Repeat up to five times • If the obstruction is still not relieved, continue alternating five back blows with five abdominal thrusts
START CPR	Start CPR if the victim becomes unresponsive <ul style="list-style-type: none"> • Support the victim carefully to the ground • Immediately activate the ambulance service • Begin CPR with chest compressions

Treatment for mild airway obstruction

Coughing generates high and sustained airway pressures and may expel the foreign body. Aggressive treatment with back blows, abdominal thrusts and chest compressions at this stage may cause harm and can worsen the airway obstruction. These treatments are reserved for victims who have signs of severe airway obstruction. Victims with mild airway obstruction should remain under continuous observation until they improve, as severe airway obstruction may subsequently develop.

Treatment for severe airway obstruction

The clinical data on choking is largely retrospective and anecdotal. For conscious adults and children over one year of age with complete airway obstruction, case reports show the effectiveness of back blows or 'slaps' and abdominal thrusts.

Approximately half of cases of airway obstruction are not relieved by a single technique. The likelihood of success is increased when combinations of back blows or slaps, and abdominal and chest thrusts are used.

Treatment of choking in an unresponsive victim

Higher airway pressures can be generated using chest thrusts compared with abdominal thrusts. Bystander initiation of chest compressions for unresponsive or unconscious victims of choking is associated with improved outcomes. Therefore, start chest compressions promptly if the victim becomes unresponsive or unconscious. After 30 compressions, attempt 2 rescue breaths, and continue CPR until the victim recovers and starts to breathe normally.

If you perform an abdominal thrust then the patient must be sent to hospital for evaluation

Aftercare and referral for medical review

- Following successful treatment of choking, foreign material may nevertheless remain in the upper or lower airways and cause complications later.
- Victims with a persistent cough, difficulty swallowing or the sensation of an object being still stuck in the throat should, therefore, seek medical advice.
- Abdominal thrusts and chest compressions can potentially cause serious internal injuries and all victims successfully treated with these measures should be examined afterwards for injury.
- Patients receiving antiplatelet and/or anticoagulant drugs are at increased risk of intra-abdominal haemorrhage and we suggest a low threshold for obtaining a senior clinical opinion and thoracoabdominal CT scan if a thoracoabdominal injury is suspected.

Resuscitation of children and victims of drowning

Many children do not receive resuscitation because potential CPR providers fear causing harm if they are not specifically trained in resuscitation for children. This fear is unfounded: it is far better to use the adult BLS sequence for resuscitation of a child than to do nothing. For ease of teaching and retention, laypeople are taught that the adult sequence may also be used for children who are not responsive and not breathing normally. The following minor modifications to the adult sequence will make it even more suitable for use in children:

- Give 5 initial rescue breaths before starting chest compressions.
- If you are on your own, perform CPR for 1 minute before going for help.
- Compress the chest by at least one third of its depth, approximately 4 cm for the infant and approximately 5 cm for an older child. Use two fingers for an infant under 1 year; use one or two hands as needed for a child over 1 year to achieve an adequate depth of compression.

The same modifications of 5 initial breaths and 1 minute of CPR by the lone CPR provider before getting help may improve outcome for victims of drowning. This modification should be taught only to those who have a specific duty of care to potential drowning victims (e.g. lifeguards).

BELMATT HEALTHCARE TRAINING



— B E L M A T T —
HEALTHCARE TRAINING

Contact Us



PHONE CALL

0207 692 8709



FOLLOW US

Belmatt Healthcare
Training



EMAIL US

info@belmatt.co.uk

Belmatt Healthcare Training Limited, provider of postgraduate training.

Registered Office: Suite 570 405 Kings Road, Chelsea, London, SW10 0BB