AQI ManuscriptTables

Michael Andreae

December 3, 2015

Prepare

##

Load Data to produce table

Load the complete case dataset produced with AQI_Import on 3Dec2015 once from the source data file and save it in /Data and clean workspace.

```
'data.frame':
                    92683 obs. of 19 variables:
##
               : Factor w/ 2 levels "no Ondan", "Ondan": 1 2 2 2 1 2 2 2 1 ...
               : logi FALSE TRUE TRUE TRUE FALSE TRUE ...
##
   $ ondL
               : Factor w/ 2 levels "no Dex", "Dex": 1 1 2 1 1 2 2 2 2 1 ...
   $ dexa
##
   $ dexL
               : logi FALSE FALSE TRUE FALSE FALSE TRUE ...
               : Factor w/ 2 levels "no Drope", "Drope": 1 1 1 1 1 1 1 1 1 1 ...
##
   $ dro
##
   $ droL
               : logi FALSE FALSE FALSE FALSE FALSE ...
               : Factor w/ 2 levels "neither", "either": 1 2 2 2 1 2 2 2 1 ...
##
   $ any
               : logi FALSE TRUE TRUE TRUE FALSE TRUE ...
##
   $ either
##
   $ pay
               : Factor w/ 4 levels "Commercial", "MEDICAID", ...: 3 1 3 3 3 1 2 1 3 1 ...
##
               : num 0.44 0.68 0.45 0.44 1.05 1.07 0.34 0.83 0.41 0.32 ...
   $ income
##
   $ incomeQ
              : Factor w/ 4 levels "very low", "low", ...: 2 3 2 2 4 4 1 3 2 1 ....
               : int 43 39 68 72 73 59 35 49 77 57 ...
##
   $ age
##
   $ age_group: Factor w/ 6 levels "19 - 49", "Under 1",..: 1 1 5 5 5 4 1 1 5 4 ...
##
               : Factor w/ 2 levels "female", "male": 2 2 1 2 1 2 2 1 2 1 ...
##
               : Factor w/ 4333 levels "0182T", "0192T", ...: 2864 3553 3752 3075 1550 3580 2569 2549 3754
   $ cpt
               : Factor w/ 5 levels "1","2","3","4",..: 5 3 2 3 4 2 3 2 3 4 ...
##
   $ ASA
  $ anes_type: Factor w/ 4 levels "General", "Neuroaxial",..: 1 1 1 1 1 1 1 1 1 1 ...
  $ practice : Factor w/ 6 levels "A","B","C","D",..: 6 6 6 6 6 6 6 6 6 ...
               : Factor w/ 1208 levels "5622", "5623",...: 50 17 4 21 27 1 5 91 71 77 ...
   $ prov
```

Build Tables

Table 1 Population characteristics

Table 1 Population characteristics describes the subpopulation in the NACOR database with information on antiemetic administration. We found statistically significant differences (p<0.001) in all comparisons (age, gender, income in the patients zip code and case characteristics like outpatient procedure, emergency procedure and case duration.

Groups	Variables	Total
Population	n	441645
Gender	female	249909
	male	191654
Age	median [IQR)	52
	19 - 49	159002
	Under 1	6357

Groups	Variables	Total
	1-18	35881
	50 - 64	116869
	65 - 79	96274
	80+	27252
Insurance	Commercial	97443
	MEDICAID	25865
	Medicare	51441
	SELF	1585
Income	income	1
	very low	2
	low	3
	middle	4
	high	5