

Anesthesia & Analgesia

A Novel Approach to Synthesize the Evidence on Analgesic Adjuvants for Postoperative Pain --Manuscript Draft--

Manuscript Number:	
Full Title:	A Novel Approach to Synthesize the Evidence on Analgesic Adjuvants for Postoperative Pain
Short Title:	A Novel Approach to Synthesize the Evidence
Article Type:	Invited Editorial
Corresponding Author:	Michael H Andrae Penn State Milton S Hershey Medical Center Hershey, Pennsylvania UNITED STATES
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Penn State Milton S Hershey Medical Center
Corresponding Author's Secondary Institution:	
First Author:	Michael H Andrae, M.D.
First Author Secondary Information:	
Order of Authors:	Michael H Andrae, M.D.
	Nathan L Pace, MD., MStat.
Order of Authors Secondary Information:	
Manuscript Region of Origin:	UNITED STATES
Requested Editor:	Honorio T. Benzon, M.D.

A Novel Approach to Synthesize the Evidence on Analgesic Adjuvants for Postoperative Pain

Authors

Michael H. Andreae, M.D., Department of Anesthesiology & Perioperative Medicine,
H187, Penn State Health Milton S. Hershey Medical Center, Penn State College of
Medicine, Hershey. PA

Nathan L Pace, MD., MStat., Department of Anesthesiology, University of Utah, Salt
Lake City, UT

Corresponding author:

Michael Andreae, MD, Address: Department of Anesthesiology, Department of
Anesthesiology, Penn State Health Milton S. Hershey Medical Center, 500 University
Drive, Hershey. PA 17033, Tel: +1 (717) 531 6140, Email:
mandreae@pennstatehealth.psu.edu

Acknowledgement

We would like to acknowledge Doleman et al for providing the data to generate the
figures.

Conflicts of Interest: None

Number of words in this Editorial (1598), Figures (2)

Keywords:

analgesic adjuvants, postoperative pain, meta-analysis, Bayesian statistics

Editorial Text

Based on the best available evidence, which analgesic adjuvant is best for a particular patient to optimize postoperative pain control? For example, how much opioid sparing will the administration of intraoperative intravenous acetaminophen afford for a typical bariatric patient at Penn State Hershey or University of Utah Medical Centers? With patients and surgical procedures across the United States as diverse as species in the rain forest, this can be a difficult question. Often randomized controlled trial yield varying estimates of the clinical effectiveness of analgesic adjuvants. RCTs results, seemingly at random, leave clinicians baffled. Doleman et al in the study “Baseline morphine consumption...” propose an ingenious new solution to solve this riddle¹. They synthesized the evidence for morphine dose reduction with adjuvants by controlling for baseline risk (morphine consumption) across surgical procedures and patient populations. With their novel approach, local audit data could be used to predict the expected average reduction in morphine consumption for any analgesic adjuvant. In the same breath, they modify and perhaps toss out the established paradigm of procedure-specific pain control.

Some variability in outcome estimates from RCTs investigating the same intervention in the similar population is to be expected. Each RCT recruits patients by a convenience sample; it is not a random sample of the entire population who might receive the adjuvant. By pure chance alone, each RCT has a chance to over- or to under estimate the effect of an intervention. Also, larger and smaller studies will lead to more or less precise estimates of effect of interventions. Meta-analysis pools available RCTs to synthesize the evidence and to increase the precision of effect estimates. This may

1
2
3
4 reduce the uncertainty in the face of seemingly contradictory results. However, if the
5
6 heterogeneity of the RCTs is very large (too inconsistent and disparate), then these
7
8 summary statistics may be inappropriate and are frequently described as mixing apples
9
10 and oranges. Excessive between-study heterogeneity in meta-analysis raises concerns
11
12 that the included studies are clinically and methodologically too different, making
13
14 pooling all identified RCTs unreasonable.
15
16
17

18
19 To explain why RCTs yield contradictory results, we can group studies by surgical
20
21 intervention. Stratification by surgical intervention makes clinical sense because
22
23 between-study heterogeneity may be smaller within each stratum. This led to the
24
25 current paradigm of procedure-specific pain control²: It is expected that different surgical
26
27 procedures cause different amounts of pain. Populations undergoing different
28
29 interventions for different diseases may vary in how they respond to pain, in their
30
31 comorbidities, pharmacokinetics etc. Clearly, a population of elderly men following
32
33 thoracotomy for lung cancer will differ from a population of young women after cesarean
34
35 section. It follows that postoperative pain control should be tailored to the specific
36
37 surgical intervention and the particular population. While this seems intuitive, it
38
39 drastically reduces the number of available studies for clinical decision making on for a
40
41 particular patient population undergoing a particular procedure. For example, on
42
43 thyroidectomy Doleman et al.¹ found only one single RCT investigating the effect of
44
45 intravenous acetaminophen. This poses a significant challenge to any evidence-based
46
47 approach to procedure-specific postoperative pain management.
48
49
50
51
52
53
54
55

56 For studies investigating adjuvants for improved postoperative pain control, the effect
57
58 estimate is the mean difference in morphine dose between those receiving and those
59
60
61
62
63
64
65

not receiving the adjuvant. The mean dose of those not receiving the adjuvant can be considered to be the baseline risk for pain. Doleman et al.¹ used Bayesian hierarchical meta-regression modeling adapted to requirements of such evidence synthesis³. They demonstrated that surgical procedure is not the main predictor of between-study variability in effect estimates for analgesic adjuvants to control postoperative pain, but baseline risk is. Restated, the benefit of the analgesic adjuvant is mainly driven by baseline morphine consumption in the control group and not by the surgical procedure. Two figures using acetaminophen data as an example will illustrate their findings.

Doleman et al.¹ found 25 RCTs investigating acetaminophen for postoperative pain control. In Figure 1, a classical forest plot is shown, ordered by surgical interventions.* The 25 RCTs have different effect sizes; the results are inconsistent, sometimes varying widely even within the same surgery.

In Figure 2, a schematic of the acetaminophen subplot of Doleman's Figure 2¹ is rendered.♦ Each study is represented by a colored icon. If surgery were the best way to explain why studies yield different results, then studies investigating the same surgical procedure should have similar results. Hence the same color icons should be grouped together (along the same value on the y-axis indicating the estimated effect). But studies in the same surgical group are far apart on the y-axis. By contrast, studies with similar baseline risk (close on the x-axis, i.e. baseline morphine consumption in the

* The 95% confidence intervals provided by Doleman et al.¹ were used to estimate the standard error.⁴ The figure was generated with the Cochrane Collaboration software RevMan⁵.

♦ This figure was generated by a R⁶ statistical software package with data provided by the authors¹.

control group) tend to have similar effects, (they are close on the y-axis, i.e. mean morphine-equivalent reduction), illustrated in a neat alignment along the regression line in Figure 2. The conclusion is that a meta-regression controlling for baseline morphine consumption in the control group is superior to stratification by surgery in explaining between study variance in results.

Doleman et al.¹ propose that this is true for all analgesic adjuvants across all populations, for all surgical interventions, as shown in the other subplots of Doleman's Figure 2¹. This contradicts the dominant paradigm of procedure-specific postoperative pain control. What is more, Doleman et al.¹ also propose that clinicians can use local audit data, collected on local surgical populations, to estimate the average reduction of morphine consumption for each adjuvant. This is a great feature to have for a new unifying theory on how to synthesize the evidence for postoperative pain control.

Doleman et al.¹ fit both a classical model and a Bayesian model that differs from the classical approach (also known as frequentist statistics). The Bayesian approach incorporates prior information and combines it with the newly observed data, much like a physician would in clinical practice. Both frequentist and Bayesian statistical models are based on subjective assumptions; the model choice itself is often more important for correct inferences. In both types of modeling, the analyst should explore the sensitivity of results and inferences to assumptions and model choices. In an available supplement, Doleman et al.¹ provided software code, data and model details. They used very weak prior assumptions and in their sensitivity analysis found their results to be robust. They made their Bayesian approach transparent, sound and trustworthy.

1
2
3
4 Based on their previous work⁷, they speculated that baseline risk might better explain
5
6 variability. Baseline risk for pain may modify the treatment effect of analgesic adjuvants
7
8 as a proxy for unmeasured patient-level characteristics⁷. Their conjecture is supported
9
10 by their meta analyses. This now suggests a set of testable hypotheses and predictions,
11
12 making their proposed novel and unifying theory attractive. Progress in science and
13
14 medicine is made by proposing models and hypotheses and then attempting to falsify
15
16 these. Indeed, investigators can use local audit data to predict the reduction in morphine
17
18 requirements and then prospectively observe the actual reduction achieved to
19
20 investigate how well the Doleman model predictions hold. Meta-analysts can replicate
21
22 the approach by Doleman et al.¹ and for example test if it can be extended to evidence
23
24 synthesis of regional anesthesia to explain and/or reduce the between study variability
25
26 of the effects of regional anesthesia versus conventional multimodal pain control for
27
28 postoperative pain control. If Doleman's hypothesis holds, it would greatly simplify
29
30 evidence synthesis and provide clinicians with much stronger and much more granular
31
32 evidence on what analgesic to use for which patient. In fact, in the second part of their
33
34 paper, Doleman et al.¹ go on to use meta-regression to discuss rather refined and
35
36 specific questions. For example, the question, if acetaminophen is more effective if
37
38 administered intravenously rather than orally, is of great financial and clinical
39
40 importance. But there are simply too few studies to address this question, if we insisted
41
42 on combining only studies comparing the effects in the same population/surgical
43
44 intervention. Doleman et al.¹ provide a league table of efficacy of analgesic adjuvants.
45
46
47 Doleman et al.¹ acknowledge the many limitations of their work. Their meta-regression,
48
49 like any meta-analysis was an observational study, even if it had an a priori defined
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

analysis plan. The confidence of their predictions and inferences will be weaker for ranges of baseline risk (baseline morphine consumption in the control group) where they have less studies for the particular analgesic adjuvant at hand. We will list these limitations using acetaminophen as an example:

- The range of surgical procedures included in the acetaminophen trials was limited, (see Doleman's Table 1¹).
- The patient mix in the acetaminophen studies are predominantly female.
- The standard morphine dose (50 mg) at which dose reduction was estimated is much higher than the mean dose (28 mg) in the included acetaminophen studies. Inferences and predictions in the higher dose ranges therefore rely on fewer studies and/or on problematic extrapolation.
- For most adjuvants studied, there was little benefit (almost equal benefit) for studies reporting low baseline consumption, which is evidenced in Figure 2 of Doleman's article¹.
- Finally, the estimates in their regression equation apply to aggregate groups not individual patients.

We and the reader may remain somewhat skeptical and cautious about the approach and inferences suggested by Doleman et al.¹ Provocative as the novel hypothesis by Doleman et al.¹ may be, it would afford great utility. It also exemplifies the benefit and potential of novel Bayesian meta-regression approaches to support clinical decision making.

Figures

Figure 1




Figure 1

Figure 1 Caption

Estimates for the mean reduction of morphine equivalent afforded by acetaminophen are shown in a classical forest plot with studies grouped by surgical interventions. The mean reduction of morphine equivalents and their 95% confidence intervals were provided by Doleman et al.. The later were used to estimate the standard error (Higgins

2011). The figure was produced with the Cochrane Collaboration software RevMan (RevMan 2014). The 25 RCTs have inconsistent effects for the drug under investigation (acetaminophen), varying widely even within the same surgery.

Figure 2




Figure 2 Caption

In Figure 2, we reproduce the acetaminophen subplot of Figure 2 of Doleman et al. Each study is represented by a dot. We coded and colored the studies by surgical procedure, according to the adjacent legend. If surgery were the best way to explain why studies yield different results, then studies investigating the same surgical procedure should have similar effects. Hence the same shape (and color) dots should be clumped together, (and the green-square-box cholecystectomy studies somewhat are). But most studies of the same color are far apart, for example orthopedic (violet-star) or hysterectomy (blue-cross-in-box) study dots are spread out over the entire range of the plot. By contrast, most studies line up neatly around the blue regression

1
2
3
4 line, when we regress the mean reduction in morphine consumption (y-axis) against the
5
6 baseline risk (baseline morphine consumption in the control group) in the x-axis. The
7
8 conclusion is that a meta-regression controlling for baseline morphine consumption in
9
10 the control group is superior to stratification by surgery in explaining between study
11
12 variance in results.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. Doleman B, Sutton A, Sherwin M, Lund J, J W. Baseline morphine consumption may explain between-study heterogeneity in meta-analyses of adjuvant analgesics and improve precision and accuracy of effect estimates. *Anesthesia and Analgesia* 2017.
2. Kehlet H, Wilkinson RC, Fischer HBJ, Camu F, Prospect Working Group. PROSPECT: evidence-based, procedure-specific postoperative pain management. *Best practice & research Clinical anaesthesiology* 2007;21:149–59. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17489225>.
3. Sutton AJ, Abrams KR. Bayesian methods in meta-analysis and evidence synthesis. *Statistical methods in medical research* 2001;10:277–303. Available at: <http://journals.sagepub.com/doi/10.1177/096228020101000404> <http://www.ncbi.nlm.nih.gov/pubmed/11491414>.
4. Higgins JPT, Deek JJ. Chapter 7: Selecting studies and collecting data. In: Higgins JPT, Green S, eds. *Cochrane handbook for systematic reviews of interventions* 5.1.0 [updated march 2011]. Cochrane Collaboration, 2011:Available from www.cochrane-handbook.org.
5. Review manager (RevMan) [Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
6. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2013. Available at: <http://www.R-project.org>.
7. Achana FA, Cooper NJ, Dias S, Lu G, Rice SJC, Kendrick D, Sutton AJ. Extending methods for investigating the relationship between treatment effect and baseline risk from pairwise meta-analysis to network meta-analysis. *Statistics in Medicine* 2013;32:752–71. Available at: <http://doi.wiley.com/10.1002/sim.5539>.

Figure 1 Forest Plot


[Click here to download Figure AcetaminophenForestplot.png](#)

Figure 2 Schematic

[Click here to download Figure Figure 2 Schematic.png](#)

