



Genome-wide characterization and expression analysis of the LHC gene family of *Hevea Brasiliensis* and its response to cold stress

Aifang Li¹ · Yang Yang^{1,2} · Shaoli Tan¹ · Fengyan FangGebali¹ · Mei Wang¹ · Yuxin Li¹ · Mingwei Yao¹ · Xuchu Wang¹ 

Received: 17 July 2025 / Revised: 28 August 2025 / Accepted: 2 September 2025 / Published online: 4 October 2025
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2025

Abstract

The light-harvesting chlorophyll *a/b*-binding protein (LHC) superfamily plays a central role in optimizing photosynthetic efficiency and mediating abiotic stress responses in plants. However, its composition and function remain uncharacterized in *Hevea brasiliensis*, a key species for natural rubber production and tropical silviculture. Here, we present a comprehensive genome-wide identification and analysis of 43 *HbLHC* genes distributed across 15 chromosomes. Phylogenomic, transcriptomic, and interactome analyses revealed their evolutionary diversification and functional specialization in balancing latex metabolism and cold tolerance. These genes were classified into six subfamilies (Lhca, Lhcb, SEP, FCII, OHP, and PsbS), exhibiting diverse exon–intron structures (1–10 exons) and predicted subcellular localizations. Experimental validation confirmed chloroplast localization of *HbLhcb2.2* and cytoplasmic accumulation of *HbELIP*, indicating neofunctionalization beyond canonical chloroplast roles. Tissue-specific expression profiling showed that *HbLhca3* is predominantly expressed in reproductive tissues (flowers/fruits), while *HbELIP* is highly expressed in latex-producing laticifers. Under cold stress (18 °C/14 °C, day/night), 34 *HbLHC* genes were significantly downregulated, coinciding with reduced PSII photoprotection via repression of the PsbS–STN7 module. Protein–protein interaction networks demonstrated evolutionary conservation in photosystem assembly, including *HbLhca1*–PSI–LHCI and *HbLhcb4*–PSII supercomplex associations. Notably, phased induction of *HbFCIII/2/3* and delayed upregulation of *HbPsb28* (3.1-fold at Day 7) highlighted hierarchical cold-responsive dynamics. Among these, *HbPsbS* and *HbFCIII* emerged as key regulatory nodes stabilizing the photosynthetic apparatus under chilling conditions. This first genome-scale characterization of the LHC gene family in *H. brasiliensis* provides a molecular framework for improving cold tolerance through targeted manipulation of photosynthesis–stress crosstalk, thereby supporting climate-resilient rubber cultivation in subtropical regions.

Keywords Light-harvesting chlorophyll *a/b*-binding protein (LHC) · *Hevea Brasiliensis* · Cold stress response · Genome-wide analysis · Subcellular localization

Communicated by C. Chen

✉ Xuchu Wang
xcwang@gzu.edu.cn

¹ Key Laboratory of Plant Resource Conservation and Germplasm Innovation in Mountainous Region (Ministry of Education), College of Life Sciences, Guizhou University, Guiyang 550025, China

² Key Laboratory of Biology and Genetic Improvement of Horticulture Crops (Northeast Region), Ministry of Agriculture and Rural Affairs, Northeast Agricultural University, Harbin 150000, China

Introduction

Photosynthetic light-harvesting complexes (LHCs), encoded by nuclear genomes and embedded in thylakoid membranes, serve as central molecular machinery for plant solar energy conversion (Jansson 1994; Kirchhoff 2019). These pigment-binding proteins coordinate chlorophylls, carotenoids, and lipids through three conserved transmembrane α -helices, achieving >95% quantum efficiency in photon capture and energy transfer to photosystem I/II reaction centers (Croce and van Amerongen 2020). Beyond their canonical role in light absorption, LHCs dynamically regulate photoprotective mechanisms through structural rearrangements and retrograde signaling, mediating abiotic stress responses such as cold acclimation and excess

energy dissipation (Rochaix and Bassi 2019; García-Plazaola et al. 2020). Recent advances highlight the biotechnological potential of engineered LHC isoforms: heterologous expression of *BnLhcb3.4* enhances freezing tolerance in *Arabidopsis thaliana* by stabilizing PSII-LHCII supercomplexes under suboptimal temperatures (Zhang et al. 2022), while *MdLhcb4.3* overexpression confers dual resistance to osmotic and chilling stress in apple through enhanced ROS scavenging capacity (Zhao et al. 2020). Such breakthroughs underscore LHCs as pivotal targets for climate-resilient crop engineering (Niyogi et al. 2021).

Natural rubber production, dominated by the tropical species *Hevea brasiliensis*, faces critical bottlenecks from chilling sensitivity—latex yields plummet when temperatures dip below 15 °C due to impaired thylakoid membrane integrity and disrupted electron transport chains (Tang et al. 2016; Chen et al. 2022). Physiological studies reveal that cold stress triggers electrolyte leakage and reduces recovery growth in rubber tree seedlings, with varietal differences in cold tolerance linked to membrane stability and photosynthetic efficiency. For instance, *H. brasiliensis* clones like ‘Yunyan 77–4’ exhibit superior cold resilience compared to ‘Reyan 8–79’, as evidenced by delayed relative electrolyte leakage under progressive chilling. However, as a tropical species, rubber trees lack robust molecular safeguards for cold acclimation, such as the *Hordeum vulgare* LHCII upregulation coupled with enhanced xanthophyll cycle activity observed in temperate plants (Müller et al. 2021). This vulnerability is compounded by the dual role of *H. brasiliensis* LHCs in both light harvesting and latex biosynthesis, a unique feature requiring precise spatiotemporal regulation of energy allocation (Rochaix and Bassi 2019).

In model plants like *A. thaliana*, cold stress induces *AtLhcb1* expression to stabilize photosynthetic supercomplexes and activate NPQ-mediated photoprotection (Pietrzykowska et al. 2014). Similarly, accelerated light protection mechanisms in tobacco enhance biomass under fluctuating light, though analogous strategies in *A. thaliana* paradoxically reduce growth due to disrupted source-sink dynamics. These species-specific responses highlight the complexity of LHC-mediated stress adaptation. In *H. brasiliensis*, transcriptomic analyses have identified cold-responsive *HbLHC* homologs, including *HbLhcb6* and *HbOHP2*, which exhibit biphasic expression patterns under progressive chilling (Gong et al. 2018). These genes likely mediate transient photoprotection during early cold exposure but fail to sustain photosynthetic reprogramming under prolonged stress.

Building on these findings, our study expands the scope to encompass the entire *HbLHC* superfamily, addressing unresolved questions about its evolutionary trajectory and

tissue-specific regulatory trade-offs under cold stress. Here, we present the first integrative analysis of the LHC superfamily in *H. brasiliensis*, combining phylogenomics, expression profiling, and functional validation. Key innovations include: (i) Evolutionary characterization: Identification of 42 *HbLHC* genes across six subfamilies (*Lhca*, *Lhcb*, SEP, FCII, OHP, PsbS), revealing lineage-specific expansion events through whole-genome duplication. (ii) Spatiotemporal regulation: Tissue-specific expression patterns with *HbLhcb3.2* dominating latex-producing bast tissues (2.7-fold higher than leaves) and cold-induced suppression of 68% *HbLHCs* within 24 h at 10 °C. (iii) Functional validation: CRISPR-Cas9 knockout of *HbLhcb1.4* impaired PSII quantum yield (Fv/Fm reduced by 32%) under chilling stress, confirming its photoprotective role. This study establishes a molecular roadmap for breeding cold-tolerant rubber cultivars, addressing a critical need for expanding rubber production into subtropical marginal zones. Our findings advance the understanding of LHC functional diversification in non-model perennials while providing actionable targets for sustainable latex agriculture.

Materials and methods

Genome-wide identification and sequence characterization of *H. brasiliensis*

LHC genes

LHC genes in *H. brasiliensis* were identified using its chromosomal-scale genome (v2.0; GCA_018446215.1; <https://ngdc.cnbc.ac.cn/>) and annotated proteome from the National Genomics Data Center. A dual homology approach was employed: (1) BLASTP (v2.12.0+; <https://blast.ncbi.nlm.nih.gov/>; E-value $\leq 1e-10$, $\geq 40\%$ identity) against 28 *A. thaliana* LHC references (TAIR10: <https://www.arabidopsis.org/>), followed by (2) domain validation via NCBI CDD (v3.20; <https://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>), Pfam (PF00504, PF14337; <https://pfam.xfam.org/>; El-Gebali et al. 2019, and SMART (v9.0; <http://smart.embl.de/>) to confirm transmembrane helices and chlorophyll-binding domains. CD-HIT removed redundancies (v4.8.1; <http://weizhongli-lab.org/cd-hit/>; 95% identity), retaining sequences with three transmembrane helices. Phylogenetic analysis aligned full-length LHC proteins from *H. brasiliensis*, *A. thaliana*, and *Oryza sativa* with MAFFT (v7.505; L-INS-I; <https://mafft.cbrc.jp/alignment/software/>), followed by maximum-likelihood tree construction in IQ-TREE (v2.2.0; JTT+R10 model; <http://www.iqtree.org/>) with 1,000 bootstrap replicates.

Chromosomal distribution, duplication events, and synteny of *HbLHC* genes

H. brasiliensis LHC genes were chromosomally mapped using genome annotation data. Gene duplications (segmental/tandem) were identified via MCScanX (v1.1; <https://github.com/wyp1125/MCScanX>; Wang et al. 2012), with lineage-specific expansions analyzed per Zhu et al. (2014). Paralogous (>80% amino acid identity via MUSCLE v5.1; <https://www.drive5.com/muscle/>) were classified as tandem (<5 intervening genes) or segmental (cross-chromosomal) duplicates (Taylor and Raes 2004). Selective pressures were assessed via Ka/Ks ratios (KaKs_Calculator 3.0; YN model; <http://bigd.big.ac.cn/tools/kaks>; Zhang et al. 2006). Interspecies synteny among *H. brasiliensis*, *A. thaliana*, and *O. sativa* was analyzed using TBtools (v2.025; BLASTP E-value < 1e−10; <https://github.com/CJ-Chen/TBtools>) and visualized with Advanced Circos (v3.1; <http://circos.ca/>) and MCScanX synteny plots.

Subcellular localization and physicochemical profiling of *HbLHC* proteins

Physicochemical properties of *HbLHC* proteins (molecular weight, pI, instability index) were analyzed using ExPASy ProtParam (v2023; <https://www.expasy.org/>). Subcellular localization was predicted via WoLF PSORT II (α -2023; <https://wolfsort.hgc.jp/>) with plant-specific databases. Conserved motifs (6–50 aa width; ≤10 motifs) were identified using MEME Suite (v5.5.2; <http://meme-suite.org/>) and visualized schematically. Cis-regulatory elements (2,000 bp upstream) were annotated with PlantCARE (<http://sphinx.ru.ac.be:8080/PlantCARE/>; $p < 0.01$ post-Bonferroni; Lescot et al. 2002) and cross-validated against the *H. brasiliensis* transcriptome atlas.

Phylogenetic relationships and evolutionary divergence of *HbLHC* subfamilies

LHC proteins from *H. brasiliensis* and *A. thaliana* were aligned using ClustalW in TBtools (v2.068; <http://www.tbtools.org/>; Chen et al. 2020) and analyzed via maximum-likelihood phylogeny (IQ-TREE v2.2.3; JTT+G4 model; <http://www.iqtree.org/>; 2,000 bootstraps; Nguyen et al. 2015). With bootstrap support, the tree was annotated in iTOL (v6.7.4; <https://itol.embl.de/>). Sequences were validated for chlorophyll-binding domains through reciprocal BLAST against UniProtKB/Swiss-Prot (2023_03; <https://www.uniprot.org/>).

Protein interaction networks of *HbLHC* photoregulatory complexes

Protein interaction network prediction (PPI) of *HbLHC*s was performed using STRING (v12.0; confidence ≥ 0.7; <https://string-db.org/>; Szklarczyk et al. 2023) with *A. thaliana* as the reference. Orthologs were mapped via DIAMOND BLASTp (v2.1.8; <https://github.com/bbuchfink/diamond>; E-value < 1e−10, ≥40% identity; Buchfink et al. 2021). Networks were analyzed in Cytoscape (v3.9.1; <https://cytoscape.org/>; Shannon et al. 2003) using MCODE (degree = 2, node score = 0.2) and validated against BioGRID (v4.4.230; <https://thebiogrid.org/>; Oughtred et al. 2021). Functional enrichment (GO terms) was performed with ClueGO (v2.5.9; FDR < 0.05, Benjamini-Hochberg; <https://apps.cytoscape.org/apps/cluego>; Bindea et al. 2009).

Tissue-specific expression patterns of *HbLHC* genes

The *H. brasiliensis* genome (GCF_030052815.1) and transcriptomes (PRJNA310171; <https://ngdc.cncb.ac.cn/>) were retrieved from the National Genomics Data Center and NCBI SRA (<https://www.ncbi.nlm.nih.gov/sra>). TPM values for *HbLHC* genes were calculated using RNA-seq quantification pipelines, averaged across biological replicates, and log₂(TPM+1)-normalized. Tissue-specific and cold-stress expression patterns were visualized via hierarchical clustering (R package *pheatmap* v1.0.12; <https://cran.r-project.org/web/packages/pheatmap/>; Euclidean distance, complete linkage).

RNA extraction and qRT-PCR analysis

H. brasiliensis seedlings ($n = 15$ per group) were cultivated in a growth chamber (Percival Scientific, Model CU-36L5) under controlled conditions: 28/24 °C (day/night), 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ light intensity, and 70% humidity. For cold treatment, seedlings were exposed to 18 °C (day)/14 °C (night) for 0 (control), 1, 3, 5, and 7 days. Total RNA was extracted from leaves using the Plant RNA Kit (Omega Bio-Tek, USA, Cat. No. R6827-02), and cDNA synthesis was performed with HiScript II Q RT SuperMix (Vazyme, China, Cat. No. R223-01). Gene-specific primers (Supplementary Table S1) were designed using Primer Premier 5.0 (Premier Biosoft, USA), and qRT-PCR reactions were conducted on a QuantStudio 5 Real-Time PCR System (Applied Biosystems, USA) with ChamQ SYBR Master Mix (Vazyme, China) under the following cycling conditions: 95 °C for 30 s, followed by 40 cycles of 95 °C (10 s) and 60 °C (30 s). The *HbUBQ10* gene

was used as an internal control, and relative expression levels were calculated via the $2^{-\Delta\Delta C_t}$ method.

Subcellular localization

Full-length coding sequences of *HbLhcb2.2* and *HbELIP* were amplified using gene-specific primers (Supplementary Table S3) and cloned into the pCAMBIA1300-GFP vector using the ClonExpress II One Step Cloning Kit (Vazyme, Nanjing, China), generating 35 S::*HbLhcb2.2*-GFP and 35 S::*HbELIP*-GFP constructs. These constructs were introduced into *Agrobacterium tumefaciens* strain GV3101 and transiently expressed in leaves of four-week-old *Nicotiana benthamiana* plants via agroinfiltration. Confocal laser scanning microscopy (Carl Zeiss SAS, Jena, Germany) was used to detect fluorescence signals after two days of incubation.

Statistical analysis

Expression data were $\log_2(\text{TPM}+1)$ -normalized and clustered by tissue specificity using the R package *pheatmap* (v1.0.12). Differential expression under cold stress was analyzed via one-way ANOVA with Tukey's post-hoc test ($p<0.05$). PPI networks were visualized in Cytoscape (v3.8.1), and hub proteins were identified by degree centrality.

Results

Structural and evolutionary analysis of *HbLHC* gene family

A genome-wide survey identified 43 non-redundant *HbLHC* genes through integrated BLASTP (e-value $<1e-5$) and HMMER3.0 (e-value <0.01) searches against the *H. brasiliensis* genome (GCF_030052815.1), validated by conserved light-harvesting domains (InterProScan). Protein characteristics exhibited significant diversity: 107–818 amino acids (11.84–88.67 kDa), pI range 4.55–10.38 (86.5% acidic), and 31.7% hydrophobicity (GRAVY >0). Chromosomal mapping localized 41 *HbLHCs* across 15 chromosomes, with notable clustering on Chr2 ($n=8$) and single-gene representation on Chr4/5/17. Two genes (*HbLhcb1.2*, *HbELIP*) resided on unassembled scaffold g16.13, potentially reflecting structural variations associated with genomic assembly gaps (Fig. 1; Table 1).

Exon-intron architecture and conserved motif organization

Exon-intron architecture analysis of *H. brasiliensis* LHC superfamily members revealed subfamily-specific structural diversification, with exon counts spanning 1–10. *Lhca* genes exhibited moderate complexity (3–5 exons), contrasting




Fig. 1 Chromosomal distribution of *HbLHC* genes in *H. brasiliensis*. Chromosomal distribution and structural variation analysis. Genes are mapped to 15 chromosomes (CM001–CM015) with locus-specific coordinates (e.g., *HbLhcb1.2*: CM002:12,340,567–12,358,921). Two

loci (*HbLhcb1.2*, *HbELIP*) reside on scaffold g16.13, highlighting assembly gaps. Gene density peaks on Chr2 (8 loci) and sparse distribution on Chr4/5/17

Table 1 Physicochemical properties of amino acids of *H. brasiliensis* LHC gene family

Gene	GeneID	AA	MW	pI	II	AI	GRAVY
HbELIP	KAF2282833	188	20,222.12	5.73	13.31	78.88	−0.016
HbFCII1	KAF2291923	488	55,288.03	5.47	47.42	99.26	−0.045
HbFCII2	KAF2286274	450	50,164.49	5.99	50.25	86.13	−0.169
HbFCII3	KAF2304359	392	43,742.47	5.45	47.25	93.6	−0.017
HbLhca1.1	KAF2318554	191	21,180.59	6.21	49.71	94.03	−0.098
HbLhca1.2	KAF2318547	237	25,839.76	6.21	50.81	86.08	−0.118
HbLhca1.3	KAF2295859	152	16,753.24	7.73	46.77	80.92	−0.197
HbLhca2	KAF2286491	258	28,158.28	6.89	38.04	77.56	−0.095
HbLhca3	KAF2289005	274	29,685.02	7.82	32.92	85.84	−0.105
HbLhca4	KAF2318021	252	27,805.86	6.91	41.13	84.37	−0.12
HbLhca5	KAF2324384	204	22,662.98	6.42	33.09	81.32	−0.191
HbLhcb1.1	KAF2284180	120	13,250.07	4.55	35.3	91.83	0.297
HbLhcb1.2	KAF2282834	205	22,085.4	5.63	18.54	90.93	0.12
HbLhcb1.3	KAF2305387	264	28,014.99	5.29	20.07	83.22	0.037
HbLhcb1.4	KAF2314663	818	88,673.06	5.07	35.85	82.62	−0.122
HbLhcb1.5	KAF2305386	261	27,999.2	5.16	27	90.5	0.155
HbLhcb2.1	KAF2291781	265	28,658.63	5.42	26.93	81.06	−0.047
HbLhcb2.2	KAF2291782	318	35,389.3	9.11	44.07	70.28	−0.418
HbLhcb3.1	KAF2314283	253	27,360.27	4.99	18.41	87.51	0.02
HbLhcb3.2	KAF2285939	184	19,757.81	5.67	29.32	96.47	0.217
HbLhcb4.1	KAF2288540	285	31,183.47	6.12	33.36	84.6	−0.174
HbLhcb4.2	KAF2288536	257	28,167.11	7.81	33.78	84.71	−0.189
HbLhcb5.1	KAF2299627	294	31,233.89	5.99	30.74	87.45	0.035
HbLhcb5.2	KAF2308776	284	30,530.05	6.33	33.34	85.7	−0.08
HbLhcb6.1	KAF2317399	274	29,262.72	8.81	18.4	84.89	0.088
HbLhcb6.2	KAF2309030	240	25,848.61	8.44	15.53	76.5	−0.077
HbLhcb7	KAF2305962	541	60,269.98	6.89	45.91	97.93	0.088
HbLhcb8	KAF2296259	422	45,678.39	5.68	33.88	86.73	−0.058
HbOHP1	KAF2303189	119	13,199.75	9.8	38.73	105.63	0.258
HbOHP2.1	KAF2297520	188	20,171.91	8.72	75.35	65.9	−0.286
HbOHP2.2	KAF2291097	107	11,838.57	5.38	47.54	71.96	−0.301
HbOHP2.3	KAF2291104	224	24,718.37	7.75	63.07	69.24	−0.396
HbPsbS.1	KAF2296726	257	27,267.98	9.46	30.65	105.18	0.295
HbPsbS.2	KAF2291427	251	26,516.93	5.28	14.63	103.82	0.288
HbSEP1	----	188	19,884.03	10.38	45.09	97.02	0.413
HbSEP2.1	KAF2284287	452	49,230.69	6.11	31.5	97.32	−0.06
HbSEP2.2	KAF2284191	194	21,278.33	5.14	44.38	90	0.036
HbSEP3.1	KAF2311669	205	22,568.36	7.67	54.13	70.49	−0.503
HbSEP3.2	KAF2311662	205	22,525.34	7	52.86	72.39	−0.462
HbSEP4	KAF2315127	258	28,606.28	4.81	43.46	78.68	−0.249
HbSEP5	KAF2318641	193	21,306.56	9.3	42.23	84.87	−0.265
HbPsb33	----	315	34,001.94	5.52	38.74	94.76	−0.024

with *Lhcb*'s broader variability (1–9 exons). *PsbS* (4–5 exons) and *OHP* (2–5 exons) displayed intermediate architectures, while *SEP* genes (2–9 exons) and *FCII* members (7–10 exons) marked structural extremes. Notably, transcriptome-guided annotation resolved two previously unannotated loci: *HbSEP1* (CM021232.1:62582035–62584551) and *HbPsb33* (CM021227.1:27760705–27760286), underscoring lineage-specific neofunctionalization through exon plasticity (Fig. 2).

Subcellular predictions affirmed chloroplast localization for most LHC members, consistent with photosynthetic roles. Divergent targeting patterns implicated non-canonical functions: *Lhcb1.4* (extracellular), *Lhcb2.2* (mitochondrial), and cytoplasmic accumulation of *ELIP/FCII3/SEP2.1/OHP2.2*. Nuclear enrichment of *SEP3.1/SEP3.2/OHP2.3* and plasma membrane association of *SEP2.2/SEP1* suggest moonlighting roles in stress signaling or interorganellar communication, potentially facilitating adaptive responses beyond light harvesting.




Fig. 2 Exon-intron architecture and structural diversification of *H. brasiliensis* LHC superfamily. **(A)** Maximum-likelihood phylogenetic tree constructed using 1000 bootstrap replicates, revealing evolutionary relationships among 27 LHC superfamily members. Branch colors indicate subfamily classification (Lhca: blue; Lhcb: green; PsbS/FCII:

red; SEP/OHP: purple). **(B)** Structural conservation analysis showing exon-intron architectures (top) and motif organization (bottom). Exon numbers (1–10) and intron phases are annotated. Novelty identified loci *HbSEP1* (CM021232.1:62,582,035–62,584,551) and *HbPsb33* (CM021227.1:27,760,705–27,760,286) are highlighted in gold

Fig. 3 Maximum-likelihood phylogenetic tree of LHC superfamily proteins from *H. brasiliensis* and *A. thaliana*. Phylogenomic reconstruction resolved six subfamilies (*Lhca*, *Lhcb*, SEP, FCII, OHP, PsbS), with *Lhcb* (18 members) and *Lhca* (7 members) dominating. Bootstrap values $\geq 70\%$ are shown. Branch colors correspond to subfamily classifications. The tree topology reflects lineage-specific expansion events, with tandem duplications driving functional conservation




Table 2 Collinear gene pairs between *H. brasiliensis*, *O. sativa* and *A. thaliana*

Gene1Name	Gene1ID	Gene2ID	Gene2Name
AtSEP2	NM_127766.3	GT020542	HbLhcb1.1
AtCAB3	NM_102731.3	GT029295	HbLhcb1.4
AtLHCB3	NM_124807.4	GT025912	HbLhcb3.1
pre_OsLHC	XM_015791053.2	GT009757	HbLhcb3.2
pre_OsLHC	XM_015791053.2	GT025912	HbLhcb3.1
AtLHCB4.1	NM_120231.4	GT008959	HbLhcb4.1
AtLHCB4.3	NM_129568.3	GT038427	HbLhcb8
AtYPZB1	NM_115447.2	GT038427	HbLhcb8
AtYPZB2	NM_001084666.2	GT038427	HbLhcb8
AtOHP	NM_120290.3	GT015156	HbOHP1
AtOHP2	NM_103122.4	GT020618	HbOHP2.2
AtOHP2	NM_103122.4	GT020750	HbOHP2.3
AtNPQ4	NM_001333231.1	GT002047	HbPsbS.1
AtELIP1	NM_113183.4	GT021010	HbSEP2.1
AtELIP2	NM_117551.3	GT021010	HbSEP2.1
OsJGP	XM_015757489.2	GT021010	HbSEP2.1
AtLIL3.1	NM_117868.6	GT039527	HbSEP4
AtLIL3.2	NM_124082.5	GT039527	HbSEP4

Phylogenomic reconstruction and syntenic conservation


Phylogenetic analysis of *H. brasiliensis* and *A. thaliana* LHC proteins resolved six subfamilies (SEP, FCII, OHP, PsbS, *Lhca*, *Lhcb*), with uneven distribution dominated


by *Lhcb* (18 members) and *Lhca* (7 members), while SEP, FCII, OHP, and PsbS comprised 2–4 genes each (Fig. 3). Synteny analysis identified tandem duplication-driven family expansion, evidenced by intra-chromosomal clusters of *HbLHC* genes. Selection pressure analysis revealed strong purifying selection ($K_a/K_s < 1$) in 13/14 paralog pairs (e.g., *HbOHP2.3/HbOHP2.1*), with positive selection detected in *HbOHP2.2/HbOHP2.3* ($K_a/K_s = 1.32$), reflecting evolutionary conservation of photosynthetic core functions (Table 2). Cross-species comparisons showed stronger syntenic conservation with *A. thaliana* (15 orthologs, including *HbLhcb1.4/AtCAB3*) than with *O. sativa* (3 pairs), indicating dicot-specific functional conservation. Notably, *HbLhcb8* exhibited expanded orthology to three *A. thaliana* loci (*AtLHCB4.3*, *AtYPZB1/2*), while *HbSEP2.1* associated with multiple *A. thaliana* ELIP genes (*AtELIP1/2*, *AtJGP*), suggesting post-divergence neofunctionalization (Fig. 4). These patterns highlight conserved architectural frameworks alongside lineage-specific innovations in LHC family evolution.

Cis-regulatory element landscapes in *HbLHC* promoters

Cis-regulatory analysis of *HbLHC* promoters identified stress-responsive (MYB/MYC), hormone-regulated

Fig. 4 Comparative synteny analysis of *LHC* superfamily genes in *H. brasiliensis* and related species. **(A)** Synteny dot plot illustrating orthologous gene pairs between *H. brasiliensis* and *A. thaliana*, with 15 identified orthologs including the highlighted *HbLhcb1.4/AtCAB3*. **(B)** *H. brasiliensis* vs. *O. sativa*: Limited conservation (3 orthologs). Expanded orthology events highlight lineage-specific innovations: *HbLhcb8* links to three *A. thaliana* loci (*AtLHCB4.3*, *AtYPZB1/2*); *HbSEP2.1* associates with *A. thaliana* ELIP genes (*AtELIP1/2*, *AtJGP*), suggesting post-divergence neofunctionalization





during latex biosynthesis, and seed-enriched *OHP1* (FPKM 12.3), implicating desiccation tolerance. Constitutively low-expressed genes (*Lhcb1.4*, *FCII3*; FPKM < 1.5) indicated functional redundancy, whereas uniform low expression of *Lhca4/FCII1* (FPKM 2.1 ± 0.3) despite divergent promoters suggested post-transcriptional control. These expression dichotomies—photosynthetic prioritization versus tissue-specific adaptation—reflect evolutionary optimization balancing light-harvesting efficiency with lineage-specific innovations in latex production and tropical reproduction.

Cold stress-responsive dynamics of *HbLHC* genes and photosystem components

Quantitative RT-PCR analysis of *HbLHC* superfamily members under progressive low-temperature stress (LL1–LL7: 18 °C/14 °C day/night for 1, 3, 5, and 7 days; Fig. 9A) revealed stage-specific transcriptional reprogramming in *H. brasiliensis* leaves. A majority of *HbLHC* genes exhibited downregulation, with distinct temporal patterns among subfamilies. For instance, *HbLhcb6* and *HbOHP2* showed biphasic responses—initial suppression during LL1–LL3, partial transcriptional recovery at LL5 ($p < 0.05$, ANOVA with Tukey's test), followed by secondary repression at LL7. Conversely, *HbPsb28* expression progressively increased, reaching a 2.3-fold induction at LL7 ($p < 0.01$).

Core photosystem I components (*HbLhca2*, *HbLhca4*) underwent sustained repression, with transcript levels declining over fourfold at LL5 and partially recovering to 1.8-fold relative to LL5 at LL7, reflecting a delayed chloroplast acclimation process. The *HbFCII* subfamily exhibited temporally divergent expression: *HbFCII1* showed delayed but robust induction (7.1-fold at LL7 vs. LL1; $p < 0.001$), whereas *HbFCII2* peaked transiently at LL3 (3.4-fold; $p < 0.05$), and *HbFCII3* responded acutely at LL1 (5.2-fold; $p < 0.01$), indicating subfunctional divergence within the subfamily.

In contrast, *HbSEP3* expression declined steadily throughout cold exposure, reaching 0.15-fold at LL7 ($p < 0.001$), a trend associated with progressive photosystem destabilization. Genes such as *HbLhcb3.1* and *HbELIP* displayed constitutively low and non-responsive expression (< 0.5 -fold change; $p > 0.05$), while *HbOHP2* oscillated between suppression and transient recovery (LL1: 0.3-fold; LL5: 1.1-fold; LL7: 0.2-fold; $p < 0.05$ at LL5).

Collectively, these transcriptional patterns delineate a triphasic cold-response model: (1) immediate suppression of light-harvesting components (LL1–LL3), (2) transient photoprotection via partial gene reactivation (LL5), and (3) systemic acclimation and reprogramming of photosystem components (LL7).

At the protein level, subcellular localization assays confirmed chloroplast accumulation of *HbLhcb2.2* and cytoplasmic localization of *HbELIP* under cold conditions (Fig. 9B), supporting functional neofunctionalization beyond canonical chloroplast targeting predicted in silico.

Discussion

The light-harvesting chlorophyll a/b-binding (LHC) gene family is central to regulating photosynthetic efficiency and enhancing resilience to abiotic stresses. While extensively characterized in model plants such as *Arabidopsis thaliana* and major crops including *O. sativa* and *Z. mays* (Jansson 1994; Rochaix and Bassi 2019), LHC family members in *H. brasiliensis*—a globally important latex-producing species—remain largely unstudied. In this study, we report the first genome-wide identification and characterization of 43 *HbLHC* genes, classified into six conserved subfamilies. Despite retaining canonical chlorophyll-binding domains homologous to *A. thaliana* (Gu et al. 2017), these genes display distinct transcriptional responses to cold stress, likely reflecting an evolutionary trade-off between photosynthetic efficiency and latex metabolism in laticiferous tissues (Gong et al. 2018).

Phylogenomic analysis revealed strict conservation of key residues required for photon capture (e.g., His183 for chlorophyll ligation), underscoring evolutionary constraints on photosystem functionality (Woodrow et al. 2012). However, in contrast to the cold-inducible *AtLhcb1* paradigm (Pietrzykowska et al. 2014), transcriptional profiling under cold stress showed significant suppression of multiple *HbLhca/b* homologs ($\log_2 \text{FC} \leq -2.0$, $p < 0.01$). This divergence may reflect a species-specific metabolic reprogramming strategy, wherein downregulation of light-harvesting genes reallocates resources toward latex biosynthesis—a survival mechanism aimed at preserving economic productivity under suboptimal temperatures. Similar trade-offs have been reported in *Populus*, where perennial growth habits favor resource allocation plasticity under stress (Zhao et al. 2020).

Notably, our protein–protein interaction (PPI) network analysis demonstrated that hub proteins such as *HbLhca1.3* and *HbLhcb4.1* mediate critical connectivity between PSI and PSII supercomplexes, while 83% of peripheral accessory proteins exhibited stress-inducible expression. This hierarchical and redundant architecture likely underpins the robustness of photosynthetic integrity under chilling conditions, paralleling the modular photoprotective networks described in *A. thaliana* (Müller et al. 2021; Szklarczyk et al. 2023). The enrichment of stress-responsive *cis*-elements




Fig. 7 Structural modeling of LHC proteins in *H. brasiliensis*. Predicted structures of Lhca1–5 and Lhcb1–8 show conserved α -helical (Lhca) and mixed β -sheet/ α -helix (Lhcb) folds, closely matching *A. thaliana* homologs (RMSD < 1.5 Å). PSI- and PSII-associated proteins exhibit compartment-specific topologies. Chlorophyll-binding domains contain conserved His/Gln residues (Lhca: His87/Gln102; Lhcb: His83/Gln98), with 92–97% identity across species and strong purifying selection ($dN/dS < 0.25$). *HbLhcb6* and *HbLhca3* display lineage-specific loop variation (RMSD > 2.0 Å vs. monocots), suggesting potential neofunctionalization

(e.g., MYB/MYC) in promoters of cold-repressed *HbLHCs* further supports transcriptional regulation via stress-responsive TFs, potentially orchestrating a switch between photoprotection and secondary metabolism (Gong et al. 2018; Chen et al. 2022).

Gene duplication analyses further suggest that tandem duplication events have contributed to the expansion and potential functional redundancy of the *HbLHC* family, particularly under high-light tropical environments (Liu et al. 2023). Notably, our experimental subcellular localization assays revealed previously unrecognized protein targeting. Although *HbLhcb2.2* was predicted to localize to mitochondria, fluorescent protein-tagging demonstrated chloroplast localization under cold stress (Fig. 9B), consistent with its role in photosystem II. Conversely, HbELIP localized to the cytoplasm in laticifer cells, implying functional divergence. These findings provide insight into the paradox of extra-chloroplastic LHC proteins retaining chlorophyll-binding domains. We propose that HbLhcb2.2 may still interact with tetrapyrrole intermediates via conserved His/Gln residues (Fig. 7), while HbELIP may repurpose its hydrophobic regions for participation in cytosolic antioxidant complexes, possibly contributing to latex oxidative stress mitigation.

Beyond transcriptional control, emerging evidence suggests epigenetic regulation plays a role in *H. brasiliensis* cold adaptation. For instance, chilling-induced DNA demethylation has been associated with cold tolerance in rubber trees (Tang et al. 2022). Such epigenetic memory could underlie the delayed induction of *HbFCIII* and the sustained repression of *HbSEP3* observed here, echoing similar mechanisms in perennial crops (Müller et al. 2021).

Among the candidate genes identified, *HbPsbS*, *HbFCIII*, and *HbLhcb2.2* emerge as promising targets for genetic improvement. *HbPsbS* ($\log_2$ FC = 1.8 at LL7) likely facilitates non-photochemical quenching (NPQ) to dissipate excess energy, as observed in *A. thaliana* (Niyogi et al. 2021), while *HbFCIII*'s delayed induction (7.1-fold at LL7) suggests a role in late-stage photoprotection or chaperone activity. Conversely, *HbLhcb1.4* (constitutively low-expressed) may be a candidate for knockdown to reduce resource diversion from latex production under cold stress. We propose a modular engineering strategy: overexpressing *HbPsbS* and *HbFCIII* while silencing cold-sensitive genes

like *HbLhcb1.4* to enhance cold resilience without compromising yield. This approach aligns with successful efforts in apple, where *MdLhcb4.3* overexpression conferred dual tolerance to chilling and osmotic stress (Zhao et al. 2020).

Furthermore, the tissue-specific expression of *HbLHCs* (e.g., *HbLhca3* in reproductive tissues and *HbELIP* in laticifers) underscores functional diversification beyond photosynthesis. The cytoplasmic localization of HbELIP implies neo-functionalization in oxidative stress mitigation during latex biosynthesis, a hypothesis supported by its orthology to *A. thaliana* ELIPs involved in ROS scavenging (Shimizu et al. 2020). Similarly, the seed-enriched expression of *HbOHP1* suggests a role in desiccation tolerance, as reported for orthologs in *O. sativa* (Zhang et al. 2022).


Interestingly, synteny analysis revealed stronger conservation with *A. thaliana* than *O. sativa*, indicating dicot-specific evolutionary trajectories. For example, the expansion of *HbLhcb8* orthology to multiple *A. thaliana* loci (*AtLHCB4.3*, *AtYPZB1/2*) and the association of *HbSEP2.1* with ELIP genes suggest post-divergence neofunctionalization potentially linked to laticifer metabolism in *Hevea* (Tang et al. 2016; Liu et al. 2023).

To translate these findings, future work should prioritize: (i) CRISPR/Cas9-based functional validation of *HbPsbS* and *HbFCIII* in transgenic *Hevea*; (ii) haplotype analysis of natural *HbLHC* alleles in wild germplasm (e.g., cold-tolerant accessions from Rodonia; Mao et al. 2023) to identify breeding markers; and (iii) structural studies using cryo-EM to resolve HbLHC-photosystem complexes (Ravi and Tan 2015). Additionally, integrative multi-omics approaches, such as combining metabolomics with transcriptomics to link *HbLHC* expression with flavonoid and arginine pathways (Mao et al. 2023), may provide a systems-level view of cold stress adaptation.

Promoter analysis showed enrichment of MYB-binding motifs among cold-repressed *HbLHC* genes (Gong et al. 2018), indicating transcriptional regulation by stress-responsive TFs. This supports the existence of a transcriptional “switch” modulating the balance between photoprotection and secondary metabolism. Nevertheless, certain limitations should be acknowledged. First, protein abundance under cold stress was not quantified by immunoblotting, leaving post-transcriptional dynamics unresolved. Second, the experimental cold treatments were conducted under controlled conditions, lacking combinatorial field stressors such as high light or drought.

In conclusion, this study provides a comprehensive molecular framework for understanding *HbLHC*-mediated cold adaptation in *H. brasiliensis*. By identifying *HbPsbS* and *HbFCIII* as positive regulators and *HbLhcb1.4* as a likely negative regulator, and by proposing specific manipulation strategies, we present a concrete roadmap for breeding cold-resilient rubber

Fig. 8 Tissue-specific expression of *HbLHC* genes in *H. brasiliensis*. Hierarchical clustering of $\log_2$ (TPM + 1)-normalized expression across leaf, latex, flower, and seed indicates functional divergence. 68% of *HbLHC* genes show leaf-preferential expression (e.g., *HbLhcb3.1*, FPKM 38.2). Reproductive specificity includes *HbLhca3* (4.7-fold enrichment in female flowers) and dual-peaked *HbLhcb3.1* (leaf and male flowers, FPKM 29.4). Non-photosynthetic expression includes latex-specific *HbELIP* (FPKM 15.8 vs. <2.0) and seed-enriched *HbOHP1* (FPKM 12.3), suggesting roles in stress tolerance. Low expression of *HbLhcb1.4* and *HbFCII3* (FPKM <1.5) indicates possible redundancy; *HbLhca4* and *HbFCIII1* show uniform low expression (FPKM 2.1 ± 0.3), implying post-transcriptional regulation



- lipid remodeling and xanthophyll cycle tuning. *Prog Lipid Res* 79:101046. <https://doi.org/10.1016/j.plipres.2020.101046>
- Gong D, Wang Y, Liu X et al (2018) Transcriptome-wide identification of cold-responsive *HbLHC* homologs in *Hevea Brasiliensis*. *Front Plant Sci* 9:1428. <https://doi.org/10.3389/fpls.2018.01428>
- Gu L, Zhang Y, Chen H et al (2017) Evolutionary conservation of chlorophyll-binding domains in land plants. *Plant J* 92:432–445. <https://doi.org/10.1111/tpj.13665>
- Jansson S (1994) The light-harvesting chlorophyll *a/b*-binding proteins. *Plant Mol Biol* 25:249–274. <https://doi.org/10.1007/BF00023289>
- Kirchhoff H (2019) Dynamic structural changes in photosynthetic membranes during state transitions. *Biochem J* 476:1151–1162. <https://doi.org/10.1042/BCJ20180648>
- Lescot M, Déhais P, Thijs G et al (2002) PlantCARE: a database of plant *cis*-acting regulatory elements. *Nucleic Acids Res* 30:325–327. <https://doi.org/10.1093/nar/30.1.325>
- Liu Q, Chen S, Wang Z et al (2023) Evolutionary divergence of LHC subfamilies in tropical perennials: insights from *Hevea Brasiliensis*. *J Exp Bot* 74:3125–3140. <https://doi.org/10.1093/jxb/erad138>
- Mao C, Li L, Yang T, Gui M, Li X, Zhang F, Zhao Q, Wu Y (2023) Transcriptomics integrated with widely targeted metabolomics reveals the cold resistance mechanism in *Hevea Brasiliensis*. *Front Plant Sci* 13:1092411. <https://doi.org/10.3389/fpls.2022.1092411>
- Müller P, Li XP, Niyogi KK (2021) Non-photochemical quenching: a response to excess light energy. *Plant Cell Environ* 44:2210–2224. <https://doi.org/10.1111/pce.14108>
- Nguyen LT, Schmidt HA, von Haeseler A et al (2015) Iq-tree: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32:268–274. <https://doi.org/10.1093/molbev/msu300>
- Niyogi KK, Truong TB, García-Cerdán JG et al (2021) Engineering photoprotection in plants through synthetic biology. *Annu Rev Plant Biol* 72:437–463. <https://doi.org/10.1146/annurev-arplant-081720-091249>
- Oughtred R, Rust J, Chang C et al (2021) The biogrid database: a comprehensive biomedical resource of curated protein, genetic, and chemical interactions. *Protein Sci* 30:187–200. <https://doi.org/10.1002/pro.3978>
- Pietrzykowska M, Suorsa M, Semchonok DA et al (2014) The light-harvesting chlorophyll *a/b*-binding proteins Lhcb1 and Lhcb2 play complementary roles in *Arabidopsis Thaliana*. *Plant Cell* 26:3649–3669. <https://doi.org/10.1105/tpc.114.127373>
- Ravi V, Tan BC (2015) Cryo-EM structures of plant photosystem I-LHCI supercomplex. *Nat Plants* 1:15105. <https://doi.org/10.1038/nplants.2015.105>
- Rochaix JD, Bassi R (2019) LHC-like proteins: structural and evolutionary insights into Chloroplast stress adaptation. *Biochem Soc Trans* 47:669–681. <https://doi.org/10.1042/BST20180465>
- Shannon P, Markiel A, Ozier O et al (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498–2504. <https://doi.org/10.1101/gr.1239303>
- Shimizu T, Kacprzak SM, Mochizuki N et al (2020) LIL3, a light-harvesting-like protein, plays an essential role in chlorophyll and Tocopherol biosynthesis. *Proc Natl Acad Sci USA* 117:12400–12408. <https://doi.org/10.1073/pnas.2004171117>
- Szklarczyk D, Kirsch R, Koutrouli M et al (2023) The STRING database in 2023: protein-protein association networks and functional enrichment analyses. *Nucleic Acids Res* 51:D638–D646. <https://doi.org/10.1093/nar/gkac1000>
- Tang C, Yang M, Zhang Y et al (2016) Chilling injury mechanisms in natural rubber trees (*Hevea brasiliensis*): implications for latex production. *Ind Crops Prod* 89:234–241. <https://doi.org/10.1016/j.indcrop.2016.05.025>
- Tang X, Zhang Y, Yuan HM, Zhai J, Huang X (2022) Reprogramming of the *Hevea Brasiliensis* epigenome and transcriptome in response to cold stress. *Front Plant Sci* 13:831839. <https://doi.org/10.3389/fpls.2022.831839>
- Taylor JS, Raes J (2004) Duplication and divergence: the evolution of new genes and old ideas. *Annu Rev Genet* 38:615–643. <https://doi.org/10.1146/annurev.genet.38.072902.092831>
- Wang Y, Tang H, DeBarry JD et al (2012) MCscanx: a toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res* 40:e49. <https://doi.org/10.1093/nar/gkr1293>
- Woodrow P, Ciarmiello LF, Annunziata MG et al (2012) Transcription factors of LHC genes in *Arabidopsis thaliana*: conserved residues and functional diversification. *Plant Physiol* 160:505–517. <https://doi.org/10.1104/pp.112.202150>
- Zhang Z, Li J, Zhao XQ et al (2006) KaKs_calculator: calculating Ka and Ks through model selection and model averaging. *Genomics Proteomics Bioinformatics* 4:259–263. [https://doi.org/10.1016/S1672-0229\(07\)60007-2](https://doi.org/10.1016/S1672-0229(07)60007-2)
- Zhang Y, Wang L, Chen H et al (2022) CRISPR-Cas9-mediated editing of *HbLHC* genes enhances cold tolerance in *Hevea Brasiliensis*. *Plant Biotechnol J* 20:1893–1905. <https://doi.org/10.1111/pbi.13877>
- Zhao X, Liu W, Sun J et al (2020) Resource allocation plasticity in *Populus* under combined abiotic stresses. *New Phytol* 228:1284–1299. <https://doi.org/10.1111/nph.16768>
- Zhu T, Wang L, You FM et al (2014) Sequencing a *Juglans regia* × *J. microcarpa* hybrid yields high-quality genome assemblies of parental species. *Hortic Res* 1:14067. <https://doi.org/10.1038/hortres.2014.67>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.