

Global no net loss of natural ecosystems

Maron, Martine; Simmonds, Jeremy S.; Watson, James E. M.; Sonter, Laura J.; Bennun, Leon; Griffiths, Victoria F.; Quétier, Fabien; von Hase, Amrei; Edwards, Stephen; Rainey, Hugo; Bull, Joseph W.; Savy, Conrad E.; Victurine, Ray; Kiesecker, Joseph; Puydarrieux, Philippe; Stevens, Todd; Cozannet, Naïg; Jones, J.P.G.

Nature Ecology and Evolution

DOI:

10.1038/s41559-019-1067-z

Published: 01/01/2020

Peer reviewed version

Cyswllt i'r cyhoeddiad / Link to publication

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Maron, M., Simmonds, J. S., Watson, J. E. M., Sonter, L. J., Bennun, L., Griffiths, V. F., Quétier, F., von Hase, A., Edwards, S., Rainey, H., Bull, J. W., Savy, C. E., Victurine, R., Kiesecker, J., Puydarrieux, P., Stevens, T., Cozannet, N., & Jones, J. P. G. (2020). Global no net loss of natural ecosystems. Nature Ecology and Evolution, 4(1), 46-49. Article 33. https://doi.org/10.1038/s41559-019-1067-z

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 - You may not further distribute the material or use it for any profit-making activity or commercial gain
 - You may freely distribute the URL identifying the publication in the public portal?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Global no net loss of natural ecosystems

- 2 Martine Maron^{1,2*}, Jeremy S. Simmonds^{1,2}, James E.M. Watson^{1,2,3}, Laura J. Sonter^{1,2}, Leon
- 3 Bennun^{4,5}, Victoria F. Griffiths⁶, Fabien Quétier⁷, Amrei von Hase⁸, Stephen Edwards⁹, Hugo
- 4 Rainey³, Joseph W. Bull¹⁰, Conrad E. Savy¹¹, Ray Victurine³, Joseph Kiesecker¹², Philippe
- 5 Puydarrieux⁹, Todd Stevens³, Naïg Cozannet¹³, Julia P.G. Jones^{14s}
- 6 1. Centre for Biodiversity and Conservation Science, The University of Oueensland, St Lucia 4072,
- 7 Australia.
- 8 2. School of Earth and Environmental Sciences, The University of Queensland, St Lucia 4072,
- 9 Australia.
- 3. Wildlife Conservation Society, Global Conservation Program, New York 10460, United States of
- 11 America
- 4. The Biodiversity Consultancy, Cambridge CB2 1SJ, United Kingdom.
- 13 5. Conservation Science Group, Department of Zoology, University of Cambridge, Cambridge CB2
- 14 3EJ, United Kingdom.
- 15 6. Department of Zoology, University of Oxford, Oxford OX1 3SZ, United Kingdom.
- 7. Biotope, 34140 Mèze, France.
- 17 8. Forest Trends, Washington, DC 20036, United States of America and Bo-Kaap 8001, South Africa.
- 18 9. International Union for Conservation of Nature (IUCN), CH-1196 Gland, Switzerland.
- 19 10. Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation,
- 20 University of Kent, United Kingdom.
- 21 11. International Finance Corporation, Nairobi, Kenya.
- 22 12. The Nature Conservancy, Global Conservation Lands Program, Colorado 80524, United States of
- 23 America.
- 24 13. Agence Française de Développement, Paris, France.
- 25 14. College of Environmental Sciences and Engineering, Bangor University, Bangor LL57 2UW,
- 26 United Kingdom.
- 27 Corresponding author: Martine Maron (m.maron@ug.edu.au)

Abstract

29

- 30 A global goal of No Net Loss (GNNL) of natural ecosystems or better has recently been
- proposed, but such a goal would require equitable translation to country-level contributions.
- 32 Given the wide variation in ecosystem depletion, these could vary from Net Gain (for
- countries where restoration is needed), to Managed Net Loss (in rare circumstances where
- 34 natural ecosystems remain extensive and human development imperative is greatest).
- National contributions and international support for implementation also must consider non-
- area targets (e.g. for threatened species) and socioeconomic factors such as the capacity to
- 37 conserve and the imperative for human development.

38

39

Main text

- 40 Momentum is building for an ambitious new commitment to be signed at the Conference of
- 41 the Parties to the Convention on Biological Diversity (CBD) in 2020 as a global framework
- for nature conservation¹⁻⁴. Notable are calls for retention of half the Earth's natural
- ecosystems^{5,6}, to be enshrined by 2030 as a target under the deal. Yet this leaves little 'room
- 44 to move'—approximately half the Earth's terrestrial ecosystems have already been lost⁷.
- Nevertheless, complete cessation of anthropogenic impacts on natural ecosystems is
- infeasible, given the imperative for socioeconomic development where current levels of
- 47 human development are low⁸. Conservation that ignores such differences among nations is
- 48 likely to be unjust⁹.
- 49 In this context, a goal of global No Net Loss (GNNL) of natural ecosystems is likely the most
- ambitious target that society can realistically achieve^{10,11}, at least by 2030. Such a goal allows
- for losses in some places and gains in others, which, taken together, ensure no further net
- decline of natural ecosystems, benefitting the species and people which rely upon them¹².
- 53 GNNL implies an absolute cessation of decline in net terms—a key distinction from the
- relative 'NNL' that characterises biodiversity offset policies¹³.
- It is far from trivial to translate a GNNL goal to effective policy mechanisms and mitigation
- approaches at the national level; indeed, the problem is akin to dividing humanity's 'carbon
- 57 budget' equitably ^{14,15}. Here, we examine how different countries might set goals for retention
- 58 and restoration as part of a contribution to achieving GNNL of natural ecosystems, using
- 59 terrestrial ecosystems as an example.

- Translating a GNNL goal to a blanket requirement for each nation to achieve NNL would
- clearly be inappropriately coarse. Instead, a GNNL target would act as an umbrella for a
- range of minimum net outcome goals adopted by each country as their respective
- contributions to GNNL (Fig. 1). Some countries support natural ecosystems across almost
- their entire extent—10 have more than 75% of original natural ecosystems according to the
- latest published human footprint¹⁶ (e.g. Suriname and Canada Fig. 1; see Methods for more
- detail), while others retain close to none of their original natural ecosystems in reasonable
- condition (68 countries including France, Italy and India have <5% remaining; Fig. 1).
- 68 Countries also vary tremendously in the imperative to convert or degrade those ecosystems in
- 69 the pursuit of needed economic development, and in their capacity to protect and restore
- 70 ecosystems. So, under a GNNL commitment, some countries might focus on restoring earlier
- 71 losses, while others might further deplete their remaining ecosystems. Thus, some countries
- 72 might commit to Net Gain, some to NNL, and in some circumstances, controlled loss, or
- drawdown, of ecosystems (here termed Managed Net Loss).
- 74 Information about depletion of natural ecosystems can help frame both country-level
- conservation goals, and policy mechanisms for achieving those goals. For example, even
- NNL is likely to be inadequate to conserve threatened species and functioning ecosystems for
- countries whose natural ecosystems are most severely depleted. Therefore, for such countries,
- Net *Gain* in the extent of their natural ecosystems is likely to be essential. For example, the
- 79 UK has only 6% of ecosystems with a Human Footprint of <4 remaining (a threshold used as
- a proxy for ecosystem intactness⁷). The UK government recently proposed biodiversity Net
- 81 Gain as a requirement for new development projects¹⁷. Similarly, France has committed to
- 82 zero net conversion of natural land¹⁸. On the other hand, those countries with largely intact
- remaining ecosystems (e.g. Suriname, Gabon) may, in some circumstances, be able to accept
- further limited and controlled depletion ('Managed Net Loss') (Fig. 1). However, even if all
- 85 countries with less than 25% of natural ecosystems remaining adopt Net Gain and seek to
- double the extent of those ecosystems through restoration, this would only contribute 4% to
- 87 global ecosystem extent. Conversely, even a small percentage of net loss from countries with
- extensive remaining natural ecosystems, such as Australia and Brazil (5,535,401 km² and
- 89 4,643,615 km², respectively), would shift a very substantial restoration burden to other
- 90 countries, if GNNL is to be achieved.
- 91 Even within countries that retain similar amounts of natural ecosystems, variation in
- 92 depletion among different ecosystems can be lower (e.g. Norway, where retention of all its

```
93
       different ecosystem types is similarly high) or higher (e.g. Chad, where some ecosystems are
 94
       much more depleted than others). In such cases, approvals for unavoidable losses of less-
 95
       depleted ecosystem types might be tied to requirements to restore other, more-depleted
       ecosystems, using compensatory policy mechanisms like biodiversity offsetting 19,20. Further
 96
       complexity is introduced by the fact that some ecosystems may be extensive within a country,
 97
 98
       but globally rare; conversely, others are highly-depleted at a country level, yet globally
 99
       common. Therefore, both country-level goal-setting, and trading losses for gains among
100
       different ecosystems within a country, must reflect this variation to ensure all ecosystem
101
       types can be adequately conserved.
102
       We use the retention of terrestrial natural ecosystems to illustrate the complexity of
103
       translating GNNL to country-level goals, and propose that a similar exercise could consider
104
       the translation of the concept to the marine realm, or indeed to non-political units such as
105
       ecoregions. However, area-based retention is only one type of target that must be set for
106
       biodiversity to be adequately conserved. For example, the number of species listed as
       threatened with extinction does not correlate strongly with the depletion of natural
107
       ecosystems within a country (Pearson's R = 0.17; Figure 1a), though species decline often
108
       lags behind habitat loss<sup>21</sup>. Therefore, further ecosystem losses even from countries with
109
110
       relatively extensive natural systems could have a disproportionately negative impact in the
       most diverse but imperilled places (e.g., Brazil; 55% ecosystems remaining but 290 globally-
111
112
       threatened species of birds, mammals and amphibians).
113
       A purely biophysical basis for conservation goal-setting in a country ignores important
114
       socioeconomic realities, which may further modify appropriate relative contributions of
115
       countries to a GNNL goal. Countries vary enormously in their levels of human development;
       people's basic needs in many countries are not currently being met<sup>12</sup>. Rapid economic growth
116
117
       for those at the bottom of the global wealth rankings is the most important goal for
118
       governments in many such countries and is essential from a human rights perspective. The
119
       countries with the most severe ecosystem depletion (and therefore requiring, in principle,
120
       biodiversity Net Gain) include many countries with the lowest Human Development Index
       (HDI) values (e.g. numerous African countries) (Fig. 2). Given that converting ecosystems
121
122
       can contribute to much needed development, and significant amounts of ecosystem
123
       degradation in poorer countries has contributed to fuelling economic growth in richer
       countries<sup>22</sup>, it is unrealistic as well as unjust for goals to be set without socio-economic
124
125
       circumstances being considered. Addressing these equity implications, while also recognising
```

- the fundamental role of nature in supporting achievement of the Sustainable Development
- Goals¹², will also be essential to secure support for a GNNL commitment.
- Given that globally, biodiversity loss already exceeds safe levels²³, NNL at the country level
- might be the minimum acceptable standard for wealthy, developed countries where standards
- of living are already high (e.g. Australia, Canada; Fig. 2). We suggest their conservation
- goals should be set such that further degradation and loss of ecosystems is halted—at least in
- net terms. This may require radical solutions including moving away from the paradigm that
- economic growth is always desirable⁹.
- 134 Countries with low HDI are more likely to face further pressure on their natural ecosystems
- to facilitate urgently-needed economic development. Therefore, even where the level of
- depletion of natural ecosystems implies a NNL goal, Managed Net Loss may be unavoidable
- for such countries (Fig. 2), at least temporarily²⁴. Countries with a low HDI may reasonably
- expect support from the international community to deliver on their contribution to a GNNL
- goal. Unfortunately, weak governance in some low HDI countries discourages such
- investment²⁵ and can limit the effectiveness of any development support²⁶ or of any in-
- country mechanisms to compensate for biodiversity losses. For example, many of the
- countries to which assistance may need to be provided score poorly on the Corruption
- Perceptions Index (Fig. 2). Achievement of global biodiversity conservation arguably is most
- sensitive not to the global goals and targets that are agreed, but to how well such complex
- challenges to their implementation are addressed²⁷.
- Our framework provides guidance on the principles through which different countries could
- identify appropriate respective contributions toward a global goal of NNL of biodiversity.
- Any agreed set of contributions must tackle the reality of both biodiversity depletion, its
- causes, and global inequity in both ongoing pressures and capacity to respond to them. Goals
- must be transparently managed to avoid the task falling inequitably upon the world's poorest
- countries, while recognising that development at the expense of biodiversity is
- unsustainable²⁸.
- Loss without limit is the paradigm under which natural ecosystems are currently being
- destroyed³. The need to clarify the overarching goal of the CBD and sharpen our
- commitments to retain, restore, and protect natural ecosystems was underscored resoundingly
- by the recent release of the IPBES global assessement²⁹. So, as the focus turns to setting post-
- 2020 conservation targets under the CBD, calls to dramatically increase their ambition^{1,30} and

to set explicit nature retention targets³ must be heeded—and a pathway to translate them to country-level contributions laid out. A GNNL goal sets a limit to the loss we—and biodiversity—can tolerate, while allowing for human development where it is most urgently needed. Any basis for country-level commitments to a GNNL goal must reflect the substantial variation among countries in the level of depletion of their natural ecosystems—but also the degree to which capacity to conserve and the imperative for human development varies globally.

We used the depletion of natural ecosystems as one proxy for biodiversity loss, and the global

165

166

167

168

169170

171

172173

174

175

176

177

178

179

180

181

182

183

184

185 186

187

188

189

158159

160

161

162

163

164

Methods

Human Footprint 2009 dataset³¹ as an indicator of this depletion. The Human Footprint is a comprehensive representation of anthropogenic threats to biodiversity, which cumulatively accounts for eight human pressures—built environments, crop lands, pasture lands, human population density, night lights, railways, major roadways, and navigable waterways³¹. It is mapped across the terrestrial surface of the globe at a 1 km² resolution, on a scale of 0 (lowest Human Footprint) to 50 (highest Human Footprint). Human Footprint values of 0-3 are representative of land that is largely devoid of infrastructure and development (although may support sparse human populations)^{7,32}. We therefore considered areas with a Human Footprint value of ≥ 4 to be transformed – in other words, no longer supporting a natural ecosystem (as per Watson, et al.⁷). For 170 countries (for which data were also available for all measures), we calculated the area of the country that is mapped with Human Footprint values of 0-3, as a proportion of the area of the country (for which Human Footprint mapping was available). This represented our measure of the proportion of the original natural ecosystems remaining in each country. We also calculated the variance in depletion of specific natural ecosystem types in each country. To do this, we used the map of global terrestrial ecoregions³³, to represent the broad ecosystem types that do or would have naturally occurred in each country. We calculated the loss of each ecoregion type per country, by overlaying the Human Footprint map (value ≥ 4). To calculate the variation in depletion among ecoregion types within each country, we used the Gini coefficient – a metric frequently used to indicate dispersion within a frequency distribution. Although most commonly used as an index of income inequality, it can be used as an index of inequality for disparate datasets; a value of 0 indicates all values are identical

190	and 1 indicates extreme disparity among values. All GIS analysis was undertaken using
191	ArcMap6.1, with spatial datasets projected to a Mollweide coordinate system.
192	To explore the extent to which countries differ in their biophysical context, we plotted the
193	proportion of the original natural ecosystems remaining in each country against the variance
194	in depletion of natural ecosystems. We also considered two other measures of the status of a
195	country's biodiversity: the number of species listed as threatened under the IUCN Red List of
196	Threatened Species (restricted to fully assessed taxa only, as of November 2018: mammals,
197	birds, amphibians; note that most taxa are poorly known, so this too is a partial measure); and
198	the total area (km²) of natural ecosystems remaining in each country.
199	To examine how countries varied in environmental and socioeconomic contexts, we
200	incorporated two further datasets into our analysis. We used the 2017 Human Development
201	Index (HDI) ³⁴ as a representation of key elements of human development at the national
202	level. This composite metric subsumes indices relating to life expectancy, education and per
203	capita income. We also considered the 2017 Corruption Perceptions Index (CPI) ³⁵ , which
204	represents relative public sector corruption levels of nations as perceived by experts and
205	businesspeople, and has been linked with the strength of a nation's democratic institutions ³⁶ .
206	We plotted these variables as they relate to a nation's level of depletion of ecosystems, to
207	examine how variation in a country's socioeconomic factors potentially affect its capacity to
208	contribute to a goal of GNNL.
209	
210	Author contributions
211	MM, JPGJ, JEMW and JSS led the writing. JSS led the data analysis. All authors developed
212	the central concepts collaboratively, and wrote and edited parts of the manuscript.
213	
214	Data availability
215	All datasets used in this analysis are available via the citations identified in the Methods
216	section. The raw data used to create Figure 1 and Figure 2 are available in Supplementary
217	Table 1.
218	Code availability
219	No custom code was used.

1	1	n
Z	Z	U

221	Comp	peting interests statement		
222	LB, F	Q and AvH receive income from commercial contracts for consultancy services related		
223	to the	development and implementation of biodiversity offset policies.		
224				
225	Ackn	owledgements		
226	This v	This work was funded via the Science for Nature and People Partnership and its support of		
227	the Co	the Compensatory Conservation Working Group. MM was supported by Australian Research		
228	Counc	Council Future Fellowship FT140100516. The work was supported by the COMBO Project		
229	(funde	(funded by the Agence Française de Développement, Fonds Français pour l'Environnement		
230	Mond	ial and the MAVA Foundation)		
231				
232	References			
233	1	Dinerstein, E. et al. A global deal for nature: guiding principles, milestones, and		
234		targets. Science Advances 5, eaaw2869, doi:10.1126/sciadv.aaw2869 (2019).		
235	2	Mace, G. M. et al. Aiming higher to bend the curve of biodiversity loss. Nature		
236		Sustainability 1, 448-451, doi:10.1038/s41893-018-0130-0 (2018).		
237	3	Maron, M., Simmonds, J. S. & Watson, J. E. M. Bold nature retention targets are		
238		essential for the global environment agenda. Nature Ecology & Evolution 2, 1194-		
239		1195, doi:10.1038/s41559-018-0595-2 (2018).		
240	4	Pimm, S. L., Jenkins, C. N. & Li, B. V. How to protect half of Earth to ensure it		
241		protects sufficient biodiversity. Science Advances 4, eaat2616,		
242		doi:10.1126/sciadv.aat2616 (2018).		
243	5	Dinerstein, E. et al. An ecoregion-based approach to protecting half the terrestrial		
244		realm. Bioscience 67, 534-545, doi:10.1093/biosci/bix014 (2017).		
245	6	Wilson, E. O. Half-earth: Our planet's fight for life. New York: Liveright Publishing		

Corporation (2016).

246

- Watson, J. E. M. *et al.* Persistent disparities between recent rates of habitat conversion
- and protection and implications for future global conservation targets. *Conservation*
- 249 *Letters* **9**, 413-421, doi:10.1111/conl.12295 (2016).
- Nilsson, M., Griggs, D. & Visbeck, M. Policy: Map the interactions between
- Sustainable Development Goals. *Nature* **534**, 320-322, doi:10.1038/534320a (2016).
- Büscher, B. et al. Half-Earth or Whole Earth? Radical ideas for conservation, and
- 253 their implications. *Oryx* **51**, 407-410, doi:10.1017/S0030605316001228 (2017).
- 254 10 Arlidge, W. N. S. *et al.* A global mitigation hierarchy for nature conservation.
- 255 *Bioscience* **68**, 336-347, doi:10.1093/biosci/biy029 (2018).
- Bull, J. W. et al. Net positive outcomes for nature. Nature Ecology & Evolution (In
- press).
- United Nations. The Sustainable Development Goals Report 2018. New York, United
- States of America (2018).
- 260 13 Maron, M. et al. The many meanings of no net loss in environmental policy. Nature
- 261 Sustainability 1, 19-27, doi:10.1038/s41893-017-0007-7 (2018).
- 262 14 Geden, O. An actionable climate target. *Nature Geoscience* **9**, 340-342,
- doi:10.1038/ngeo2699 (2016).
- Holz, C., Kartha, S. & Athanasiou, T. Fairly sharing 1.5: national fair shares of a
- 265 1.5 °C-compliant global mitigation effort. *International Environmental Agreements*:
- 266 *Politics, Law and Economics* **18**, 117-134, doi:10.1007/s10784-017-9371-z (2018).
- Venter, O. et al. Global terrestrial Human Footprint maps for 1993 and 2009.
- Scientific Data 3, 160067, doi:10.1038/sdata.2016.67 (2016).
- 269 17 Government of United Kingdom (DEFRA). Government to mandate 'biodiversity net
- 270 gain', https://deframedia.blog.gov.uk/2019/03/13/government-to-mandate-
- biodiversity-net-gain/> (2019).
- 272 18 Government of France. *Plan biodiversité*, <a href="https://www.ecologique-
- solidaire.gouv.fr/plan-biodiversite> (2018).

- Bull, J. W., Hardy, M. J., Moilanen, A. & Gordon, A. Categories of flexibility in
- biodiversity offsetting, and their implications for conservation. *Biological*
- 276 *Conservation* **192**, 522-532, doi:https://doi.org/10.1016/j.biocon.2015.08.003 (2015).
- 277 20 Maron, M. et al. Taming a wicked problem: resolving controversies in biodiversity
- offsetting. *Bioscience* **66**, 489-498, doi:10.1093/biosci/biw038 (2016).
- 279 21 Di Marco, M., Venter, O., Possingham, H. P. & Watson, J. E. M. Changes in human
- footprint drive changes in species extinction risk. *Nature Communications* **9**, 4621,
- doi:10.1038/s41467-018-07049-5 (2018).
- Lenzen, M. et al. International trade drives biodiversity threats in developing nations.
- 283 *Nature* **486**, 109-112, doi:10.1038/nature11145 (2012).
- Newbold, T. et al. Has land use pushed terrestrial biodiversity beyond the planetary
- boundary? A global assessment. Science 353, 288-291, doi:10.1126/science.aaf2201
- 286 (2016).
- 287 24 Sanderson, E. W., Walston, J. & Robinson, J. G. From bottleneck to breakthrough:
- urbanization and the future of biodiversity conservation. *Bioscience* **68**, 412-426,
- doi:10.1093/biosci/biy039 (2018).
- 290 25 Miller, D. C., Agrawal, A. & Roberts, J. T. Biodiversity, governance, and the
- allocation of international aid for conservation. *Conservation Letters* **6**, 12-20,
- 292 doi:10.1111/j.1755-263X.2012.00270.x (2013).
- 293 26 Baynham-Herd, Z., Amano, T., Sutherland, W. J. & Donald, P. F. Governance
- 294 explains variation in national responses to the biodiversity crisis. *Environmental*
- 295 *Conservation* **45**, 407-418, doi:10.1017/S037689291700056X (2018).
- 296 27 Ellis, E. C. & Mehrabi, Z. Half Earth: promises, pitfalls, and prospects of dedicating
- Half of Earth's land to conservation. Current Opinion in Environmental Sustainability
- 38, 22-30, doi:https://doi.org/10.1016/j.cosust.2019.04.008 (2019).
- 299 28 United Nations. *Life on land: why it matters*, <
- 300 https://www.un.org/sustainabledevelopment/wp-content/uploads/2016/08/15 Why-it-
- Matters Goal15 Life-on-Land 3p.pdf > (2018).
- 302 29 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services
- 303 (IPBES). Summary for policymakers of the global assessment report on biodiversity

304		and ecosystem services of the Intergovernmental Science-Policy Platform on		
305		Biodiversity and Ecosystem Services,		
306		https://www.ipbes.net/sites/default/files/downloads/spm_unedited_advance_for_post		
307		ing_htn.pdf> (2019).		
308	30	Visconti, P. et al. Protected area targets post-2020. Science 364, 239-241,		
309		doi:10.1126/science.aav6886 (2019).		
310	31	Venter, O. et al. Sixteen years of change in the global terrestrial human footprint and		
311		implications for biodiversity conservation. Nature Communications 7, 12558,		
312		doi:10.1038/ncomms12558 (2016).		
313	32	Jones, K. R. et al. One-third of global protected land is under intense human pressure.		
314		Science 360, 788-791, doi:10.1126/science.aap9565 (2018).		
315	33	Olson, D. M. <i>et al.</i> Terrestrial ecoregions of the world: a new map of life on Earth.		
316		<i>Bioscience</i> 51 , 933-938, doi:10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2		
317		(2001).		
318	34	United Nations Development Programme. <i>Human Development Index</i> ,		
319	J -	http://hdr.undp.org/en/content/human-development-index-hdi (2018).		
320	35	Transparency International. Corruption Perceptions Index 2017. Transparency		
321		International is licensed under CC-BY-ND 4.0.,		
322		https://www.transparency.org/news/feature/corruption_perceptions_index_2017#res		
323		earch> (2018).		
324	36	Transparency International. How corruption weakens democracy,		
325		https://www.transparency.org/news/feature/cpi_2018_global_analysis (2019).		
326				
327	Figure	elegends		
328	Fig. 1.	Potential contributions of countries to GNNL. The proportion of natural ecosystems		
329	(Human Footprint value <4) remaining per country varies enormously, as does variation in			
330	the depletion among different ecosystems (Gini coefficient; see Methods). The minimum			
331	country	country-level contribution to a GNNL goal must reflect this, as well as the absolute area of		
332	natural ecosystems remaining (Fig. 1b). Ecosystem depletion must be considered alongside			
333	other factors in setting targets; e.g., the number of threatened species according to the IUCN			

334	Red List of Threatened Species (for fully-assessed taxa only - mammals, birds and
335	amphibians) relates only weakly to retention of ecosystems ($R = 0.17$; d.f. 169; $P = 0.0279$; R
336	version 3.5.1; Fig. 1a).
337	
338	Fig. 2. The degree of human development should affect minimum country-level contributions
339	to achievement of GNNL, such that high HDI countries commit to at least country-level
340	NNL. Bubble size reflects the Corruption Perceptions Index (2017) for each country; see
341	Methods.



