

Monte-Carlo simulations - Conditional Poisson Distribution

March 26, 2019

Contents


1	Description	1
2	Bias function	2
3	Bias of $\hat{\lambda}$	5
4	CV of $\hat{\lambda}$	8

1 Description


To investigate small-sample performance of the bias correction, we carried out Monte-Carlo simulations. To this purpose, we generate datasets according to description of the model. A dataset \mathbf{X}^k constitutes of N samples where every sample is a 0-1 vector which carries a randomly generated MOI from conditional Poisson distribution. The sample sizes N are chosen in an increasing fashion in order to see the improvement in performance in an asymptotic manner. For each combination of values for set of model parameters, $S = 10,000$ datasets are generated. Afterwards, results for each dataset is derived, i.e., MLE, TBCMLe (theoretical bias-corrected MLE) and HBCMLe (heuristic bias-corrected MLE). Finally, considering the empirical distribution of these statistics, the mean and variance of each statistic from 10,000 datasets are calculated.

2 Bias function


A




B




C



D



E



F






Figure 1: **Bias function.** The figure shows the behavior of average approximation of theoretical bias from $S = 10,000$ simulated datasets (solid lines) versus the theoretical bias at the true parameter (long-dashed lines). Different sample sizes are specified with different colors.


A




B




C



D



E



F






Figure 2: **Bias function.** Same as Figure 1


A



B



C



D






Figure 3: **Bias function.** Same as Figure 1

3 Bias of $\hat{\lambda}$


A




B




C



D



E



F






Figure 4: **Bias in %**-The figure shows the bias in % of different versions of bias-corrected MLE of mean MOI parameter along with MLE of $\psi = \frac{\lambda}{1-e^{-\lambda}}$ for different lineage frequency distributions. Different colors correspond to different sample sizes N . The solid, long-dashed and dot-dashed lines correspond to MLE, TBCMLE (theoretical bias-corrected MLE) and HBCMLE (heuristic bias-corrected MLE), respectively.


A




B




C



D



E



F




Figure 5: Same as Figure 4






Figure 6: Same as Figure 4

4 CV of $\hat{\lambda}$


A




B




C



D



E



F






Figure 7: Same as Figure 4 but for coefficient of variation.


A




B




C



D



E



F




Figure 8




Figure 9