


# Our Universe is Chaotic (Probably)

Hashir H. Khan

13 May 2023

# Our World is Chaotic




(a) Double Pendulum




(b) Turbulent Flow

Figure: Chaos in mechanical and fluid motion.

# Our World is Chaotic



(a) Cardiac Arrhythmia



(b) Logistic Map

Figure: Chaos in the human body and population models.

# Our World is Chaotic

Our world, at large, is characterized by **nonlinear, chaotic systems**.

Question

Answer

# Our World is Chaotic

Our world, at large, is characterized by **nonlinear, chaotic systems**. In fact, it is hard to find macroscopic systems which do **not** have a capacity for chaos.

Question

Answer

# Our World is Chaotic

Our world, at large, is characterized by **nonlinear, chaotic systems**. In fact, it is hard to find macroscopic systems which do **not** have a capacity for chaos.

## Question

But what is **chaos**? Is the universe itself chaotic?

## Answer

# Our World is Chaotic

Our world, at large, is characterized by **nonlinear, chaotic systems**. In fact, it is hard to find macroscopic systems which do **not** have a capacity for chaos.

## Question

But what is **chaos**? Is the universe itself chaotic?

## Answer

This and lots more to follow in the talk.

# Structure of the Talk

## 1 The Bianchi IX Cosmology



# Structure of the Talk

- 1 The Bianchi IX Cosmology
- 2 Measures of Chaos

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - ① The BKL Approximation

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - ① The BKL Approximation
  - ② Painleve Analysis

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - i The BKL Approximation
  - ii Painleve Analysis
  - iii Lyapunov Exponents

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - ❶ The BKL Approximation
  - ❷ Painleve Analysis
  - ❸ Lyapunov Exponents
  - ❹ Fractals

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - ❶ The BKL Approximation
  - ❷ Painleve Analysis
  - ❸ Lyapunov Exponents
  - ❹ Fractals
  - ❺ Curvature

# Structure of the Talk

- ① The Bianchi IX Cosmology
- ② Measures of Chaos
  - ❶ The BKL Approximation
  - ❷ Painleve Analysis
  - ❸ Lyapunov Exponents
  - ❹ Fractals
  - ❺ Curvature
- ③ Implications For Our Universe

# The Bianchi IX Cosmology

- Characterized by **homogeneity** and **anisotropy**.



Figure: Example of a homogeneous, anisotropic shape.



# The Bianchi IX Cosmology

- Characterized by **homogeneity** and **anisotropy**.
- Homogeneous: Every point location looks the same.



Figure: Example of a homogeneous, anisotropic shape.

# The Bianchi IX Cosmology

- Characterized by **homogeneity** and **anisotropy**.
- Homogeneous: Every point location looks the same.
- Anisotropic: Does not look the same in all directions.



Figure: Example of a homogeneous, anisotropic shape.

# The Bianchi IX Cosmology

- The metric:

$$ds^2 = -dt^2 + \sum_k (l_k)^2 \sigma_k^2.$$

# The Bianchi IX Cosmology

- The metric:

$$ds^2 = -dt^2 + \sum_k (l_k)^2 \sigma_k^2.$$

- Here,  $k = 1, 2, 3$  where, (in Euler angle coordinates:  $0 \leq \psi \leq 4\pi$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \phi \leq 2\pi$ )
  - $\sigma_1 = (0, \sin \psi, -\cos \psi \sin \theta)$ ,
  - $\sigma_2 = (0, \cos \psi, \sin \psi \sin \theta)$ ,
  - $\sigma_3 = (-1, 0, \cos \theta)$ ,

# The Bianchi IX Cosmology

- The metric:

$$ds^2 = -dt^2 + \sum_k (l_k)^2 \sigma_k^2.$$

- Here,  $k = 1, 2, 3$  where, (in Euler angle coordinates:  $0 \leq \psi \leq 4\pi$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \phi \leq 2\pi$ )
  - $\sigma_1 = (0, \sin \psi, -\cos \psi \sin \theta)$ ,
  - $\sigma_2 = (0, \cos \psi, \sin \psi \sin \theta)$ ,
  - $\sigma_3 = (-1, 0, \cos \theta)$ ,
- The  $l_k$ 's, henceforth denoted by  $a(t)$ ,  $b(t)$ ,  $c(t)$ , are the scale factors.

# The Bianchi IX Cosmology

- The three equations of evolution are

$$\frac{d^2}{d\tau^2}(\ln a^2) = (b^2 - c^2)^2 - a^4$$

and cyclically,  $a \rightarrow b \rightarrow c \rightarrow a$ .

# The Bianchi IX Cosmology

- The three equations of evolution are

$$\frac{d^2}{d\tau^2}(\ln a^2) = (b^2 - c^2)^2 - a^4$$

and cyclically,  $a \rightarrow b \rightarrow c \rightarrow a$ .

- Here, the new coordinate time  $\tau$  is related to  $t$  by  $dt = abc d\tau$ .

# The Bianchi IX Cosmology

- The three equations of evolution are

$$\frac{d^2}{d\tau^2}(\ln a^2) = (b^2 - c^2)^2 - a^4$$

and cyclically,  $a \rightarrow b \rightarrow c \rightarrow a$ .

- Here, the new coordinate time  $\tau$  is related to  $t$  by  $dt = abc d\tau$ .
- Aim is to go back in time to the initial singularity ( $t = 0$ ,  $\tau = -\infty$ ) and observe **chaos** in the scale factors.




# Measures of Chaos


## Question

What is **chaos**?

## Answer



(a) No Chaos



(b) Chaos


# Measures of Chaos

## Question


What is **chaos**?

## Answer

Roughly speaking, chaos is a sensitive dependence on initial conditions.



(a) No Chaos



(b) Chaos

# Measures of Chaos


## Question

What is **chaos**?


## Answer

Roughly speaking, chaos is a sensitive dependence on initial conditions.

A system is chaotic if, given initial conditions with a small uncertainty, we are unable to predict the state of the system after a (long enough) finite time interval.



(a) No Chaos



(b) Chaos

# Break for Remembering Stuff

If you forgot everything up till now, just remember the following:

- Our universe is characterized by homogeneous, anisotropic Bianchi IX cosmology.

# Break for Remembering Stuff

If you forgot everything up till now, just remember the following:

- Our universe is characterized by homogeneous, anisotropic Bianchi IX cosmology.
- Chaos is a sensitive dependence on initial conditions.

# The BKL Approximation

- Belinski–Khalatnikov–Lifshitz (BKL) at the Landau Institute for Theoretical Physics.

# The BKL Approximation

- Belinski–Khalatnikov–Lifshitz (BKL) at the Landau Institute for Theoretical Physics.
- Start with the Kasner (Bianchi I) solution:

$$ds^2 = -dt^2 + \sum_k t^{2p_k} dx_k^2,$$

where  $-\frac{1}{3} \leq p_1 \leq 0 \leq p_2 \leq \frac{2}{3} \leq p_3 \leq 1$  and  $\sum_k p_k = \sum_k p_k^2 = 1$ .

# The BKL Approximation

- Belinski–Khalatnikov–Lifshitz (BKL) at the Landau Institute for Theoretical Physics.
- Start with the Kasner (Bianchi I) solution:

$$ds^2 = -dt^2 + \sum_k t^{2p_k} dx_k^2,$$

where  $-\frac{1}{3} \leq p_1 \leq 0 \leq p_2 \leq \frac{2}{3} \leq p_3 \leq 1$  and  $\sum_k p_k = \sum_k p_k^2 = 1$ .

- This universe expands in one direction and contracts in two as we go back in time.



# The BKL Approximation

- Belinski–Khalatnikov–Lifshitz (BKL) at the Landau Institute for Theoretical Physics.
- Start with the Kasner (Bianchi I) solution:

$$ds^2 = -dt^2 + \sum_k t^{2p_k} dx_k^2,$$

where  $-\frac{1}{3} \leq p_1 \leq 0 \leq p_2 \leq \frac{2}{3} \leq p_3 \leq 1$  and  $\sum_k p_k = \sum_k p_k^2 = 1$ .

- This universe expands in one direction and contracts in two as we go back in time.
- Introduce the BKL parameter  $u$ :

$$p_1 = -\frac{u}{u^2 + u + 1}, \quad p_2 = \frac{u+1}{u^2 + u + 1}, \quad p_3 = \frac{u(u+1)}{u^2 + u + 1}$$

# The BKL Approximation

- Describe Bianchi IX using a series of Kasner epochs.

# The BKL Approximation

- Describe Bianchi IX using a series of Kasner epochs.
- The epoch-change map (called the **Gauss Map**) is

$$u_{n+1} = \frac{1}{u_n - [u_n]},$$

where  $[u_n]$  is the fractional part of  $u_n$ .

# The BKL Approximation

- Describe Bianchi IX using a series of Kasner epochs.
- The epoch-change map (called the **Gauss Map**) is

$$u_{n+1} = \frac{1}{u_n - [u_n]},$$

where  $[u_n]$  is the fractional part of  $u_n$ .

- Slight aside: the **metric entropy**  $h$  of a discrete map characterizes its chaoticity by describing how nearby trajectories diverge on average.

$$h = \frac{\langle \ln(\Delta_t / \Delta_0) \rangle}{t}$$

# The BKL Approximation

- Describe Bianchi IX using a series of Kasner epochs.
- The epoch-change map (called the **Gauss Map**) is

$$u_{n+1} = \frac{1}{u_n - [u_n]},$$

where  $[u_n]$  is the fractional part of  $u_n$ .

- Slight aside: the **metric entropy**  $h$  of a discrete map characterizes its chaoticity by describing how nearby trajectories diverge on average.

$$h = \frac{\langle \ln(\Delta_t / \Delta_0) \rangle}{t}$$

- Using some measure theory, it can be found that

$$h = \frac{\pi^2}{6(\ln 2)^2}$$

# The BKL Approximation

- Describe Bianchi IX using a series of Kasner epochs.
- The epoch-change map (called the **Gauss Map**) is

$$u_{n+1} = \frac{1}{u_n - [u_n]},$$

where  $[u_n]$  is the fractional part of  $u_n$ .

- Slight aside: the **metric entropy**  $h$  of a discrete map characterizes its chaoticity by describing how nearby trajectories diverge on average.

$$h = \frac{\langle \ln(\Delta_t / \Delta_0) \rangle}{t}$$

- Using some measure theory, it can be found that

$$h = \frac{\pi^2}{6(\ln 2)^2}$$

- Since  $h > 0$  implies exponential divergence, the **Gauss Map** is chaotic.

# The BKL Approximation

- Numerical estimates show that the BKL approximation describes Bianchi IX very accurately near the singularity.

# The BKL Approximation

- Numerical estimates show that the BKL approximation describes Bianchi IX very accurately near the singularity.
- There is a **drawback** however. For all its accuracy, it remains an approximation and tells nothing about the chaoticity of Bianchi IX itself.



# Painleve Analysis

- A test to characterize whether a system is **integrable**.

# Painleve Analysis

- A test to characterize whether a system is **integrable**.
- Integrability: System has enough first integrals.

# Painleve Analysis

- A test to characterize whether a system is **integrable**.
- Integrability: System has enough first integrals.
- The only **movable singularities** of the general solution are poles  $\implies$  system is integrable.

# Painleve Analysis

- A test to characterize whether a system is **integrable**.
- Integrability: System has enough first integrals.
- The only **movable singularities** of the general solution are poles  $\implies$  system is integrable.
- Movable singularity: a singularity in the solution which depends on the value of the arbitrary coefficient.

# Painleve Analysis

- A test to characterize whether a system is **integrable**.
- Integrability: System has enough first integrals.
- The only **movable singularities** of the general solution are poles  $\implies$  system is integrable.
- Movable singularity: a singularity in the solution which depends on the value of the arbitrary coefficient.
- Example:

$$\frac{du}{dx} + \frac{u^2}{x} = 0, \quad u = \frac{1}{c + \ln x}$$

# Painleve Analysis

- After performing a Painleve test, we find that the general solution of the Bianchi IX model has essential singularities of the form  $\sim e^{1/\tau}$ .

# Painleve Analysis

- After performing a Painleve test, we find that the general solution of the Bianchi IX model has essential singularities of the form  $\sim e^{1/\tau}$ .
- Bianchi IX is therefore non-integrable.

# Painleve Analysis

- After performing a Painleve test, we find that the general solution of the Bianchi IX model has essential singularities of the form  $\sim e^{1/\tau}$ .
- Bianchi IX is therefore non-integrable.
- Drawback: Non-integrability does not imply chaos.



# Lyapunov Exponents

- Lyapunov exponents are the metric entropy for continuous systems.

# Lyapunov Exponents

- Lyapunov exponents are the metric entropy for continuous systems.
- Certain numerical estimates show that there is a positive Lyapunov exponent in Bianchi IX.

# Lyapunov Exponents

- Lyapunov exponents are the metric entropy for continuous systems.
- Certain numerical estimates show that there is a positive Lyapunov exponent in Bianchi IX.
- Other estimates however show that the largest Lyapunov exponent is 0.

# Lyapunov Exponents

- Lyapunov exponents are the metric entropy for continuous systems.
- Certain numerical estimates show that there is a positive Lyapunov exponent in Bianchi IX.
- Other estimates however show that the largest Lyapunov exponent is 0.
- These contradictory results therefore do not shed light on any chaos present in Bianchi IX.

# Fractals

- Fractals are self-similar structures.




Figure: The Sierpinski Triangle.

# Fractals

- Fractals are self-similar structures.
- For each fractal, we can specify a fractal dimension.




Figure: The Sierpinski Triangle.

# Fractals

- **Fractals** are self-similar structures.
- For each fractal, we can specify a **fractal dimension**.
- An example is the **box-counting dimension** which describes the rate at which the number of boxes required to completely cover a fractal grows compared to the rate at which the size of each box shrinks.




Figure: The Sierpinski Triangle.

- By identifying a set of initial conditions in phase space, which when evolved backwards in time, show an oscillatory behaviour close to the singularity, we can identify our fractal set.



# Fractals

- By identifying a set of initial conditions in phase space, which when evolved backwards in time, show an oscillatory behaviour close to the singularity, we can identify our fractal set.
- If evolution of these states causes them to remain in this set, we have an invariant set.

- By identifying a set of initial conditions in phase space, which when evolved backwards in time, show an oscillatory behaviour close to the singularity, we can identify our fractal set.
- If evolution of these states causes them to remain in this set, we have an invariant set.
- Numerical estimates have shown that the fractal dimension of this invariant set is  $1.86 \pm 0.01$ .

# Fractals

- By identifying a set of initial conditions in phase space, which when evolved backwards in time, show an oscillatory behaviour close to the singularity, we can identify our fractal set.
- If evolution of these states causes them to remain in this set, we have an invariant set.
- Numerical estimates have shown that the fractal dimension of this invariant set is  $1.86 \pm 0.01$ .
- Dimension is a fraction  $\implies$  invariant set is a fractal  $\implies$  chaos exists.

- By identifying a set of initial conditions in phase space, which when evolved backwards in time, show an oscillatory behaviour close to the singularity, we can identify our fractal set.
- If evolution of these states causes them to remain in this set, we have an invariant set.
- Numerical estimates have shown that the fractal dimension of this invariant set is  $1.86 \pm 0.01$ .
- Dimension is a fraction  $\implies$  invariant set is a fractal  $\implies$  chaos exists.
- **Drawback:** As with all other numerical methods, this remains analytically unproven.

- Introduce a Hamiltonian for Bianchi IX:

$$2H = -p_t^2 + p_\zeta^2 + \left( \frac{p_\phi}{\sinh \zeta} \right)^2$$

- Introduce a Hamiltonian for Bianchi IX:

$$2H = -p_t^2 + p_\zeta^2 + \left( \frac{p_\phi}{\sinh \zeta} \right)^2$$

- Here, we have ignored the potential term in the Hamiltonian.

- Introduce a Hamiltonian for Bianchi IX:

$$2H = -p_t^2 + p_\zeta^2 + \left( \frac{p_\phi}{\sinh \zeta} \right)^2$$

- Here, we have ignored the potential term in the Hamiltonian.
- Note that the spatial part of the Hamiltonian includes the inverse of the hyperbolic plane metric:

$$ds^2 = dx^2 + (\sinh^2 x) dy^2$$

- Introduce a Hamiltonian for Bianchi IX:

$$2H = -p_t^2 + p_\zeta^2 + \left( \frac{p_\phi}{\sinh \zeta} \right)^2$$

- Here, we have ignored the potential term in the Hamiltonian.
- Note that the spatial part of the Hamiltonian includes the inverse of the hyperbolic plane metric:

$$ds^2 = dx^2 + (\sinh^2 x) dy^2$$

- Solutions are geodesics on the hyperbolic plane.



- Using results from measure theory and differential geometry, it has been shown that this geodesic motion is chaotic.

- Using results from measure theory and differential geometry, it has been shown that this geodesic motion is chaotic.
- In particular, a small finite set will spread almost everywhere on the hyperbolic plane under the equations of motion.

- Using results from measure theory and differential geometry, it has been shown that this geodesic motion is chaotic.
- In particular, a small finite set will spread almost everywhere on the hyperbolic plane under the equations of motion.
- Therefore, the evolution of Bianchi IX is chaotic.

- Using results from measure theory and differential geometry, it has been shown that this geodesic motion is chaotic.
- In particular, a small finite set will spread almost everywhere on the hyperbolic plane under the equations of motion.
- Therefore, the evolution of Bianchi IX is chaotic.
- **Drawback:** Ignoring the potential term is an approximation which may not be valid.

# Implications For Our Universe

- If the Bianchi IX model describes our universe and is indeed chaotic, we cannot specify a set of initial conditions which brought us to our universe today.

# Implications For Our Universe

- If the Bianchi IX model describes our universe and is indeed chaotic, we cannot specify a set of initial conditions which brought us to our universe today.
- In particular, we CANNOT tell how the universe began.

# The End

Our universe is (probably) chaotic.

~ Hashir H. Khan  
(the H stands for Hashir H. Khan)