

HW8

Chapters 11 and 12

TA Solutions

The code below just loads some packages and makes it so that enough digits are printed that you won't get confused by rounding errors.

```
library(dplyr) # functions like summarize
library(ggplot2) # for making plots
library(gridExtra)
library(GGally)
library(readr)
library(car)
options("pillar.sigfig" = 10) # print 10 significant digits in summarize output
```

Problem 1: Galapagos (Adapted from Sleuth3 12.20)

Quote from book:

The data [read in below] come from a 1973 study. (Data from M. P. Johnson and P. H. Raven, "Species Number and Endemism: The Galapagos Archipelago Revisited," *Science* 179 (1973): 893-5.) The number of species on an island is known to be related to the island's area. Of interest is what other variables are also related to the number of species, after island area is accounted for. The data set includes the following variables:

- Island: a character vector indicating the island
- Total: total number of observed species
- Native: number of native species
- Area: area (km²)
- Elev: elevation (m)
- DistNear: distance from nearest island (km)
- DistSc: distance from Santa Cruz (km)
- AreaNear: area of nearest island (km²)

In this analysis, our response variable is **Native**, the number of native species. You will use **Area**, **Elev**, **DistNear**, **DistSc**, and **AreaNear** as possible explanatory variables.

```
galapagos <- read_csv("http://www.evanlray.com/data/sleuth3/ex1220_galapagos.csv")


## Parsed with column specification:
## cols(
##   Island = col_character(),
##   Total = col_double(),
##   Native = col_double(),
##   Area = col_double(),
##   Elev = col_double(),
##   DistNear = col_double(),
##   DistSc = col_double(),
##   AreaNear = col_double()
## )
```

```
head(galapagos)
```

```
## # A tibble: 6 x 8
##   Island      Total Native  Area  Elev DistNear DistSc AreaNear
##   <chr>      <dbl>  <dbl> <dbl> <dbl>  <dbl>  <dbl>  <dbl>
## 1 Baltra      58    23 25.09  332    0.6    0.6    1.84
## 2 Bartolome   31    21  1.24  109    0.6   26.3  572.33
## 3 Caldwell     3     3  0.21  114    2.8   58.7    0.78
## 4 Champion    25     9  0.1   46    1.9   47.4    0.18
## 5 Coamano      2     1  1.05  130    1.9    1.9  903.82
## 6 Daphne Major 18    11  0.34  119     8     8    1.84
```


(a) Make a pairs plot of the data including only the variables you will use in your analysis (with the response variable last).

```
ggpairs(galapagos %>% select(Area, Elev, DistNear, DistSc, AreaNear, Native))
```



(b) Identify a set of transformations for all variables in the model so that the regression conditions appear to be fairly well satisfied. In doing this, consider pairs plots of the transformed data and plots of residuals vs explanatory variables in a regression model including all transformed explanatory variables. (You should have both of these types of plots.) After this step, you should feel fairly confident that any models you fit will either have approximately linear relationships among transformed variables, or know how you will handle non-linearity by adding polynomial terms in the model. You should also feel confident that the variance of residuals is fairly constant across values of explanatory variables. Note that DistSc includes some 0 values. A common trick in cases like this is to add 1 to the observed values of that variable before doing transformations (this means that things like a log transformation are an option).

```
galapagos_transformed <- galapagos %>%
  transmute(
    log_Area = log(Area),
    log_Elev = log(Elev),
    log_DistNear = log(DistNear),
    log_DistSc = log(DistSc + 1),
    log_AreaNear = log(AreaNear),
    Native_0.25 = Native^0.25
  )
ggpairs(galapagos_transformed)
```




```

lm_full <- lm(Native_0.25 ~ log_Area + log_Elev + log_DistNear + log_DistSc + log_AreaNear,
  data = galapagos_transformed)
galapagos_transformed <- galapagos_transformed %>%
  mutate(
    resid = residuals(lm_full)
  )
p1 <- ggplot(data = galapagos_transformed, mapping = aes(x = log_Area, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p2 <- ggplot(data = galapagos_transformed, mapping = aes(x = log_Elev, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p3 <- ggplot(data = galapagos_transformed, mapping = aes(x = log_DistNear, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p4 <- ggplot(data = galapagos_transformed, mapping = aes(x = log_DistSc, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p5 <- ggplot(data = galapagos_transformed, mapping = aes(x = log_AreaNear, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p6 <- ggplot(data = galapagos_transformed, mapping = aes(x = resid)) +
  geom_density() +
  theme_bw()
grid.arrange(p1, p2, p3, p4, p5, p6)


## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'

```



Grader note: a log transformation for Native is also OK. The fourth root transformation looks slightly better, but log also works, and has interpretation advantages. The output with `log(Native)` as the response is included as the second output for each part of the homework for reference.


```
galapagos_transformed2 <- galapagos %>%
  transmute(
    log_Area = log(Area),
    log_Elev = log(Elev),
    log_DistNear = log(DistNear),
    log_DistSc = log(DistSc + 1),
    log_AreaNear = log(AreaNear),
    Native_log = log(Native)
  )
ggpairs(galapagos_transformed2)
```



```
lm_full2 <- lm(Native_log ~ log_Area + log_Elev + log_DistNear + log_DistSc + log_AreaNear,
  data = galapagos_transformed2)
galapagos_transformed2 <- galapagos_transformed2 %>%
  mutate(
    resid = residuals(lm_full2)
  )
p1 <- ggplot(data = galapagos_transformed2, mapping = aes(x = log_Area, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p2 <- ggplot(data = galapagos_transformed2, mapping = aes(x = log_Elev, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p3 <- ggplot(data = galapagos_transformed2, mapping = aes(x = log_DistNear, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p4 <- ggplot(data = galapagos_transformed2, mapping = aes(x = log_DistSc, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
p5 <- ggplot(data = galapagos_transformed2, mapping = aes(x = log_AreaNear, y = resid)) +
  geom_point() +
  geom_smooth() +
  theme_bw()
```

```
p6 <- ggplot(data = galapagos_transformed2, mapping = aes(x = resid)) +
  geom_density() +
  theme_bw()
grid.arrange(p1, p2, p3, p4, p5, p6)
```

```
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
## `geom_smooth()` using method = 'loess' and formula 'y ~ x'
```




(c) Check for influential observations, outliers or high leverage observations.


With Native^{0.25} transformation:

```
galapagos_transformed <- galapagos_transformed %>%
  mutate(
    obs_index = row_number(),
    h = hatvalues(lm_full),
    studres = rstudent(lm_full),
    D = cooks.distance(lm_full)
  )
ggplot(data = galapagos_transformed, mapping = aes(x = obs_index, y = h)) +
  geom_hline(yintercept = 2 * 6 / nrow(galapagos_transformed)) +
```


```
geom_point() +  
theme_bw()
```




```
ggplot(data = galapagos_transformed, mapping = aes(x = obs_index, y = studres)) +  
  geom_point() +  
  theme_bw()
```

```
ggplot(data = galapagos_transformed, mapping = aes(x = obs_index, y = D)) +  
  geom_point() +  
  theme_bw()
```




```
galapagos_transformed <- galapagos_transformed %>%  
  mutate(suspicious = obs_index %in% c(5, 15))  
ggpairs(galapagos_transformed, mapping = aes(color = suspicious), columns = 1:6)
```




With log(Native) transformation:


```
galapagos_transformed2 <- galapagos_transformed2 %>%
  mutate(
    obs_index = row_number(),
    h = hatvalues(lm_full12),
    studres = rstudent(lm_full12),
    D = cooks.distance(lm_full12)
  )
ggplot(data = galapagos_transformed2, mapping = aes(x = obs_index, y = h)) +
  geom_hline(yintercept = 2 * 6 / nrow(galapagos_transformed2)) +
  geom_point() +
  theme_bw()
```




```
ggplot(data = galapagos_transformed2, mapping = aes(x = obs_index, y = studres)) +  
  geom_point() +  
  theme_bw()
```



```
ggplot(data = galapagos_transformed2, mapping = aes(x = obs_index, y = D)) +  
  geom_point() +  
  theme_bw()
```



```
galapagos_transformed2 <- galapagos_transformed2 %>%  
  mutate(suspicious = obs_index %in% c(5, 15))  
ggpairs(galapagos_transformed2, mapping = aes(color = suspicious), columns = 1:6)
```




(d) Use all subsets regression to identify a set of models with similar ability to model these data well, based on your transformed variables. If necessary, perform this step both with and without the outliers or influential observations included.

With Native^{0.25} transformation:

```
library(leaps)
```

```
## Warning: package 'leaps' was built under R version 3.6.3
```

```
candidate_models <- regsubsets(Native_0.25 ~ log_Area + log_Elev + log_DistNear +  
  log_DistSc + log_AreaNear, data = galapagos_transformed)  
plot(candidate_models)
```



```
summary(candidate_models)
```

```
## Subset selection object
## Call: regsubsets.formula(Native_0.25 ~ log_Area + log_Elev + log_DistNear +
##   log_DistSc + log_AreaNear, data = galapagos_transformed)
## 5 Variables (and intercept)
##           Forced in Forced out
## log_Area      FALSE      FALSE
## log_Elev       FALSE      FALSE
## log_DistNear   FALSE      FALSE
## log_DistSc     FALSE      FALSE
## log_AreaNear   FALSE      FALSE
## 1 subsets of each size up to 5
## Selection Algorithm: exhaustive
##           log_Area log_Elev log_DistNear log_DistSc log_AreaNear
## 1  ( 1 ) "*"      " "      " "          " "          " "
## 2  ( 1 ) "*"      " "      "*"          " "          " "
## 3  ( 1 ) "*"      " "      "*"          " "          "*"
## 4  ( 1 ) "*"      "*"      "*"          " "          "*"
## 5  ( 1 ) "*"      "*"      "*"          "*"          "*"


```

```
summary(candidate_models)$bic
```

```
## [1] -36.15035 -34.91195 -33.51806 -30.22219 -26.89620
```



```
candidate_models2 <- regsubsets(Native_0.25 ~ log_Area + log_Elev + log_DistNear +
  log_DistSc + log_AreaNear, data = galapagos_transformed %>% filter(!suspicious))
plot(candidate_models2)
```



```
summary(candidate_models2)
```

```
## Subset selection object
## Call: regsubsets.formula(Native_0.25 ~ log_Area + log_Elev + log_DistNear +
##   log_DistSc + log_AreaNear, data = galapagos_transformed %>%
##   filter(!suspicious))
## 5 Variables (and intercept)
##           Forced in Forced out
## log_Area      FALSE      FALSE
## log_Elev      FALSE      FALSE
## log_DistNear  FALSE      FALSE
## log_DistSc    FALSE      FALSE
## log_AreaNear  FALSE      FALSE
## 1 subsets of each size up to 5
## Selection Algorithm: exhaustive
##           log_Area log_Elev log_DistNear log_DistSc log_AreaNear
## 1  ( 1 ) "*"      " "      " "          " "          " "
## 2  ( 1 ) "*"      " "      " "          "*"          " "
## 3  ( 1 ) "*"      "*"      " "          "*"          " "
## 4  ( 1 ) "*"      "*"      "*"          "*"          " "
## 5  ( 1 ) "*"      "*"      "*"          "*"          "*"

```

```
summary(candidate_models2)$bic
```

```
## [1] -40.33967 -42.38225 -40.04481 -36.84316 -33.51199
```

When all observations are included, the following three models have roughly similar performance:

Model 1: log Area, log DistNear, and log AreaNear as explanatory variables

Model 2: log Area and log DistNear as explanatory variables

Model 3: log Area as the only explanatory variable

When one outlier and one high leverage observation are omitted, the following three models have roughly similar performance:


Model 1: log Area, log DistSc, and log Elev as explanatory variables

Model 2: log Area and log DistSc as explanatory variables

Model 3: log Area as the only explanatory variable

With log(Native) transformation:

```
# library(leaps)
candidate_models2_1 <- regsubsets(Native_log ~ log_Area + log_Elev + log_DistNear +
  log_DistSc + log_AreaNear, data = galapagos_transformed2)
plot(candidate_models2_1)
```



```
summary(candidate_models2_1)
```


```
## Subset selection object
## Call: regsubsets.formula(Native_log ~ log_Area + log_Elev + log_DistNear +
##       log_DistSc + log_AreaNear, data = galapagos_transformed2)
## 5 Variables (and intercept)
##           Forced in Forced out
## log_Area      FALSE      FALSE
## log_Elev      FALSE      FALSE
## log_DistNear  FALSE      FALSE
## log_DistSc    FALSE      FALSE
## log_AreaNear  FALSE      FALSE
## 1 subsets of each size up to 5
## Selection Algorithm: exhaustive
##           log_Area log_Elev log_DistNear log_DistSc log_AreaNear
## 1  ( 1 ) "*"      " "      " "          " "          " "
## 2  ( 1 ) "*"      " "      " "          " "          "*"
## 3  ( 1 ) "*"      " "      "*"          " "          "*"
## 4  ( 1 ) "*"      "*"      "*"          " "          "*"
## 5  ( 1 ) "*"      "*"      "*"          "*"          "*"

```

```
summary(candidate_models2_1)$bic
```

```
## [1] -28.79992 -26.11391 -23.53182 -20.38876 -16.99462
```

```
candidate_models2_2 <- regsubsets(Native_log ~ log_Area + log_Elev + log_DistNear +
  log_DistSc + log_AreaNear, data = galapagos_transformed2 %>% filter(!suspicious))
plot(candidate_models2_2)
```



```
summary(candidate_models2_2)
```

```
## Subset selection object
## Call: regsubsets.formula(Native_log ~ log_Area + log_Elev + log_DistNear +
##   log_DistSc + log_AreaNear, data = galapagos_transformed2 %>%
##   filter(!suspicious))
## 5 Variables (and intercept)
##           Forced in Forced out
## log_Area      FALSE      FALSE
## log_Elev      FALSE      FALSE
## log_DistNear  FALSE      FALSE
## log_DistSc    FALSE      FALSE
## log_AreaNear  FALSE      FALSE
## 1 subsets of each size up to 5
## Selection Algorithm: exhaustive
##           log_Area log_Elev log_DistNear log_DistSc log_AreaNear
## 1 ( 1 ) "*"      " "      " "          " "      " "
## 2 ( 1 ) "*"      " "      " "          "*"      " "
## 3 ( 1 ) "*"      "*"      " "          "*"      " "
## 4 ( 1 ) "*"      "*"      " "          "*"      "*"
## 5 ( 1 ) "*"      "*"      "*"          "*"      "*"

```

```
summary(candidate_models2_2)$bic
```

```
## [1] -36.14739 -36.74069 -34.61666 -31.81287 -28.67545
```

When all observations are included, the following two models have roughly similar performance:

Model 1: log Area as the only explanatory variables (BIC=-28.8)

Model 2: log Area and log AreaNear as explanatory variables (BIC=-26.1)

When one outlier and one high leverage observation are omitted, the following three models have roughly similar performance:

Model 1: log Area, log DistSc, and log Elev as explanatory variables (BIC=-34.6)

Model 2: log Area and log DistSc as explanatory variables (BIC=-36.7)

Model 3: log Area as the only explanatory variable (BIC=-36.1)

Note: Model 2 and Model 3 are the same models identified for the fourth root transformation for Native, both with and without the outlier and high leverage point. If someone makes a log transformation, you don't need to take off points, but make a note that they should investigate a fourth root transformation - it is more stabilizing in this case for outliers and leverage points (removing them from the analysis doesn't have the same impact that it does for the log transformation) They may not really see that until part (d).

(e) Obtain the model fits for all models you identified in part (d) as explaining the data about as well as each other, and print the model summaries.

With $\text{Native}^{0.25}$ transformation

```
fit1 <- lm(Native_0.25 ~ log_Area + log_DistNear + log_AreaNear, data = galapagos_transformed)
summary(fit1)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area + log_DistNear + log_AreaNear,
##     data = galapagos_transformed)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.72395 -0.19768  0.05207  0.17589  0.42334
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.89972    0.07354  25.834 < 2e-16 ***
## log_Area       0.15514    0.01572   9.872 2.77e-10 ***
## log_DistNear -0.05232    0.03292  -1.589   0.124
## log_AreaNear -0.02200    0.01640  -1.341   0.191
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.279 on 26 degrees of freedom
## Multiple R-squared:  0.7921, Adjusted R-squared:  0.7681
## F-statistic: 33.02 on 3 and 26 DF,  p-value: 5.091e-09
```

```
fit2 <- lm(Native_0.25 ~ log_Area + log_DistNear, data = galapagos_transformed)
summary(fit2)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area + log_DistNear, data = galapagos_transformed)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.84087 -0.19373  0.04474  0.21246  0.46082
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.86380    0.06949  26.820 < 2e-16 ***
## log_Area       0.15005    0.01547   9.696 2.74e-10 ***
## log_DistNear -0.04713    0.03317  -1.421   0.167
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2831 on 27 degrees of freedom
## Multiple R-squared:  0.7777, Adjusted R-squared:  0.7613
## F-statistic: 47.24 on 2 and 27 DF,  p-value: 1.524e-09
```

```
fit3 <- lm(Native_0.25 ~ log_Area, data = galapagos_transformed)
summary(fit3)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area, data = galapagos_transformed)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.81591 -0.15749  0.05753  0.23017  0.42722
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.80867    0.05869  30.818 < 2e-16 ***
## log_Area       0.14834    0.01571   9.445 3.34e-10 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2882 on 28 degrees of freedom
## Multiple R-squared:  0.7611, Adjusted R-squared:  0.7526
## F-statistic: 89.21 on 1 and 28 DF,  p-value: 3.342e-10
```

```
fit1a <- lm(Native_0.25 ~ log_Area + log_DistSc + log_AreaNear,
            data = galapagos_transformed %>% filter(!suspicious))
summary(fit1a)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area + log_DistSc + log_AreaNear,
##     data = galapagos_transformed %>% filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
```

```
## -0.36159 -0.15993 0.00683 0.14532 0.36227
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.068e+00  1.095e-01  18.879 6.61e-16 ***
## log_Area      1.471e-01  1.337e-02  11.004 7.36e-11 ***
## log_DistSc    -6.729e-02  2.989e-02  -2.251  0.0338 *
## log_AreaNear -1.513e-05  1.477e-02  -0.001  0.9992
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2347 on 24 degrees of freedom
## Multiple R-squared:  0.846, Adjusted R-squared:  0.8267
## F-statistic: 43.94 on 3 and 24 DF, p-value: 6.651e-10

fit2a <- lm(Native_0.25 ~ log_Area + log_DistSc, data = galapagos_transformed %>% filter(!suspicious))
summary(fit2a)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area + log_DistSc, data = galapagos_transformed %>%
##     filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.36155 -0.15995  0.00682  0.14536  0.36234
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.06787    0.10681   19.36 < 2e-16 ***
## log_Area      0.14711    0.01265   11.63 1.39e-11 ***
## log_DistSc    -0.06729    0.02926   -2.30  0.0301 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.23 on 25 degrees of freedom
## Multiple R-squared:  0.846, Adjusted R-squared:  0.8337
## F-statistic: 68.66 on 2 and 25 DF, p-value: 6.994e-11
```

```
fit3a <- lm(Native_0.25 ~ log_Area, data = galapagos_transformed %>% filter(!suspicious))
summary(fit3a)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area, data = galapagos_transformed %>%
##     filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.41110 -0.18423  0.04419  0.22051  0.37839
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
```

```
## (Intercept)  1.84874    0.05213   35.47 < 2e-16 ***
## log_Area     0.14488    0.01361   10.64 5.67e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2482 on 26 degrees of freedom
## Multiple R-squared:  0.8134, Adjusted R-squared:  0.8062
## F-statistic: 113.3 on 1 and 26 DF,  p-value: 5.674e-11
```

With log(Native transformation):

```
fit2_1 <- lm(Native_log ~ log_Area, data = galapagos_transformed2)
summary(fit2_1)
```

```
##
## Call:
## lm(formula = Native_log ~ log_Area, data = galapagos_transformed2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.2471 -0.2140  0.1364  0.4145  0.9884
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.23288    0.13646  16.363 7.27e-16 ***
## log_Area     0.29155    0.03652   7.984 1.08e-08 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.67 on 28 degrees of freedom
## Multiple R-squared:  0.6948, Adjusted R-squared:  0.6839
## F-statistic: 63.74 on 1 and 28 DF,  p-value: 1.076e-08
```

```
fit2_2 <- lm(Native_log ~ log_Area + log_AreaNear, data = galapagos_transformed2)
summary(fit2_2)
```

```
##
## Call:
## lm(formula = Native_log ~ log_Area + log_AreaNear, data = galapagos_transformed2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.07427 -0.27298  0.04037  0.41418  0.92891
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.27597    0.14733  15.448 6.30e-15 ***
## log_Area     0.29863    0.03778   7.904 1.69e-08 ***
## log_AreaNear -0.03177    0.03937  -0.807   0.427
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```



```
##
## Residual standard error: 0.6743 on 27 degrees of freedom
## Multiple R-squared:  0.702, Adjusted R-squared:  0.6799
## F-statistic: 31.8 on 2 and 27 DF,  p-value: 7.987e-08

fit2_1a <- lm(Native_log ~ log_Area + log_DistSc + log_Elev,
              data = galapagos_transformed2 %>% filter(!suspicious))
summary(fit2_1a)

##
## Call:
## lm(formula = Native_log ~ log_Area + log_DistSc + log_Elev, data = galapagos_transformed2 %>%
##     filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.99171 -0.33379 -0.05635  0.35918  0.87608
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  4.01055    1.25788   3.188 0.003950 **
## log_Area      0.36111    0.07885   4.580 0.000121 ***
## log_DistSc   -0.12303    0.06389  -1.926 0.066087 .
## log_Elev     -0.25903    0.25180  -1.029 0.313873
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.5022 on 24 degrees of freedom
## Multiple R-squared:  0.8196, Adjusted R-squared:  0.797
## F-statistic: 36.34 on 3 and 24 DF,  p-value: 4.383e-09

fit2_2a <- lm(Native_log ~ log_Area + log_DistSc, data = galapagos_transformed2 %>% filter(!suspicious))
summary(fit2_2a)

##
## Call:
## lm(formula = Native_log ~ log_Area + log_DistSc, data = galapagos_transformed2 %>%
##     filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.06862 -0.31592 -0.01162  0.25476  0.81829
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.73900    0.23350  11.73 1.17e-11 ***
## log_Area      0.28514    0.02765  10.31 1.71e-10 ***
## log_DistSc   -0.12406    0.06396  -1.94  0.0638 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.5027 on 25 degrees of freedom
## Multiple R-squared:  0.8116, Adjusted R-squared:  0.7965
## F-statistic: 53.85 on 2 and 25 DF,  p-value: 8.68e-10
```

```
fit2_3a <- lm(Native_log ~ log_Area, data = galapagos_transformed2 %>% filter(!suspicious))
summary(fit2_3a)
```

```
##
## Call:
## lm(formula = Native_log ~ log_Area, data = galapagos_transformed2 %>%
##     filter(!suspicious))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.15996 -0.32394  0.08423  0.35402  0.85969
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.33501    0.11104  21.028 < 2e-16 ***
## log_Area      0.28102    0.02899   9.693 4.04e-10 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.5288 on 26 degrees of freedom
## Multiple R-squared:  0.7832, Adjusted R-squared:  0.7749
## F-statistic: 93.95 on 1 and 26 DF,  p-value: 4.043e-10
```

(f) Summarize what your analysis has to say about the association of each of the explanatory variables in the data set with the response, after accounting for the explanatory variables in your models. Indicate which of your findings are consistent across the various models considered and which depend on the details of your analysis.

With Native^{0.25} transformation:

All of the models with low BIC showed very strong evidence of a positive association between an island's area and the number of native species found on the island, among islands similar to those in this study. This result held whether or not one high leverage observation and one outlier were included. If those two observations were removed, there was some weak evidence of a negative association between distance from Santa Cruz and the number of native species in the population of islands similar to those in this study, after accounting for the area of the island; this finding is not reliable since it depends on the removal of those two observations. After accounting for the size of the island, there was not evidence of an association between any of the other explanatory variables and the number of native species on the island.

With log(Native) transformation:

All of the models with low BIC showed very strong evidence of a positive association between an island's area and the number of native species found on the island, among islands similar to those in this study. This result held whether or not one high leverage observation and one outlier were included. This is the most reliable finding, since this transformation was more sensitive to the removal of observations (outliers/leverage points). Regardless, after accounting for the size of the island, there was not evidence of an association between any of the other explanatory variables and the number of native species on the island. This is consistent with the conclusions using the first response variable transformation, which is encouraging.

(g) In this part we'll think through what's going on in a model that includes only your transformed Elev variable and your transformed Area variable.

i. Fit a model that has your (potentially transformed) Native as the response and your (potentially transformed) Elev and Area variables as explanatory variables. Print the model summary and also use the avPlots function to create added variables plots for these variables.


With $\text{Native}^{0.25}$ transformation:

```
fit_both <- lm(Native_0.25 ~ log_Area + log_Elev, data = galapagos_transformed)
summary(fit_both)
```

```
##
## Call:
## lm(formula = Native_0.25 ~ log_Area + log_Elev, data = galapagos_transformed)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.8167 -0.1842  0.0616  0.2340  0.4390
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   1.95900     0.58985   3.321 0.002579 **
## log_Area       0.15724     0.03821   4.115 0.000327 ***
## log_Elev      -0.03081     0.12025  -0.256 0.799759
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2931 on 27 degrees of freedom
## Multiple R-squared:  0.7617, Adjusted R-squared:  0.744
## F-statistic: 43.15 on 2 and 27 DF,  p-value: 3.903e-09
```

```
avPlots(fit_both)
```

Added-Variable Plots




With $\log(\text{Native})$ transformation:

```
fit_both2 <- lm(Native_log ~ log_Area + log_Elev, data = galapagos_transformed2)
summary(fit_both2)
```

```
##
## Call:
## lm(formula = Native_log ~ log_Area + log_Elev, data = galapagos_transformed2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.2504 -0.2812  0.1135  0.4521  1.0356
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   2.83622    1.36815   2.073 0.047839 *
## log_Area       0.32724    0.08864   3.692 0.000994 ***
## log_Elev      -0.12364    0.27893  -0.443 0.661107
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.6799 on 27 degrees of freedom
## Multiple R-squared:  0.697, Adjusted R-squared:  0.6745
## F-statistic: 31.05 on 2 and 27 DF, p-value: 9.992e-08
```

```
avPlots(fit_both2)
```

Added-Variable Plots



ii. Fit a model that has (potentially transformed) Elev as the response and (potentially transformed) Area as the only explanatory variable. Add the residuals from this model to your data set with transformed variables.

This doesn't depend on the y transformation

```
fit_elev <- lm(log_Elev ~ log_Area, data = galapagos_transformed)
galapagos_transformed <- galapagos_transformed %>%
  mutate(
    resid_elev = residuals(fit_elev)
  )

galapagos_transformed2 <- galapagos_transformed2 %>%
  mutate(
    resid_elev = residuals(fit_elev)
  )
```

iii. Fit a model that has (potentially transformed) Native as the response and (potentially transformed) Area as the only explanatory variable. Add the residuals from this model to your data set with transformed variables.

With $\text{Native}^{0.25}$ transformation:

```
fit_native <- lm(Native_0.25 ~ log_Area, data = galapagos_transformed)
galapagos_transformed <- galapagos_transformed %>%
  mutate(
    resid_native = residuals(fit_native)
  )
```


With $\log(\text{Native})$ transformation:

```
fit_native2 <- lm(Native_log ~ log_Area, data = galapagos_transformed2)
galapagos_transformed2 <- galapagos_transformed2 %>%
  mutate(
    resid_native2 = residuals(fit_native2)
  )
```

iv. Make a plot that has the residuals from part ii on the horizontal axis and the residuals from part iii on the vertical axis. Compare this plot to the added variable plot for Elev from part i.

With $\text{Native}^{0.25}$ transformation:


```
ggplot(data = galapagos_transformed, mapping = aes(x = resid_elev, y = resid_native)) +
  geom_point() +
  theme_bw()
```



This plot matches the added variable plot above.

With $\log(\text{Native})$ transformation:

```
ggplot(data = galapagos_transformed2, mapping = aes(x = resid_elev, y = resid_native2)) +  
  geom_point() +  
  theme_bw()
```



This plot matches the added variable plot above.

v. Fit a linear model that has the residuals from part iii as the response and the residuals from part ii as the explanatory variable. Print out the model summary. Compare the coefficient estimate for the slope to the coefficient estimate for Elev from your model in part i.

With Native^{0.25} transformation:

```
av_fit <- lm(resid_native ~ resid_elev, data = galapagos_transformed)
summary(av_fit)
```

```
##
## Call:
## lm(formula = resid_native ~ resid_elev, data = galapagos_transformed)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.8167 -0.1842  0.0616  0.2340  0.4390
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -1.528e-17  5.255e-02   0.000    1.000
## resid_elev  -3.081e-02  1.181e-01  -0.261    0.796
##
```



```
## Residual standard error: 0.2878 on 28 degrees of freedom
## Multiple R-squared:  0.002425,    Adjusted R-squared:  -0.0332
## F-statistic: 0.06806 on 1 and 28 DF,  p-value: 0.7961
```

The coefficient estimate from this fit is the same as the coefficient estimate for elevation in the fit from part i.

With log(Native) transformation:

```
av_fit2 <- lm(resid_native2 ~ resid_elev, data = galapagos_transformed2)
summary(av_fit2)
```

```
##
## Call:
## lm(formula = resid_native2 ~ resid_elev, data = galapagos_transformed2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.2504 -0.2812  0.1135  0.4521  1.0356
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  4.278e-17  1.219e-01   0.000    1.000
## resid_elev  -1.236e-01  2.739e-01  -0.451    0.655
##
## Residual standard error: 0.6676 on 28 degrees of freedom
## Multiple R-squared:  0.007225,    Adjusted R-squared:  -0.02823
## F-statistic: 0.2038 on 1 and 28 DF,  p-value: 0.6552
```

The coefficient estimate from this fit is the same as the coefficient estimate for elevation in the fit from part i.