

Development of an algorithm that uses piecewise cubic functions to approximate a function.

For a cubic function, we will need 4 points $ax^3 + bx^2 + cx + d$

Picking the 2 middle points requires calculating the points between a and b that are $\frac{1}{3}$ and $\frac{2}{3}$ of the way between the endpoints. For a given interval defined by $[a, b]$, $\frac{1}{3}$ of the width of the interval is $\frac{b-a}{3}$.

$$\text{For } m_1, \text{ we have } m_1 = a + \frac{b-a}{3} = \frac{3a+b-a}{3} = \frac{2a+b}{3}$$

$$\text{For } m_2, \text{ we have } m_2 = a + 2 * \frac{b-a}{3} = \frac{3a+2b-2a}{3} = \frac{a+2b}{3}$$

$$\text{Next, we set } \int_a^b f(x) dx = A_1 f(a) + A_2 f(m_1) + A_3 f(m_2) + A_4 f(b)$$

Now, we calculate the results of integrating some simple functions ($1, x, x^2, x^3$) from a to b.

$$\int_a^b 1 dx = [x]_a^b = b - a = A_1 + A_2 + A_3 + A_4$$

$$\int_a^b x dx = \left[\frac{x^2}{2}\right]_a^b = \frac{b^2-a^2}{2} = aA_1 + \frac{2a+b}{3}A_2 + \frac{a+2b}{3}A_3 + bA_4$$

$$\int_a^b x^2 dx = \left[\frac{x^3}{3}\right]_a^b = \frac{b^3-a^3}{3} = a^2A_1 + \left(\frac{2a+b}{3}\right)^2A_2 + \left(\frac{a+2b}{3}\right)^2A_3 + b^2A_4$$

$$\int_a^b x^3 dx = \left[\frac{x^4}{4}\right]_a^b = \frac{b^4-a^4}{4} = a^3A_1 + \left(\frac{2a+b}{3}\right)^3A_2 + \left(\frac{a+2b}{3}\right)^3A_3 + b^3A_4$$

Converting these equations into an augmented matrix gives us:

$$\left| \begin{array}{cccc|c} 1 & 1 & 1 & 1 & b-a \\ a & \frac{2a+b}{3} & \frac{a+2b}{3} & b & \frac{b^2-a^2}{2} \\ a^2 & \left(\frac{2a+b}{3}\right)^2 & \left(\frac{a+2b}{3}\right)^2 & b^2 & \frac{b^3-a^3}{3} \\ a^3 & \left(\frac{2a+b}{3}\right)^3 & \left(\frac{a+2b}{3}\right)^3 & b^3 & \frac{b^4-a^4}{4} \end{array} \right|$$

First, we subtract $a \cdot R_1$ from R_2 , $a^2 \cdot R_1$ from R_3 , and $a^3 \cdot R_1$ from R_4 to get:

$$\left| \begin{array}{cccc|c} 1 & 1 & 1 & 1 & b-a \\ 0 & \frac{b-a}{3} & \frac{2b-2a}{3} & b-a & \frac{(b-a)^2}{2} \\ 0 & \frac{(b-a)(5a+b)}{9} & \frac{(a+b)^2-9a^2}{9} & b^2-a^2 & \frac{(a-b)^3(2a+b)}{3} \\ 0 & \frac{(2a+b)^3-27a^3}{27} & \frac{(a+2b)^3-27a^3}{27} & b^3-a^3 & \frac{3a^4-4a^3b+b^4}{4} \end{array} \right|$$

Next, we simplify R2 by dividing it by $\frac{b-a}{3}$

1	1	1	1	$\frac{b-a}{2}$
0	1	2	3	$\frac{3b-3a}{2}$
0	$\frac{(b-a)(5a+b)}{9}$	$\frac{(a+b)^2-9a^2}{9}$	b^2-a^2	$\frac{(a-b)^3(2a+b)}{3}$
0	$\frac{(2a+b)^3-27a^3}{27}$	$\frac{(a+2b)^3-27a^3}{27}$	b^3-a^3	$\frac{3a^4-4a^3b+b^4}{4}$

Next we subtract R2 from R1, $\frac{(b-a)(5a+b)}{9}$ *R2 from R3, and $\frac{(2a+b)^3-27a^3}{27}$ *R2 from R4

1	0	-1	-2	$\frac{a-b}{2}$
0	1	2	3	$\frac{3b-3a}{2}$
0	0	$\frac{2(a-b)^2}{9}$	$\frac{2(a-b)^2}{3}$	$\frac{(b-1)^3}{6}$
0	0	$\frac{2(a-b)^2(2a+b)}{9}$	$\frac{2(a-b)^2(5a+4b)}{9}$	$\frac{-(a-b)^3(11a+7b)}{36}$

Next, we simplify by multiplying R3 by $\frac{9}{2(a-b)^2}$

1	0	-1	-2	$\frac{a-b}{2}$
0	1	2	3	$\frac{3b-3a}{2}$
0	0	1	3	$\frac{(3b-3a)}{4}$
0	0	$\frac{2(a-b)^2(2a+b)}{9}$	$\frac{2(a-b)^2(5a+4b)}{9}$	$\frac{-(a-b)^3(11a+7b)}{36}$

Next, we add R3 to R1, and subtract 2*R3 from R2 and $\frac{2(a-b)^2(2a+b)}{9}$ *R3 from R4

1	0	0	1	$\frac{b-a}{4}$
0	1	0	-3	0
0	0	1	3	$\frac{(3b-3a)}{4}$
0	0	0	$\frac{2(b-a)^3}{9}$	$\frac{(a-b)^4}{36}$

Next, we simplify R4 by multiplying it by $\frac{9}{2(b-a)^3}$

1	0	0	1	$\frac{b-a}{4}$
0	1	0	-3	0
0	0	1	3	$\frac{3(b-a)}{4}$
0	0	0	1	$\frac{b-a}{8}$

Finally, we subtract R4 from R1, add 3*R4 to R2 and subtract 3*R4 from R3

$$\left| \begin{array}{cccc|c} 1 & 0 & 0 & 0 & \frac{b-a}{8} \\ 0 & 1 & 0 & 0 & \frac{3(b-a)}{8} \\ 0 & 0 & 1 & 0 & \frac{3(b-a)}{8} \\ 0 & 0 & 0 & 1 & \frac{b-a}{8} \end{array} \right|$$

$$\begin{aligned} \text{So, } \int_a^b f(x) dx &= \frac{b-a}{8} f(a) + \frac{3(b-a)}{8} f(m_1) + \frac{3(b-a)}{8} f(m_2) + \frac{b-a}{8} f(b) \\ &= \frac{b-a}{8} \left[f(a) + 3f\left(\frac{2a+b}{3}\right) + 3f\left(\frac{a+2b}{3}\right) + f(b) \right] \end{aligned}$$

Example 1

Exact

$$\int_0^{11} x dx = \left[\frac{x^2}{2} \right]_0^{11} = \left[\frac{121}{2} - 0 \right] = \frac{121}{2} = 60.5$$

Approximation

$$\begin{aligned} \int_0^{11} x dx &= \frac{11-0}{8} \left[f(0) + 3f\left(\frac{2(0)+11}{3}\right) + 3f\left(\frac{0+2(11)}{3}\right) + f(11) \right] \\ &= \frac{11}{8} \left[0 + 3f\left(\frac{11}{3}\right) + 3f\left(\frac{22}{3}\right) + 11 \right] \\ &= \frac{11}{8} [11 + 11 + 22] = 11/8(44) = \frac{121}{2} = 60.5 \end{aligned}$$

Error

$$\text{Exact} - \text{Approximation} = 60.5 - 60.5 = 0$$

Example 2

Exact

$$\int_{12}^{13} x^{-2} dx = \left[-\frac{1}{x} \right]_{12}^{13} = -\frac{1}{13} - \left(-\frac{1}{12} \right) = -\frac{12}{156} + \frac{13}{156} = \frac{1}{156} \approx .0064102564102564$$

Approximation

$$\begin{aligned} \int_{12}^{13} x^{-2} dx &= \frac{13-12}{8} \left[f(12) + 3f\left(\frac{2(12)+13}{3}\right) + 3f\left(\frac{12+2(13)}{3}\right) + f(13) \right] \\ &= \frac{1}{8} \left[\frac{1}{144} + 3f\left(\frac{37}{3}\right) + 3f\left(\frac{38}{3}\right) + \frac{1}{169} \right] = \frac{1}{8} \left[\frac{313}{24336} + \frac{27}{1369} + \frac{27}{1444} \right] \\ &= \frac{1}{8} \left(\frac{616773301}{12027070224} \right) = \frac{616773301}{96216561792} \approx .0064102612846771 \end{aligned}$$

Error

$$\text{Exact - Approximation} = \frac{1}{156} - \frac{616773301}{96216561792} = -\frac{469}{96216561792} \approx -4.87442 \cdot 10^{-9}$$

Example 3

Exact

$$\int_3^6 3x^2 + \frac{x}{7} dx = \left[x^3 + \frac{x^2}{14} \right]_3^6 = \left(216 + \frac{18}{7} \right) - \left(27 + \frac{9}{14} \right) = \frac{1530}{7} - \frac{387}{14} = \frac{2673}{14} \\ \approx 190.9285714$$

Approximation

$$\int_3^6 3x^2 + \frac{x}{7} dx = \frac{6-3}{8} \left[f(3) + 3f\left(\frac{2(3)+6}{3}\right) + 3f\left(\frac{3+2(6)}{3}\right) + f(6) \right] \\ = \frac{3}{8} \left[\frac{192}{7} + 3f(4) + 3f(5) + \frac{762}{7} \right] = \frac{3}{8} \left[\frac{954}{7} + \frac{1020}{7} + \frac{1590}{7} \right] \\ = \frac{3}{8} \left(\frac{3564}{7} \right) = \frac{2673}{14} \approx 190.9285714$$

Error

$$\text{Exact - Approximation} = \frac{2673}{14} - \frac{2673}{14} = 0$$

Example 4

Exact

$$\int_{-2}^5 x^3 - 5 dx = \left[\frac{x^4}{4} - 5x \right]_{-2}^5 = \left(\frac{5^4}{4} - 25 \right) - \left(\frac{-2^4}{4} + 10 \right) = \frac{525}{4} - 14 = \frac{469}{4} = 117.25$$

Approximation

$$\int_{-2}^5 x^3 - 5 dx = \frac{5-(-2)}{8} \left[f(-2) + 3f\left(\frac{2(-2)+5}{3}\right) + 3f\left(\frac{-2+2(5)}{3}\right) + f(5) \right] \\ = \frac{7}{8} \left[f(-2) + 3f\left(\frac{1}{3}\right) + 3f\left(\frac{8}{3}\right) + f(5) \right] \\ = \frac{7}{8} \left[-13 + 3\left(\frac{-134}{27}\right) + 3\left(\frac{377}{27}\right) + 120 \right] \\ = \frac{7}{8} [107 + 27] = \frac{938}{8} = \frac{469}{4} = 117.25$$

Error

$$\text{Exact - Approximation} = 117.25 - 117.25 = 0$$

However, this algorithm is only exact for powers up to 3. It is not exact for powers higher than 3.

Example 5

Exact

$$\int_1^3 x^4 dx = \left[\frac{x^5}{5} \right]_1^3 = \left[\frac{243}{5} - \frac{1}{5} \right] = \frac{242}{5} = 48.4$$

Approximation

$$\begin{aligned} \int_1^3 x^4 dx &= \frac{3-(1)}{8} [f(1) + 3f(\frac{2(1)+3}{3}) + 3f(\frac{1+2(3)}{3}) + f(3)] \\ &= \frac{1}{4} [1 + 3f(\frac{5}{3}) + 3f(\frac{7}{3}) + 81] \\ &= \frac{1}{4} [82 + \frac{5^4}{3^3} + \frac{7^4}{3^3}] \\ &= \frac{1}{4} [82 + \frac{3026}{27}] \\ &= \frac{1310}{27} \approx 48.\overline{518} \end{aligned}$$

Error

$$\text{Exact - Approximation} = \frac{242}{5} - \frac{1310}{27} = \frac{-16}{135} \approx -0.118\overline{518}$$