

Shell-model fits for Sn isotopes

M. Honma,^{*1} T. Otsuka,^{*2,*3} T. Mizusaki,^{*4} and M. Hjorth-Jensen^{*5}

[NUCLEAR STRUCTURE, shell model]

The Sn isotopes provide us with an ideal test for conventional shell-model calculations on the basis of the microscopic effective interaction theory and also the knowledge of nuclear forces. Because of the $Z=50$ semi-magic structure, the deformation is not significant, and the shell-model description within a limited one-major shell is expected to be reasonable. In fact, shell-model calculations using an effective interaction derived microscopically from a realistic nucleon-nucleon potential have been successfully applied for the analysis of new experimental data and also the prediction for unobserved properties¹⁾.

However, such a microscopic approach is not necessarily successful in the middle of the neutron shell as shown in Ref.²⁾ for example, where the excitation energy of the 2_1^+ state is predicted to be too high by 0.4 MeV for ^{118}Sn . On the other hand, it has been shown that the microscopic effective interaction can be improved for a practical use by fitting to the experimental data. Therefore, it is interesting to investigate how accurate the structure of Sn isotopes can be described by using such a fitted effective interaction. We report the results of such an attempt.

The adopted model space consists of five single-particle orbits $1d_{5/2}$, $0g_{7/2}$, $0h_{11/2}$, $2s_{1/2}$ and $1d_{3/2}$ in which the Sn isotopes are described by valence neutrons, and the $T = 1$ part of the two-body interactions is relevant. We start with an effective interaction obtained in a microscopic way as described in Ref.³⁾ using the $N^3\text{LO}$ interaction⁴⁾. A series of fitting calculations was carried out by taking 313 experimental energy data including 29 binding energies. We utilized the Linear-Combination (LC) method as in our previous approach⁵⁾. We found that only a small modification was necessary to improve the microscopic interaction. In fact a reasonable convergence was obtained by varying only 22 LC's of 165 Hamiltonian parameters (5 single-particle energies and 160 two-body matrix elements). In the final fit, the rms deviation of 147 keV was obtained.

As an example of the results, the excitation energies of the yrast states of even- N isotopes are shown in Fig.1. One can find a reasonable agreement between the experimental data and the shell-model results throughout the isotope chain. The quality of the

fit is similar for odd- N isotopes. The predicted ground-state spin of ^{101}Sn is $7/2^+$ rather than $5/2^+$, in good agreement with recent experimental observation.⁶⁾ The fit of the binding energy is basically successful for heavier isotopes, but the shell-model result begins to show underbinding as approaching ^{100}Sn . This result suggests the development of collectivity which can not be described by the present model space. Such a picture is consistent with the large $B(E2)$ values below $N = 64$ observed in recent extensive measurements⁷⁾.

This fitted interaction can be a good starting point for extending the model space so as to describe the ^{100}Sn core excitations and also for the study of more neutron-rich isotopes.




Fig. 1. Comparison of excitation energies of even-spin yrast states between the experimental data (symbols) and the shell-model results (lines). Experimental data are taken from Ref.⁸⁾ Open symbols indicate that the spin assignment is uncertain. The shell-model results are obtained by using the efficient shell-model code MSHELL⁹⁾.

References

- 1) A. Banu *et al.*: Phys. Rev. C **72**, 061305(R) (2005).
- 2) P. Guazzoni *et al.*: Phys. Rev. C **78**, 064608 (2008).
- 3) M. Hjorth-Jensen *et al.*: Phys. Rep. **261**, 125 (1995).
- 4) D. R. Entem *et al.*: Phys. Rev. C **68**, 041001(R) (2003).
- 5) M. Honma *et al.*: Phys. Rev. C **80**, 064323 (2009).
- 6) I. G. Darby *et al.*: Phys. Rev. Lett. **105**, 162502 (2010).
- 7) A. Ekström *et al.*: Phys. Rev. Lett. **101**, 012502 (2008).
- 8) Data extracted using the NNDC WorldWideWeb site from the ENSDF database.
- 9) T. Mizusaki: RIKEN Accel. Prog. Rep. **33**, 14 (2000).

^{*1} Center for Mathematical Sciences, University of Aizu
^{*2} Department of Physics, University of Tokyo
^{*3} Center for Nuclear Studies, University of Tokyo
^{*4} Institute of Natural Sciences, Senshu University
^{*5} Department of Physics and Center of Mathematics for Applications, University of Oslo