Modelling of Orthophosphate and Zinc Inputs to Rivers from Diffuse and Point Sources using a Bayesian Load Apportionment Model with Time-Varying Parameters

Maximillian McCourt

University of Edinburgh
June 2026

Executive Summary

Effective management of river water quality in England depends on whether pollution originates from point sources, such as sewage treatment work (STW) effluents, or diffuse sources, such as agricultural runoff, as each requires fundamentally different regulatory responses. Under the Water Framework Directive, many English river bodies continue to fail Good Ecological Status, yet the balance between point and diffuse sources has shifted considerably since 2000 as point-source regulation has tightened. Understanding where this balance now stands, and how it has evolved, is essential for targeting future investment effectively.

This report investigates point and diffuse-source contributions to orthophosphate and dissolved zinc concentrations at two contrasting sites: the Thames at Teddington Weir, a large urbanised catchment with a well-documented history of STW regulation, and the Wyre at St Michaels, a smaller predominantly agricultural catchment, over the 2000–2024 period. The Load Apportionment Model was applied in a Bayesian framework to quantify uncertainty in source apportionment, and extended to allow point-source contributions to change over time rather than assuming them fixed, better reflecting the evolving regulatory and catchment conditions of a 25-year monitoring record.

For zinc, persistent point-source dominance was found at the Thames and diffuse-source dominance at the Wyre throughout the period. However, the monitoring data for zinc do not show the clear contrast between point and diffuse behaviour that the framework relies upon, and uncertainty in these apportionments is too wide to reliably guide compliance decisions at either site. Further methodological development would be needed before zinc source apportionment can inform targeted Environment Agency (EA) intervention.

The time-varying framework consistently outperformed the traditional fixed-parameter approach at sites where catchment conditions changed most rapidly, and is recommended as the preferred tool for long-term source apportionment analysis at regulated catchments. Extending this approach to all 135 Harmonised Monitoring Scheme sites across England would allow the EA to systematically identify catchments where point-source infrastructure investment would yield the greatest water quality improvement, versus those where diffuse source management has become the binding constraint on achieving Good Ecological Status.

Contents

1 Introduction **4**

2 Data Overview and Pre-processing **4**
2.1 Water Quality Concentration Data . . . 4
2.2 Flow Data . . . 4
2.3 Censored Observations . . . 4

3 Exploratory Data Analysis **5**
3.1 Generalised Additive Model (GAM) . . . 5
3.1.1 Model Selection and Results . . . 5

4 Bayesian LAM **7**
4.1 Evaluation of Model . . . 7
4.2 Results . . . 8

5 Bayesian LAM with time-varying parameters (TV-LAM) **9**
5.1 Evaluation of Model . . . 10
5.2 Results . . . 10

6 Model Comparison **12**
6.1 Residual Temporal Bias . . . 12
6.2 In-Sample Goodness of Fit . . . 12

7 Discussion **13**
7.1 Main Findings . . . 13
7.2 Limitations . . . 13
7.3 Future Directions . . . 14

Appendix A GAM Diagnostics: Residual vs Fitted graphs **16**

Appendix B Bayesian LAM Model Diagnostics **18**

Appendix C Bayesian LAM Posterior Parameter Estimates **34**

Appendix D Bayesian LAM Posterior Median Point Source contribution to total load **34**

Appendix E Bayesian TV-LAM Fitting Diagnostics **35**

Appendix F Bayesian TV-LAM Scalar Parameter Estimates and Convergence Diagnostics **38**

Appendix G Observation-level CRPS Comparison **39**

List of Figures

1 Orthophosphate concentrations, 2000–2024 . . . 5
2 Zinc concentrations, 2000–2024 . . . 5
3 Log concentration vs log flow, all site-determinand combinations . . . 6
4 Residuals vs fitted: Thames orthophosphate interaction GAM . . . 6
5 Bayesian LAM residuals vs flow: Thames orthophosphate, 2000–2004 . . . 7
6 Bayesian LAM residuals vs time: Thames orthophosphate, 2000–2004 . . . 8
7 Bayesian LAM QQ plot: Thames orthophosphate, 2000–2004 . . . 8
8 Bayesian LAM posterior predictive check: Thames orthophosphate, 2000–2004 . . . 8
9 Windowed point-source contribution (%): orthophosphate . . . 8
10 Windowed point-source contribution (%): zinc . . . 9
11 TV-LAM traceplots: Thames orthophosphate . . . 10
12 TV-LAM residuals vs time: Thames orthophosphate . . . 10
13 TV-LAM posterior predictive check: Thames orthophosphate . . . 10
14 TV-LAM posterior median A_t: orthophosphate . . . 11
15 TV-LAM posterior median A_t: zinc . . . 11
16 TV-LAM point-source contribution (%): orthophosphate . . . 11
17 TV-LAM point-source contribution (%): zinc . . . 11
18 Observation-level CRPS comparison: Thames orthophosphate . . . 13
19 Appendix: Residuals vs fitted: orthophosphate GAM models . . . 16
20 Appendix: Residuals vs fitted: zinc GAM models . . . 17
21 Appendix: Bayesian LAM residuals vs flow: Thames orthophosphate, 2005–2024 . . . 18
22 Appendix: Bayesian LAM residuals vs flow: Wyre orthophosphate, 2000–2024 . . . 19
23 Appendix: Bayesian LAM residuals vs flow: Thames zinc, 2000–2024 . . . 20
24 Appendix: Bayesian LAM residuals vs flow: Wyre zinc, 2005–2024 . . . 21
25 Appendix: Bayesian LAM residuals vs time: Thames orthophosphate, 2005–2024 . . . 22
26 Appendix: Bayesian LAM residuals vs time: Wyre orthophosphate, 2000–2024 . . . 23
27 Appendix: Bayesian LAM residuals vs time: Thames zinc, 2000–2024 . . . 24
28 Appendix: Bayesian LAM residuals vs time: Wyre zinc, 2005–2024 . . . 25
29 Appendix: QQ plots: Thames orthophosphate LAM residuals, 2005—24 . . . 26
30 Appendix: QQ plots: Wyre orthophosphate LAM residuals, 2000–2024 . . . 27
31 Appendix: QQ plots: Thames zinc LAM residuals, 2000–2024 . . . 28
32 Appendix: QQ plots: Wyre zinc LAM residuals, 2005–2024 . . . 29
33 Appendix: Posterior predictive checks: Thames orthophosphate LAM, 2005–2024 . . . 30
34 Appendix: Posterior predictive checks: Wyre orthophosphate LAM, 2000–2024 . . . 31
35 Appendix: Posterior predictive checks: Thames zinc LAM, 2000–2024 . . . 32
36 Appendix: Posterior predictive checks: Wyre zinc LAM, 2005–2024 . . . 33
37 Appendix: TV-LAM traceplots: Wyre orthophosphate, Thames zinc, Wyre zinc . . . 35
38 Appendix: TV-LAM residuals vs time: Wyre orthophosphate, Thames zinc, Wyre zinc . . . 36
39 Appendix: TV-LAM posterior predictive checks: Wyre orthophosphate, Thames zinc, Wyre zinc . . . 37
40 Appendix: Observation-level CRPS comparison: remaining site-determinand combinations . . . 39

List of Tables

1 Data availability and censoring summary . . . 4
2 REML scores: additive vs interaction GAMs . . . 6
3 GAM smooth term complexity and significance . . . 7
4 Prior specifications for Bayesian LAM and TV-LAM . . . 7
5 Mean CRPS: Windowed baseline LAM vs TV-LAM . . . 12
6 Appendix: Bayesian LAM posterior parameter estimates: orthophosphate . . . 34
7 Appendix: Bayesian LAM posterior parameter estimates: zinc . . . 34
8 Appendix: Posterior median point-source contribution (%), windowed LAM . . . 34
9 Appendix: TV-LAM scalar parameter estimates: orthophosphate . . . 38
10 Appendix: TV-LAM scalar parameter estimates: zinc . . . 38

1 Introduction

E ective management of river water quality in England requires the ability to distinguish between point-source pollutants, such as sewage treatment work (STW) e uents and industrial discharges, and di use-source pollutants, such as agricultural runo and road drainage. Under the EU Water Framework Directive (WFD; 2000/60/EC), river bodies across England are required to achieve Good Ecological Status (GES), yet many continue to fail these standards [10, 15]. Identifying the dominant pollution source is therefore essential for targeting mitigation e orts e ectively.

Excess orthophosphate promotes harmful algal growth and eutrophication, causing dissolved oxygen uctuations that disrupt aquatic ecosystems [6]. Elevated concentrations of dissolved metals such as zinc are directly toxic to aquatic organisms and pose risks to human health [9]. As point-source contributions have declined following regulatory intervention, notably mandatory phosphorus stripping at larger STWs under the Urban Wastewater Treatment Directive, di use sources are increasingly the binding constraint on achieving Good Ecological Status. Quantifying the relative contributions of point and di use sources over time has therefore become a central water management challenge.

The Load Apportionment Model (LAM), introduced by [2], exploits the contrasting hydrological signatures of these two source types, expressing total pollutant concentration as the sum of two power-law functions of river ow. Requiring only routine concentration and ow monitoring data, the LAM o ers a parsimonious and interpretable framework for source apportionment. However, its assumption of xed parameters is inappropriate over long time series where catchment conditions evolve, a limitation motivating this study. We therefore propose a Bayesian implementation of the LAM to formally quantify uncertainty in source apportionment estimates, and a time-varying extension (TV-LAM) to accommodate structural non-stationarity in the ow-concentration relationship.

This report applies both frameworks to orthophosphate and zinc at contrasting sites on the rivers Thames and Wyre, comparing their ability to characterise point and di use source contributions over a 25-year period and evaluating whether accounting for temporal variation meaningfully affects source apportionment conclusions.

2 Data Overview and Pre-processing

2.1 Water Quality Concentration Data

Spot-sample water quality data were obtained from the Harmonised Monitoring Scheme (HMS) network, which provides a long-term record of nutrient and metal concentrations at 135 sites on England's principal rivers [13]. Although GB-wide o cial reporting ceased in 2013, the Environment Agency (EA) has continued sampling at the same sites under its own monitoring programme, with all samples analysed by EA laboratories and their predecessors, ensuring consistency across the full 2000 2024 study period [14].

We focus our analysis on two determinands: orthophosphate ($mg\,l^{-1}$) and dissolved zinc ($\mu g\,l^{-1}$), chosen to represent contrasting nutrient and metal pollutant types with di ering point/di use source characteristics. These concentrations are investigated at two contrasting sites: Thames at Teddington Weir, a large, urbanised, heavily regulated catchment with a well-documented history of STW phosphorus stripping; and Wyre at St Michaels, a smaller, predominantly agricultural catchment expected to exhibit greater di use-source in uence. This contrast provides a meaningful test of the LAM framework across di erent source regimes. Both sites were sampled at least monthly over the 2000 2024 study period, with known gaps during the 2001 foot-and-mouth outbreak and 2020 2021 COVID-19 lockdowns [14].

Observations falling below implausibly low thresholds, less than $0{:}005\,mg\,l^{-1}$ for orthophosphate and less than $0{:}5\,\mu g\,l^{-1}$ for dissolved zinc, were removed as likely analytical errors. This resulted in the removal of one orthophosphate observation; no zinc records were a ected.

2.2 Flow Data

Daily mean river discharge ($m^3 s^{-1}$) was obtained from the Environment Agency Hydrological Data Explorer API and matched to each water quality spot sample by date [1]. Each concentration observation was paired to the daily mean ow on the corresponding sampling date. Where multiple samples existed on the same date, these raw concentrations were averaged prior to log transformation.

River ow data included Estimated (17.8%), Suspect (9.0%), and Unchecked (0.7%) quality- agged readings alongside Good-quality observations (72.6%). Estimated readings re ect standard EA in lling procedures and are not indicative of measurement error. Suspect readings showed no systematic association with high- ow gauge exceedance, and therefore all quality categories were retained.

2.3 Censored Observations

Table 1 summarises the proportion of censored observations at each site-determinand combination.

Table 1: Data availability summary for key sites and determinands. Censoring indicates observations recorded below the limit of detection.

Determinand	Proportion of censored data	
	Thames at Teddington Weir	Wyre at St Michaels
Orthophosphate	0.0083	0.0034
Dissolved Zinc	0.1962	0.2222

Zinc exhibits substantially more censored observations (around 20% at both sites) than orthophosphate (< 1%), for which the true concentration is known only to lie below the recorded limit of detection (LoD). Censored zinc observations are concentrated in the earlier part of the study period, becoming increasingly sparse after approximately 2015, consistent with improvements in EA analytical detection capability over time, particularly from 2014 onwards

[9]. Naively substituting these values (for example, with half the LoD) or discarding them entirely introduces informative selection bias, as below-detection observations are systematically associated with lower true concentrations. Instead, each censored observation is encoded as a left-censored interval $(-1, \log_{10}(d_i)]$, where d_i is the recorded LoD, allowing models used in this report to integrate over this uncertainty when estimating parameters rather than treating below-detection values as known.

3 Exploratory Data Analysis

To model orthophosphate and zinc concentrations effectively, we first examine their temporal trends and relationship with river flow. Figures 1 and 2 show the concentrations of orthophosphate and zinc over time at both sites.

Figure 1: Log-scaled orthophosphate (mg/l) measurements over years 2000–2024 at primary sites Thames at Teddington Weir and Wyre at St Michaels.

Figure 2: Log-scaled zinc (µg/l) measurements over years 2000–2024 at primary sites Thames at Teddington Weir and Wyre at St Michaels.

Orthophosphate declines clearly over the 25-year period at both sites, with the Thames exhibiting a particularly sharp reduction between 2005 and 2010 before stabilising. Zinc trends are more site-specific: the Thames shows a general decline, while the Wyre decreases until around 2012 before rising steadily to 2024. These long-run trends indicate that the LAM's assumption of fixed parameters is inappropriate over the full period, motivating the use of shorter time windows for the baseline model (Section 4), and a time-varying extension thereafter (Section 5).

Figure 3 shows the relationship between log river flow and log concentration for both determinands at both sites.

Orthophosphate exhibits a clear negative relationship with flow at both sites, consistent with dilution of a predominantly point-source signal, with diminishing returns at higher flows. Zinc, by contrast, shows a more complex and site-variable relationship: concentrations initially decrease at low flows before rising again at high flows. This upturn is consistent with zinc having a stronger diffuse-source component, whereby high-flow events that dilute point-source inputs simultaneously mobilise zinc from agricultural land and urban runoff.

The contrasting flow-concentration relationships between the two determinands motivate a modelling framework capable of separately quantifying point-source and diffuse-source contributions.

3.1 Generalised Additive Model (GAM)

Generalised additive models (GAMs) are generalised linear models that allow nonlinear functions of covariates to be incorporated in regression equations, making them well-suited for capturing nonlinear temporal trends [19]. While GAMs cannot separately quantify point-source and diffuse-source contributions, they serve here as a diagnostic tool to assess whether the flow-concentration relationship changes over time, a finding that directly motivates the time-varying extension of the LAM developed in Section 5.

We fit both an additive model,

$$\log_{10}(C_i) = \alpha + f_1(m_i) + f_2(t_i) + f_3(\log Q_i) + \varepsilon, \quad \varepsilon \sim N(0, \sigma^2), \tag{1}$$

and an interaction model,


$$\begin{aligned} \log_{10}(C_i) = \alpha &+ f_1(m_i) + f_2(t_i) \\ &+ f_3(\log Q_i) + f_4(t_i, \log Q_i) + \varepsilon, \\ &\varepsilon \sim N(0, \sigma^2), \end{aligned} \tag{2}$$

where Q_i is river discharge ($m^3 s^{-1}$), t_i is the year-date in decimal form, and m_i is a factor variable representing months of the year. Each smooth f_j is represented as a penalised regression spline, with smoothing parameters estimated by restricted maximum likelihood (REML) [19].

Models were constructed using the mgcv [19] library in R [17], with censored observations handled via the cnorm() family as described in Section 2.3. Basis dimension sufficiency was confirmed for all site-determinand combinations using gam.check().

3.1.1 Model Selection and Results

Both models were fitted for each site-determinand combination and compared via REML score (Table 2).



(a) Thames orthophosphate

(b) Wyre orthophosphate

(c) Thames zinc

(d) Wyre zinc

Figure 3: Log10-scaled concentration against log10-scaled river ow (m^3=s) for orthophosphate (mg=l) and zinc (g=l) at Thames at Teddington Weir and Wyre at St Michaels.

Table 2: REML scores for additive and interaction GAM models across all site-determinand combinations. More negative values indicate better t.

Site	Det.	Additive	Interaction
Thames	Phosphate	74:3	79:4
Thames	Zinc	42:6	44:4
Wyre	Phosphate	179:4	320:7
Wyre	Zinc	3:7	7:7

The interaction model achieved a lower REML score in all cases. The interaction term $f_4(t_i; \log Q_i)$ was statistically signi cant for orthophosphate at both sites (Thames: $p < 0:001$, Wyre: $p < 0:001$), with REML improvements of 5.1 and 141.3 respectively, indicating that the relationship between river ow and orthophosphate concentration changes meaningfully over the 25-year period. For zinc, the interaction term was not signi cant at either site (Thames: $p = 0:189$, Wyre: $p = 0:163$), with smaller REML improvements of 1.8 and 4.0. The interaction model was therefore selected for orthophosphate and the additive model for zinc.

Residual diagnostics for the orthophosphate interaction model at the Thames are shown in Figure 4.

Figure 4: Residuals vs tted values for the interaction GAM t-ted to orthophosphate at the Thames, with smoothed blue LOESS trend line.

The residuals vs tted plot shows no systematic trend or heteroskedasticity, indicating the model assumptions are adequately satis ed. Results for other site-determinand model cases were broadly similar (see Appendix A).

Table 3 summarises the estimated degrees of freedom and signi cance of each smooth term across all nal models.

All terms were statistically signi cant ($p < 0:05$) except the seasonal term $f_1(m_i)$ for zinc at the Wyre (edf = 0:00, $p = 0:483$), suggesting no meaningful seasonal pattern in zinc at this site. The temporal trend $f_2(t_i)$ was highly signi cant and nonlinear for both orthophosphate models (edf 10:50 at the Thames and 10:25 at the Wyre), consistent with the long-term concentration decline visible in Figure 1. The interaction term $f_4(t_i; \log Q_i)$ showed notably higher complexity at the Wyre (edf = 12:65) than the Thames (edf = 2:30), suggesting the ow-concentration relationship has changed more substantially over time at the rural site.

Table 3: Estimated degrees of freedom (edf) and p-values for each smooth term across all nal GAM models (additive and interaction of each site-determinand case).

		Orthophosphate Interaction model		Zinc Additive model	
Term	Site	edf	p	edf	p
$f_1(m_i)$	Thames	6.31	<0.001	4.52	<0.001
	Wyre	2.70	<0.001	0.00	0.483
$f_2(t_i)$	Thames	10.50	<0.001	1.00	<0.001
	Wyre	10.25	<0.001	3.84	<0.001
$f_3(\log Q_i)$	Thames	2.45	<0.001	3.64	<0.001
	Wyre	3.79	<0.001	4.02	<0.001
$f_4(t_i; \log Q_i)$	Thames	2.30	<0.001		
	Wyre	12.65	<0.001		

These ndings establish that signi cant temporal non-stationarity is present in all site-determinand combinations, and the ow-concentration relationship for orthophosphate changes structurally over time. This motivates both the use of short time-windows in the baseline LAM (Section 4) and the time-varying extension developed in Section 5.

4 Bayesian LAM

The Load Apportionment Model (LAM), introduced by [2], provides the mechanistic framework to separately quantify point and di use-source contributions, expressing total pollutant concentration as the sum of two power-law functions of river ow.

Model. The LAM is expressed in load form, where the total pollutant load F_i ($mg\,s^{-1}$) at observation i is modelled as

$$F_i = A\ Q_i^B + C^0\ Q_i^D; \qquad (3)$$

where Q_i is river discharge ($m^3 s^{-1}$), A and B are the point-source parameters, and C^0 and D are the di use-source parameters. Since concentration C_i ($mg\,m^{-3}$) equals load divided by volumetric ow, this can be expressed as

$$C_i = A\ Q_i^{B-1} + C^0\ Q_i^{D-1} + \varepsilon_i; \quad \varepsilon_i \sim N(0; \sigma_i^2): \qquad (4)$$

where the rst term $A\ Q_i^{B-1}$ represents point-source contributions and the second term $C^0\ Q_i^{D-1}$ represents di use-source contributions.

As the LAM assumes xed parameters, it cannot accommodate long-term trends in catchment conditions. Models were therefore tted to ve-year windows: 2000 2004, 2005 2009, 2010 2014, 2015 2019, and 2020 2024, with the exception of 2000 2004 of Wyre zinc, for which monitoring commenced in 2003.

Left-censored observations are incorporated by treating the likelihood contribution as $P(y_i < d_i)$, integrating over $(-1; \log_{10}(d_i))$, rather than substituting or excluding these values.

Priors. Weakly informative priors were used, with means informed by [2] for phosphorus and applied identically to zinc in the absence of analogous prior information (see Table 4).

Table 4: Prior speci cations for models. Sign constraints are imposed as hard bounds. Parameters A and C are assigned half-normal priors as their scale depends on site-speci c load magnitudes.

Parameter	Constraint	Prior $N(\cdot;\ \cdot)$
A	> 0	$N(0;\ 10)$
B	[0; 1]	$N(0{:}5;\ 5)$
C^0	> 0	$N(0;\ 10)$
D	> 1	$N(1{:}5;\ 5)$

4.1 Evaluation of Model

All 19 models achieved satisfactory convergence when constructed in brms [5] in R [17], with $\hat{R} < 1{:}01$ and e ective sample size exceeding 400 for all parameters across all site-determinand-window combinations [11]. Trace plots for these models con rmed adequate chain mixing throughout.

Residual diagnostics (Figures 5, 6 and 7) and posterior predictive checks (Figure 8) were performed on all models, with results shown in this paper for the Thames-Orthophosphate combination from 2000 2004 as an example of goodness-of- t of our models. Results for all other baseline models can be seen in Appendix B.

Figure 5: LAM Residual vs Flow of the Bayesian LAM on Thames-Orthophosphate data in 2000 2004, with smoothed blue LOESS trend line.

The LOESS smoothing line of the residual vs ow plot in Figure 5 is relatively at around the zero-line, indicating the model captures the relationship between orthophosphate concentration and river ow in the Thames in 2000 2004 reasonably well. This was re ected in most site-determinand-period model combinations (see Appendix B.1), which generally exhibit similar at LOESS behaviour.

The LOESS smoothing line of the residual vs time plot in Figure 6 however shows a systematic downward trend over time, suggesting residual temporal non-stationary is present. Even after splitting the data into shorter 5-year periods, this model systematically overpredicts later in the period and underpredicts earlier. Systematic trends such as these were shown in several models trained, indicating that a clear limitation in our LAM framework is its inability to handle temporal non-stationarity. This systematic residual trend

was present across most site-determinand combinations, as shown in Appendix B.2.

Figure 6: Residual vs Time of the Bayesian LAM on Thames-Orthophosphate data in 2000 2004, with smoothed blue LOESS trend line.

Figure 7: LAM QQ-plot of the Bayesian LAM on Thames-Orthophosphate data in 2000 2004. Dashed line = theoretical normal quantiles.

The residuals mainly follow a straight line in the QQ plot (Figure 7), suggesting that the model residuals mainly follow a normal distribution. There is however some deviation in the lower tail, suggesting a slight skewness in the lower quantiles of our concentration data. Minor lower-tail deviations were present across most models (Appendix B.3), but the normality assumption is broadly satis ed.

The posterior predictive density plot of simulated and actual $\log_{10}$ concentration here shows that the model's simulated densities envelope the observed curve reasonably well, thus indicating our model ts the data reasonably well. Posterior predictive checks for all remaining site-determinand-window combinations are shown in Appendix B.4, and demonstrate reasonably comparable t.

4.2 Results

Following [2], the primary metric used to evaluate our model is the percentage point-source contribution to total pollutant load. This is evaluated at median river ow of

Figure 8: Posterior predictive density of simulated $\log_{10}$(Orthophosphate) concentration from Bayesian LAM at the Thames (light blue) and actual $\log_{10}$(Orthophosphate) concentration at the Thames (dark blue) in 2000 2004.

each time-period Q_{med}, and calculated as

$$PS\% = \frac{A \; Q_{med}^{B\,1}}{A \; Q_{med}^{B\,1} + C^{\,0} \; Q_{med}^{D\,1}} \; 100: \qquad (5)$$

In the Bayesian framework, this quantity is derived from the joint posterior distribution of $(A; B; C^{\,0}; D)$, yielding a full posterior distribution for PS% from which credible intervals are obtained directly.

Figures 9 and 10 present posterior median PS% estimates with 80% credible intervals across all site-determinand-window combinations (full detail of these values are shown in Appendix D).

Figure 9: Estimated point-source contribution (%) of orthophosphate in the Thames and Wyre over 5-year periods from 2000 to 2024. Circle/Triangle re ects posterior median estimate, and vertical lines represent 80% of corresponding posterior median estimate.

Orthophosphate. The Thames shows clear point-source dominance in 2000 2004 (PS% = 72.8%, 80% CI: 52.7 91.2%), which decreases consistently to 25.2% (8.6

57.1%) by 2020 2024. This trend suggests the impact of progressive STW phosphorus removal of point-source pollutants in the river Thames. The Wyre shows di use-source prominence across all time periods, logical given its rural, agriculturally-dominated location. Orthophosphates also show clear decline from 2000 2004 (PS% = 44.2% (32.3 58.4%) to 2020 2024 (PS% = 0.4 19.5%). Credible intervals for the Wyre are notably smaller than for the Thames across all time periods, re ected by more chemical observations being taken at this site.

Figure 10: Estimated point-source contribution (%) of zinc in the Thames and Wyre over 5-year periods from 2000 to 2024. Circle/Triangle re ects posterior median estimate, and vertical lines represent 80% of corresponding posterior median estimate.

Zinc. The Thames does not show a clear trend over time in point-source contribution of zinc, with most time periods suggesting a contribution of around 25-75%. 2020 2024 however shows the lowest point-source dominance (PS%=11.5%, 80% CI: 4.8 25.5%). This trend is also apparent in the river Wyre, where the 2020 2024 period experienced its lowest point-source contribution of 6.9% (80% CI: 2.4 19.1%), while all other period estimates at median river ow were between 14.9 27.4%.

While these results do indeed suggest clear changes in point and di use-source pollution in these rivers over this 25-year period, there are clear limitations in these models that mean we should be careful before making any assumptions based on these results. Primarily, even after splitting our data into 5-year periods in an attempt to deal with temporal trends, there still exists non-stationarity in our data (as shown in Figure 6). As a result, we aim to develop a model that can handle non-stationary data.

5 Bayesian LAM with time-varying parameters (TV-LAM)

Even within ve-year windows, systematic residual trends persist in the baseline LAM (Figure 6), indicating the static parameter assumption remains violated. Examining baseline posterior estimates, B and D vary across periods but with substantially overlapping credible intervals in all site-determinand cases, suggesting this variation is largely indistinguishable from noise (Appendix C). By contrast, A shows a systematic decline whose magnitude consistently exceeds posterior uncertainty, particularly at the Thames for orthophosphate, where it falls from 3.96 (95% CI: 2.45 6.66) to 0.78 (0.07 2.06). This motivates a framework in which A alone is treated as time-varying.

These results align with ecological background. A is an intercept-like scale parameter capturing the magnitude of point-source inputs, which genuinely changes over time as STW infrastructure improves and regulatory pressure evolves. B, C^0 and D re ect more stable catchment properties and show no systematic trend across periods (Appendix C). We therefore opt for a framework in which A is modelled as a time-varying parameter while B, C^0 and D are treated as xed constants.

Model. Mathematically, this Bayesian LAM with time-varying parameters is written as

$$C_i = A_{t_i} \; Q_i^{B\,1} + C^0 \; Q_i^{D\,1} + "_i; \quad "_i \;\; N(0; \;\; ^2) \qquad (6)$$

Random walk structure. The trajectory of A_t is modelled on the log-scale to enforce positivity, de ned as $_t = \log A_t$. This follows a rst-order random walk structure [8] such that, for $t = 2; :::; N$:

$$_t = _{t\,1} + _A \; ^A_t; \quad ^A_t \overset{iid}{} N(0; 1); \qquad (7)$$

where $_A$ is a variance hyperparameter controlling the rate of temporal change, such that larger values allow adaptation to changing catchment conditions.

Priors. Hyperparameters and initial states are assigned the following priors:

$$_1 \;\; N(0; 2); \qquad (8)$$

$$B \;\; N(0{:}5; 5)j_{\,[0;1]}; \quad D \;\; N(1{:}3; 0{:}5)j_{\,>1}; \qquad (9)$$

$$N^{\,+}(0; 1); \quad _A \;\; N^{\,+}(0; 0{:}2): \qquad (10)$$

The prior on $_1$ was initialised to be non-informative across the range of plausible values of A from the baseline Bayesian LAM (evident by the range of A values, 0.27 15.33, for each site-determinand's rst period) (Appendix C). Hyperprior $_A$ is assigned weakly informative half-normal distributions, allowing plausible temporal drift in A_t. Priors on B and D are identical to the baseline LAM, following [2].

In the Bayesian LAM baseline model, the estimate of C^0 varies largely based on determinand, falling in the range of 0.05 0.11 for orthophosphate, and 1.76 3.90 for zinc. As C^0 varies substantially by determinand (0.05 0.11 for orthophosphate versus 1.76 3.90 for zinc) we use determinand-speci c priors:

$$\text{Orthophosphate: } \log C^0 \;\; N(\,2; 1) \qquad (11)$$

$$\text{Zinc: } \log C^0 \;\; N(0{:}7; 1) \qquad (12)$$

Model implementation. This model was constructed using Stan [18] and run in R [17].

5.1 Evaluation of Model

All four TV-LAM models achieved satisfactory convergence, with all scalar parameters obtaining $\hat{R}$ 1:01 and e ective sample sizes of over 400 (with the exception of , obtaining 346 for the zinc model at the Thames) [11]. See Appendix F for convergence results of these scalar parameters for each model. No divergent transitions were observed.

Traceplots of these scalar parameters were investigated for each TV-LAM, with Orthophosphate at the Thames model shown in Figure 11 (with all other TV-LAM traceplots found in Appendix E.1).

Figure 11: Traceplots of scalar parameters $\log A_{init}$; B; C^{0}; D; ; $_{A}$ estimated for TV-LAM for orthophosphate at the Thames (constructed using Stan).

As shown in the traceplot, all scalar parameters experience good mixing, with each chain exhibiting stationary, well-mixed behaviour around a stable posterior region. Traceplots of other determinand-site TV-LAM combinations also showed good mixing, with B at both zinc sites exhibiting posterior mass concentrated near the upper boundary of its constrained support $[0; 1]$, indicating the constraint is active and that B is poorly identi ed for zinc (see Appendix E.1).

Residuals were plotted over time for each TV-LAM, with Orthophosphate at the Thames model shown in Figure 12 (with all other TV-LAM residual plots found in Appendix E.2).

Figure 12: Bayesian TV-LAM Residual vs Time on Thames-Orthophosphate data with smoothed blue LOESS trend line.

The residual vs time plot of orthophosphate concentration at the Thames (Figure 12) is a key indicator that the time-varying LAM framework mitigates the prior issues caused by non-stationarity in our baseline Bayesian LAM (Figure 6). The LOESS smoothing line sticks consistently to the zero-line, indicating that our modelling framework mitigates any major systematic temporal trend. Other determinand-site combinations also exhibit adequate stationarity, with relatively at LOESS lines across all site-determinand combinations (see Appendix E.2).

Posterior predictive checks are shown for orthophosphate at the Thames (Figure 13), with remaining site-determinand combinations in Appendix E.3.

Figure 13: Posterior predictive density of simulated $\log_{10}$(Orthophosphate) concentration from TV-LAM at the Thames (light blue) and actual $\log_{10}$(Orthophosphate) concentration at the Thames (dark blue).

The posterior predictive density mostly envelopes the central mass of the observed Thames-orthophosphate concentration distribution but does not reproduce the bimodal structure, which likely re ects distinct regulation regime changes pooled across the full 25-year period. The Wyre-orthophosphate model also captured the observed density well. Both zinc models however produced simulated densities with peaks approximately half the height of the observed distribution, suggesting the models overestimate residual variance. This is likely due to a single xed being unable to capture heteroskedasticity across a 25-year period with varying levels of change in A_t. Posterior predictive checks for the remaining site-determinand combinations (see Appendix E.3) show similar patterns, with orthophosphate ts capturing the observed density well and both zinc models overestimating residual variance.

5.2 Results

We now evaluate the point-source contribution to total pollutant load for each site-determinand TV-LAM. PS_t% is computed using Equation 5, evaluated at the full-period site-speci c median ow Q_{med}, such that the resulting trajectory re ects only structural change in the point-source intercept A_t over time. This di ers slightly from the windowed baseline LAM, where PS% is evaluated at the window-speci c median ow, providing a more consistent reference condition

across the full 25-year period. Figures 14 and 15 present the posterior median trajectory of A_t with 80% credible intervals for orthophosphate and zinc respectively, providing the mechanistic basis for the PS% trends discussed below.

Figure 14: Posterior median estimates of A $_t$ for orthophosphate at the Thames (red) and Wyre (blue) with 80% credible intervals.

Figure 15: Posterior median estimates of A $_t$ for zinc at the Thames (red) and Wyre (blue) with 80% credible intervals.

As the scale parameter of point-source inputs diminishes over time, point-source contribution to total load falls correspondingly, independently of changes in river ow. The Thames orthophosphate trajectory in particular, falling from approximately 5 in 2000 to below 1 by 2010 and stabilising thereafter, re ects a structural shift in point-source loading consistent with regulatory intervention rather than gradual ecological change.

Figures 16 and 17 present posterior median PS% estimates with 80% credible intervals for the Thames and Wyre sites.

Orthophosphate. The Thames shows clear point-source dominance from 2000 2006, with the 80% credible interval of the posterior estimate lying consistently above 75%. Point-source dominance then declines sharply between approximately 2007 2010, broadly coinciding with the mandatory introduction of phosphorus stripping at larger sewage treatment works under the Urban Wastewater Treatment Directive [16, 3], before stabilising at around 50% with 80% credible interval spanning 25 75% throughout to 2024.

The Wyre exhibits primarily di use-source dominance throughout the period, with initial posterior median estimates of point-source contribution between 25 55% and relatively narrow credible intervals of around 5-10% in width, re ecting the greater volume of observations at this site. From approximately 2018, there is a distinct and sustained decline in point-source contribution from around 30% to almost 0% by 2024, suggesting that total orthophosphate load in the Wyre at St Michaels has become almost entirely di use-source dominated by the end of the study period. This is consistent with its rural, agriculturally-dominated catchment.

Figure 16: Posterior median estimates of point-source contribution (%) of orthophosphate in the Thames (red) and Wyre (blue). The corresponding shaded regions represent 80% credible interval in these posterior estimates.

These ndings are broadly consistent with the windowed baseline LAM (Section 4.2), which estimated Thames orthophosphate point-source contribution declining from 72.8% (80% CI: 52.7 91.2%) in 2000 2004 to 25.2% (8.6 57.1%) by 2020 2024, suggesting suitable speci cation of the TV-LAM framework despite the methodological di erences in ow reference condition.

Figure 17: Posterior median estimates of point-source contribution (%) of zinc in the Thames (red) and Wyre (blue). The corresponding shaded regions represent 80% credible interval in these posterior estimates.

Zinc. Thames zinc remains predominantly point-source dominated throughout the period, with posterior median estimates broadly declining from around 75 80% in

2000 toward 50 60% by 2024, though 80% credible intervals remain wide, spanning approximately 50 90% in earlier periods and 30 85% by 2020 2024. The Wyre exhibits consistently lower point-source contribution, with posterior medians primarily in the region of 12 25% and very wide credible intervals spanning close to 0 60% for much of the period. The substantially wider credible intervals for both sites relative to orthophosphate re ect the larger proportion of censored zinc observations (around 20% at both sites) and the boundary behaviour of B identi ed in Section 5.1, which together limit the precision of source apportionment for this determinand. These zinc results should therefore be interpreted with considerably more caution than the orthophosphate ndings.

These ndings di er from the windowed baseline LAM (Section 4.2) in the estimated level of point-source contribution, with the baseline suggesting Thames zinc point-source contribution declined from 39.4% (80% CI: 18.7 62.7%) in 2000 2004 to 11.5% (4.8 25.5%) by 2020 2024, considerably lower than the TV-LAM trajectory. Both models nonetheless agree directionally, with point-source contribution declining over the period and uncertainty remaining high throughout.

6 Model Comparison

6.1 Residual Temporal Bias

As made evident in Section 4.1, a fundamental limitation of the windowed baseline Bayesian LAM is temporal non-stationarity. Even within ve-year periods, a systematic downward LOESS trend in the residuals (Figure 6) indicates that the static parameter assumption is violated. By modelling A_t as a time-varying parameter following a rst-order random walk, the TV-LAM explicitly accommodates this non-stationarity, eliminating the systematic residual bias as evidenced by the adequately at LOESS smoothing line in Figure 12 (with corresponding plots for all other site-determinand combinations shown in Appendix E.2).

Beyond mitigating residual bias, the TV-LAM framework removes the need for arbitrary temporal windowing. In the baseline LAM, the stationarity assumption necessitates restricting model tting to short time periods in which the ow-concentration relationship is approximately stable. The TV-LAM instead accommodates the full 25-year observation record within a single coherent model, eliminating the arbitrary choice of window boundaries and the information loss that occurs at each boundary transition.

6.2 In-Sample Goodness of Fit

In-sample goodness of t for both modelling strategies is assessed using continuous ranked probability score (CRPS), a negatively-oriented scoring rule, computed on the 2000 2024 observation period [12]. CRPS evaluates a predicted probability distribution against the actual observed value, assessing both calibration and sharpness. This comparison is asymmetric: each baseline window is tted only to the observations it is evaluated on, whereas the TV-LAM characterises the full record simultaneously, structurally favouring the baseline. The TV-LAM's CRPS advantage at the Thames sites is consequently a conservative lower bound on its true performance gain, while the baseline's marginal Wyre advantage may partly re ect this structural bene t rather than genuine superior predictive calibration.

Table 5 shows the mean CRPS values of the baseline LAM and TV-LAMs for each site-determinand combination.

Table 5: Mean CRPS comparison between the windowed baseline LAM and TV-LAM across all site-determinand combinations, evaluated on the shared 2000 2024 observation period. Lower values indicate better t. Wyre zinc TV-LAM evaluated from 2005 onward to match baseline LAM time-windows. Bold denotes the better- tting model.

Site	Determinand	Baseline	TV-LAM
Thames	Orthophosphate	0.105	**0.089**
Wyre	Orthophosphate	**0.089**	0.095
Thames	Zinc	0.079	**0.065**
Wyre	Zinc	**0.076**	0.082

The TV-LAM achieves lower mean CRPS for both Thames orthophosphate and Thames zinc, the two site-determinand combinations exhibiting the strongest long-run decline in A_t (Figures 14 and 15). The baseline achieves lower CRPS for Wyre orthophosphate and zinc, where A_t remains comparatively stable throughout the period, and thus the windowed approach is well-speci ed within each ve-year period. These results are ecologically coherent: the TV-LAM's advantage materialises precisely where structural non-stationarity is greatest, and is absent where the baseline's simpler parameterisation is adequate.

Notably, the baseline's CRPS advantage at the Wyre is marginal: 0.089 vs 0.095 (reduction of approximately 6.3%) for orthophosphate and 0.076 vs 0.082 (7.9%) for zinc, whereas the TV-LAM's advantage at the Thames is more pronounced, with CRPS reductions of 15.2% for orthophosphate and 17.7% for zinc relative to the baseline. Given that each baseline window is tted to approximately 20% of the observations available to the TV-LAM, these CRPS reductions represent a meaningful improvement in goodness of t by a model operating under greater constraint.

Figure 18 shows observation-level CRPS for the windowed baseline LAM and TV-LAMs for orthophosphate at the Thames.

Figure 18 reveals that the aggregate CRPS di erences in Table 5 are not uniformly distributed across the observation period but are concentrated in speci c temporal regions. For Thames orthophosphate, the TV-LAM's advantage is most pronounced between approximately 2007 2014, a period of rapid decline in A_t (Figure 14), and where the baseline model is most severely stressed by non-stationarity. For Thames zinc, the TV-LAM maintains a consistent advantage throughout the period, consistent with the persistent but noisier downward trend in A_t at this site. For the Wyre site-determinand combinations, the LOESS lines cross repeatedly with no sustained advantage for either model, corroborating the marginal aggregate di erences reported

Figure 18: LOESS-smoothed observation-level CRPS for the windowed baseline LAM (red) and TV-LAM (blue) for orthophosphate at the Thames.

in Table 5. Taken together, Figures 14 and 18 establish that the TV-LAM's goodness-of- t advantage is mechanistically linked to the rate of structural change in point-source inputs: where A_t evolves rapidly, the TV-LAM characterises the concentration record more accurately than a sequence of static windowed models.

Overall, these results indicate that the TV-LAM is the preferred framework for characterising source apportionment over non-stationary long-term periods, while the windowed baseline remains adequate where catchment conditions are approximately stable. Whether these patterns generalise beyond the Thames and Wyre, and what they imply for EA monitoring and investment decisions, is considered in the Discussion (Section 7).

7 Discussion

7.1 Main Findings

The central nding of this analysis is that the Bayesian LAM and TV-LAM frameworks together document ecologically coherent, site and determinand-speci c trajectories in point-source contribution over the 2000 2024 period.

For orthophosphate, both frameworks document a regulatory success story consistent with the progressive implementation of phosphorus stripping at larger STWs under the Urban Wastewater Treatment Directive [16, 3]. At the Wyre, TV-LAM posterior estimates shift from approximately 50% point-source contribution in 2000 to near 0% by 2024, re- ecting near-complete elimination of STW dominance in a predominantly rural catchment with comparatively few and smaller STW discharges. At the Thames, point-source contribution declines sharply between 2007 2010 before stabilising at around 50%, suggesting that the substantially larger aggregate STW load on this major urban catchment prevents di use-source dominance from being achieved, an outcome broadly consistent with national HMS trends and prior LAM studies on UK lowland rivers [2, 4].

For zinc, both models indicate persistent point-source dominance at the Thames and di use-source dominance at the Wyre throughout, though wide 80% credible intervals limit the precision of these apportionments considerably. While the directional agreement between models provides some con dence, the credible interval width is such that compliance-relevant thresholds cannot be reliably distinguished. From an EA operational perspective, the wide credible intervals mean point and di use contributions to zinc cannot be distinguished with su cient precision for compliance decisions at either site.

The TV-LAM successfully characterises these trajectories across the full 25-year study period, and eliminates the systematic residual temporal bias that persists even within ve-year windows in the baseline Bayesian LAM. TV-LAM consistently tting our data better than baseline LAM in periods of large structural change in A_t indicates that the random walk prior successfully captures non-stationarity rather than absorbing residual variation. Where catchment structure is stable, as at the Wyre, the simpler windowed baseline remains an adequate and more parsimonious alternative.

7.2 Limitations

Several limitations of this analysis warrant acknowledgement. The TV-LAM treats C^0 as xed because modelling it as time-varying alongside A_t produced posterior non-identi ability, indicating the data cannot simultaneously identify two freely-evolving trajectories from a single concentration- ow relationship. Consequently, the framework cannot detect structural changes in di use-source inputs over time, a meaningful ecological caveat particularly for the Wyre, where agricultural land management practices have evolved considerably over the study period.

For zinc speci cally, B was found to concentrate posterior mass at the upper boundary of its constrained support $[0; 1]$ at both sites, indicating the point/di use decomposition may itself be poorly identi ed for this determinand. This likely re ects a structural limitation of applying the LAM to zinc: B concentrating at its upper boundary suggests the point-source term is becoming load-increasing with ow, which is physically inconsistent with dilution of a constant point-source input and indicates the two-term power-law decomposition may be poorly suited to this determinand. The wide credible intervals for zinc should therefore be understood as re ecting both high censoring rates and the weaker hydrological separability of point and di use sources for zinc.

Furthermore, posterior predictive checks revealed that all TV-LAMs overestimated residual variance, likely because a single xed cannot capture heteroskedasticity across a 25-year period of varying structural change in A_t. CRPS values should also be interpreted as comparative goodness-of- t metrics on training data only: each baseline window is tted to approximately 20% of the observations the TV-LAM must simultaneously characterise, conferring a structural training advantage on the baseline that precludes direct comparison as a measure of predictive ability. Inferring out-of-sample performance would require leave-one-out cross-validation on matched held-out data. Results span only two sites and two determinands, and as such ndings should not be directly

extrapolated to other catchments.

7.3 Future Directions

The TV-LAM framework developed here was applied to two sites and two determinands; extending this to all 135 HMS sites across England would establish whether the long-term declines in point-source contribution identified at the Thames and Wyre are representative of the broader river network. This would allow the EA to identify catchments that remain point-source dominated for orthophosphate, and thus inform decisions on where further STW investment would yield the greatest water quality improvement. From a modelling perspective, the fixed assumption could be addressed by treating observational variance as time-varying, better capturing heteroskedasticity across periods of rapid structural change in A_t. For zinc specifically, future work might consider whether alternative model formulations would yield better-identified source apportionment. The modified LAM of [7], which accounts for upstream inflow contributions, represents one such candidate for determinands with more complex catchment source pathways.

References

[1] Environment Agency. Hydrology Data Explorer. 2025. url: https://environment.data.gov.uk/hydrology/explore (visited on 05/28/2026).

[2] M. J. Bowes et al. Modelling of phosphorus inputs to rivers from diffuse and point sources. In: Science of the Total Environment (2008). url: https://pubmed.ncbi.nlm.nih.gov/18367235/.

[3] Michael J. Bowes et al. Changes in point and diffuse source phosphorus inputs to the River Frome (Dorset, UK) from 1966 to 2006. In: Science of the Total Environment (2009).

[4] Michael J. Bowes et al. Identifying priorities for nutrient mitigation using river concentration flow relationships: The Thames basin, UK. In: Journal of Hydrology (2014).

[5] Paul-Christian Bürkner. brms: An R Package for Bayesian Multilevel Models Using Stan. In: Journal of Statistical Software 80 (2017).

[6] S. R. Carpenter et al. Nonpoint Pollution of Surface Waters with Phosphorus and Nitrogen. In: Ecological Applications (1998).

[7] Dingjiang Chen, Randy A. Dahlgren, and Jun Lu. A modified load apportionment model for identifying point and diffuse source nutrient inputs to rivers from stream monitoring data. In: Journal of Hydrology 501 (2013).

[8] James Durbin and Siem Jan Koopman. Time Series Analysis by State Space Methods. Oxford: Oxford University Press, 2001.

[9] Environment Agency. Interim H4 indicator 2024: exposure and adverse effects of chemicals on wildlife in the environment Supporting information and data. Accessed: June 2026. 2024. url: https://assets.publishing.service.gov.uk/media/664f2af14f29e1d07fadcd89/Exposure_and_adverse_effects_of_chemicals_on_wildlife_in_the_environment_-_interim_H4_indicator_2024.pdf.

[10] European Commission. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy (Water Framework Directive). Official Journal of the European Union, L 327. 2000.

[11] A. Gelman and D.B. Rubin. Inference from iterative simulation using multiple sequences. In: Statistical Science (1992).

[12] Tilmann Gneiting and Adrian E. Raftery. Strictly Proper Scoring Rules, Prediction, and Estimation. In: Journal of the American Statistical Association (2007). url: https://sites.stat.washington.edu/raftery/Research/PDF/Gneiting2007jasa.pdf.

[13] GOV.UK. ENV-16 Harmonised Monitoring Scheme datasets. url: https://www.gov.uk/government/statistical-data-sets/env-16-harmonised-monitoring-scheme-datasets (visited on 05/28/2026).

[14] GOV.UK. State of the water environment: long-term trends in river quality in England: 2024. In: GOV.UK (2025). url: https://www.gov.uk/government/publications/state-of-the-water-environment-indicator-b3-supporting-evidence/state-of-the-water-environment-long-term-trends-in-river-quality-in-england-2024 (visited on 06/08/2026).

[15] J. Hilton et al. How green is my river? A new paradigm of eutrophication in rivers. In: Science of the Total Environment (2006).

[16] Helen P. Jarvie, Colin Neal, and Paul J. A. Withers. Sewage-effluent phosphorus: a greater risk to river eutrophication than agricultural phosphorus? In: Science of the Total Environment (2006).

[17] R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria, 2023. url: https://www.R-project.org/.

[18] Stan Development Team. Stan Modeling Language Users Guide and Reference Manual. 2023. url: https://mc-stan.org.

[19] Simon N. Wood. Generalized Additive Models: An Introduction with R. 2nd ed. Chapman and Hall/CRC, 2017.

Own Work Declaration

Proo ng: Generative AI was used to assist in spelling, grammar, and other checking in the following way: Claude (Anthropic), June 2026 - used for suggesting word-level edits; all changes reviewed and accepted or rejected.

Project studies: Claude (Anthropic), June 2026 - used for debugging assistance during coding.

All statistical modelling, analytical decisions, interpretation of results, and conclusions are my own.

Appendix

Word Count: 4942


Appendix A GAM Diagnostics: Residual vs Fitted graphs



(a) Thames orthophosphate, additive



(b) Wyre orthophosphate, additive



(c) Wyre orthophosphate, interaction

Figure 19: Residuals vs tted for orthophosphate GAM models. Thames orthophosphate interaction model shown in main body (Figure 4). Smooth line tted via LOESS; dashed red line at zero.

(a) Thames zinc, additive

(b) Thames zinc, interaction

(c) Wyre zinc, additive

(d) Wyre zinc, interaction

Figure 20: Residuals vs tted for zinc GAM models. Smooth line tted via LOESS; dashed red line at zero.

Appendix B Bayesian LAM Model Diagnostics


Appendix B.1 Bayesian LAM Residuals vs Flow




(a) Thames orthophosphate, 2005 09



(b) Thames orthophosphate, 2010 14



(c) Thames orthophosphate, 2015 19



(d) Thames orthophosphate, 2020 24

Figure 21: Residuals vs log ow for Thames orthophosphate baseline LAM, 2005 2024. LOESS smooth in blue; dashed line at zero.

(a) Wyre orthophosphate, 2000 04

(b) Wyre orthophosphate, 2005 09

(c) Wyre orthophosphate, 2010 14

(d) Wyre orthophosphate, 2015 19

(e) Wyre orthophosphate, 2020 24

Figure 22: Residuals vs log ow for Wyre orthophosphate baseline LAM, 2000 2024.

(a) Thames zinc, 2000 04

(b) Thames zinc, 2005 09

(c) Thames zinc, 2010 14

(d) Thames zinc, 2015 19

(e) Thames zinc, 2020 24

Figure 23: Residuals vs log ow for Thames zinc baseline LAM, 2000 2024.

(a) Wyre zinc, 2005 09

(b) Wyre zinc, 2010 14

(c) Wyre zinc, 2015 19

(d) Wyre zinc, 2020 24

Figure 24: Residuals vs log ow for Wyre zinc baseline LAM, 2005 2024. No 2000 04 window tted due to monitoring commencing 2003.

Appendix B.2 Bayesian LAM Residuals vs Time



(a) Thames orthophosphate, 2005 09

(b) Thames orthophosphate, 2010 14

(c) Thames orthophosphate, 2015 19

(d) Thames orthophosphate, 2020 24

Figure 25: Residuals vs time for Thames orthophosphate baseline LAM, 2005 2024. Systematic LOESS trend indicates residual non-stationarity.

(a) Wyre orthophosphate, 2000 04

(b) Wyre orthophosphate, 2005 09

(c) Wyre orthophosphate, 2010 14

(d) Wyre orthophosphate, 2015 19

(e) Wyre orthophosphate, 2020 24

Figure 26: Residuals vs time for Wyre orthophosphate baseline LAM, 2000 2024. Systematic LOESS trend indicates residual non-stationarity.

(a) Thames zinc, 2000 04

(b) Thames zinc, 2005 09

(c) Thames zinc, 2010 14

(d) Thames zinc, 2015 19

(e) Thames zinc, 2020 24

Figure 27: Residuals vs time for Thames zinc baseline LAM, 2000 2024. Systematic LOESS trend indicates residual non-stationarity.

(a) Wyre zinc, 2005 09

(b) Wyre zinc, 2010 14

(c) Wyre zinc, 2015 19

(d) Wyre zinc, 2020 24

Figure 28: Residuals vs time for Wyre zinc baseline LAM, 2005 2024. Systematic LOESS trend indicates residual non-stationarity.

Appendix B.3 Bayesian LAM QQ Plots

(a) Thames orthophosphate, 2005 09

(b) Thames orthophosphate, 2010 14

(c) Thames orthophosphate, 2015 19

(d) Thames orthophosphate, 2020 24

Figure 29: QQ plots for Thames orthophosphate baseline LAM residuals, 2005 2024. Dashed line = theoretical normal quantiles.

(a) Wyre orthophosphate, 2000 04

(b) Wyre orthophosphate, 2005 09

(c) Wyre orthophosphate, 2010 14

(d) Wyre orthophosphate, 2015 19

(e) Wyre orthophosphate, 2020 24

Figure 30: QQ plots for Wyre orthophosphate baseline LAM residuals, 2000 2024. Dashed line = theoretical normal quantiles.

(a) Thames zinc, 2000 04

(b) Thames zinc, 2005 09

(c) Thames zinc, 2010 14

(d) Thames zinc, 2015 19

(e) Thames zinc, 2020 24

Figure 31: QQ plots for Thames zinc baseline LAM residuals, 2000 2024. Dashed line = theoretical normal quantiles.