

Sums of powers of integers with a fixed number of binary transitions, and the moments of striped random generators

Mikhail Hogrefe* Claude†

August 2026

Abstract

For $0 \leq k \leq n-1$, let $T_{n,k}$ be the set of integers in $[0, 2^n)$ whose n -bit binary representations have exactly k transitions (adjacent unequal bits), and let $S_{n,k,p} = \sum_{m \in T_{n,k}} m^p$. We show that the generating polynomial of the p -th central power sums over k satisfies a *triangular* linear recursion in n , and therefore admits an explicit eigen-expansion: a closed form as a finite combination of n -th powers of the eigenvalues $2^j(1 + (-1)^j t)$ with explicit rational coefficients in the transition-marking variable t . Extracting the coefficient of t^k yields exact formulas for $S_{n,k,p}$ with $O(k+p)$ terms, and the recursion itself evaluates all k simultaneously in $O(np^2)$ big-integer operations. We give fully explicit results for $p \leq 4$, correct a previously conjectured special case, and derive the exact variance and kurtosis—both for the uniform distribution on $T_{n,k}$ and for the “striped” random generators that motivated the question, in which each bit repeats the previous one with a fixed probability. In the striped model the formulas collapse to short rational expressions; for mean run length m , writing $\beta = 1 - 2/m$, the kurtosis converges as $n \rightarrow \infty$ to $\frac{3(2-\beta)(6+\beta)(8+7\beta)}{5(8-\beta)(2+\beta)^2}$. All formulas have been verified in exact arithmetic against exhaustive enumeration.

1 Setup and notation

Fix $n \geq 1$. Every $m \in [0, 2^n)$ has bits $b_{n-1} \cdots b_0$, and its number of transitions is $\tau(m) = \#\{j \in [1, n-1] : b_j \neq b_{j-1}\}$. Let $T_{n,k} = \{m : \tau(m) = k\}$ and

$$S_{n,k,p} = \sum_{m \in T_{n,k}} m^p.$$

Counting is classical: a string with k transitions is determined by its transition set (a k -subset of the $n-1$ boundaries) and its bottom bit, so $|T_{n,k}| = 2 \binom{n-1}{k}$.

It is convenient to center. Write each bit as a sign $\epsilon_i = 2b_i - 1 \in \{\pm 1\}$ and set

$$c(m) = 2m - (2^n - 1) = \sum_{i=0}^{n-1} \epsilon_i 2^i.$$

*M.H. posed the questions: the moments of Malachite’s striped random generators, and the conjectured special case revisited within.

†Claude (Anthropic) worked out the mathematics and wrote the note. Any errors are the second author’s.

Factor out the bottom sign: $\epsilon_i = \epsilon_0 \prod_{j \leq i} s_j$, where $s_j = \epsilon_j \epsilon_{j-1} \in \{\pm 1\}$ equals -1 exactly at transitions. The map $m \leftrightarrow (\epsilon_0, s_1, \dots, s_{n-1})$ is a bijection, $\tau(m) = \#\{j : s_j = -1\}$ does not depend on ϵ_0 , and

$$c(m) = \epsilon_0 Y, \quad Y = Y(s) = \sum_{i=0}^{n-1} 2^i \prod_{j=1}^i s_j. \quad (1)$$

Summing over $\epsilon_0 = \pm 1$ shows that all odd central power sums vanish (this is the complement symmetry $m \mapsto 2^n - 1 - m$ of $T_{n,k}$), and that everything is encoded in the polynomials

$$\widehat{M}_p(n) = \widehat{M}_p(n)(t) := \sum_{s \in \{\pm 1\}^{n-1}} t^{\#\{j : s_j = -1\}} Y(s)^p,$$

via

$$\sum_{m \in T_{n,k}} c(m)^p = \begin{cases} 2[t^k] \widehat{M}_p(n), & p \text{ even,} \\ 0, & p \text{ odd,} \end{cases} \quad S_{n,k,p} = \frac{1}{2^p} \sum_{\substack{j=0 \\ j \text{ even}}}^p \binom{p}{j} (2^n - 1)^{p-j} \cdot 2[t^k] \widehat{M}_j(n). \quad (2)$$

The second identity is the binomial expansion of $m^p = ((c + 2^n - 1)/2)^p$, with the odd- j terms dropped.

2 The triangular recursion and its eigen-expansion

Theorem 1. *The polynomials $\widehat{M}_p$ satisfy, for $n \geq 2$,*

$$\widehat{M}_p(n) = \sum_{j=0}^p \binom{p}{j} 2^j (1 + (-1)^j t) \widehat{M}_j(n-1), \quad \widehat{M}_p(1) = 1. \quad (3)$$

Consequently, since the coefficient matrix is lower triangular with distinct eigenvalues $\lambda_j = \lambda_j(t) = 2^j (1 + (-1)^j t)$,

$$\widehat{M}_p(n) = \sum_{j=0}^p c_{p,j}(t) \lambda_j(t)^{n-1}, \quad (4)$$

where the rational functions $c_{p,j}$ are determined by

$$c_{p,j} = \frac{\sum_{i=j}^{p-1} \binom{p}{i} 2^i (1 + (-1)^i t) c_{i,j}}{\lambda_j - \lambda_p} \quad (j < p), \quad c_{p,p} = 1 - \sum_{j < p} c_{p,j}.$$

Proof. Splitting off the boundary variable s_1 in (1) gives the self-similarity $Y = 1 + 2s_1 \widetilde{Y}$, where $\widetilde{Y}$ is the same functional of $(s_2, \dots, s_{n-1})$. Raise to the p -th power, expand, and sum first over $s_1 \in \{\pm 1\}$ using $\sum_{s=\pm 1} t^{[s=-1]} s^j = 1 + (-1)^j t$, then over the remaining variables; this is (3). The expansion (4) is the standard solution of a triangular linear recursion with distinct eigenvalues; matching the coefficient of λ_j^{n-1} on both sides of (3) gives the formula for $c_{p,j}$ with $j < p$, and the initial condition $\widehat{M}_p(1) = 1$ fixes $c_{p,p}$. $\square$

Theorem 1 answers the algorithmic question at once: iterating (3) computes $\widehat{M}_0, \dots, \widehat{M}_p$ as polynomials in t , hence $S_{n,k,p}$ for every k simultaneously, in $O(np^2)$ arithmetic operations on big integers—versus $\Theta(2^n)$ for enumeration. The expansion (4) is moreover a closed form, uniform in n ; the tables for $p \leq 4$ are as follows (each was generated by exact rational-function arithmetic and verified against exhaustive enumeration; note the $t \mapsto -t$ symmetry between $c_{p,j}$ and $c_{p,p-j}$ up to powers of 2).

$$\widehat{M}_0(n) = (1+t)^{n-1}, \quad (5)$$

$$\widehat{M}_1(n) = \frac{(1+t)^n - 2^n(1-t)^n}{3t-1}, \quad (6)$$

$$\widehat{M}_2(n) = \frac{(3-t)(1+t)^{n-1}}{3(1-3t)} + \frac{2^{n+1}(1-t)^{n+1}}{(3t-1)(3t+1)} + \frac{4^n(3+t)(1+t)^{n-1}}{3(1+3t)}, \quad (7)$$

$$\widehat{M}_3(n) = \frac{(t-7)(t+1)(1+t)^{n-1}}{(3t-1)(9t-7)} + \frac{(t+3)2^n(1-t)^n}{(3t-1)(3t+1)} + \frac{(t+3)4^n(1+t)^n}{(3t-1)(3t+1)} + \frac{(t-7)8^n(1-t)^n}{(3t-1)(9t-7)}, \quad (8)$$

$$\begin{aligned} \widehat{M}_4(n) = & \frac{(15-t)(7-5t)(1+t)^{n-1}}{15(1-3t)(7-9t)} + \frac{8(7+t)2^{n-1}(1-t)^{n+1}}{(3t-1)(3t+1)(9t+7)} + \frac{8(t^2-9)4^{n-1}(1+t)^{n-1}}{3(9t^2-1)} \\ & + \frac{32(t-7)8^{n-1}(1-t)^{n+1}}{(9t^2-1)(9t-7)} + \frac{16(15+t)(7+5t)16^{n-1}(1+t)^{n-1}}{15(1+3t)(7+9t)}. \end{aligned} \quad (9)$$

Each summand has poles, but the sum is a polynomial in t of degree $n-1$; the singularities cancel identically, which is a useful internal check.

Remark 2 (Krawtchouk form). There is an equivalent “static” formulation that avoids the recursion. Expanding $Y^p = \sum_{i_1, \dots, i_p} 2^{i_1 + \dots + i_p} \prod_j s_j^{e_j}$ with $e_j = \#\{\ell : i_\ell \geq j\}$ and summing over s shows, for even p ,

$$\sum_{m \in T_{n,k}} c(m)^p = 2 \sum_{i_1, \dots, i_p=0}^{n-1} 2^{i_1 + \dots + i_p} K_k(O(i); n-1),$$

where, if $i_{(1)} \leq \dots \leq i_{(p)}$ is the sorted tuple, $O(i) = (i_{(2)} - i_{(1)}) + (i_{(4)} - i_{(3)}) + \dots + (i_{(p)} - i_{(p-1)})$ is the alternating gap sum, and $K_k(d; N) = \sum_j (-1)^j \binom{d}{j} \binom{N-d}{k-j}$ is the binary Krawtchouk polynomial, i.e. $[t^k](1+t)^{N-d}(1-t)^d$. The eigen-expansion results from evaluating this p -fold sum by nested geometric series.

3 Extracting a fixed number of transitions

All denominators in (7)–(9) are products of polynomials *linear* in t , so coefficient extraction is mechanical.

Lemma 3 (extraction). *For any $A, B \geq 0$ and constant a ,*

$$[t^k] \frac{(1+t)^A(1-t)^B}{1-at} = \sum_{i=0}^k a^{k-i} \sum_j (-1)^j \binom{B}{j} \binom{A}{i-j}.$$

Terms with several linear factors in the denominator are first split by partial fractions in t . Applied to (4), this produces, for each fixed even p , an explicit formula for $\sum_{T_{n,k}} c^p$ with $O(k+p)$ terms; the relevant values of a are ± 3 for $p = 2$ and $\pm 3, \pm \frac{9}{7}$ for $p = 4$.

For $p = 2$ the assembled result is compact. Writing $\Sigma_{n,k,2} = \sum_{m \in T_{n,k}} (2m - 2^n + 1)^2$:

$$\Sigma_{n,k,2} = 2 \sum_{i=0}^k \left\{ \frac{3^{k-i}}{3} \left[3 \binom{n-1}{i} - \binom{n-1}{i-1} \right] + \frac{(-3)^{k-i} 4^n}{3} \left[3 \binom{n-1}{i} + \binom{n-1}{i-1} \right] - 2^n (-1)^i (3^{k-i} + (-3)^{k-i}) \binom{n+1}{i} \right\}, \quad (10)$$

and then

$$S_{n,k,2} = \frac{\Sigma_{n,k,2}}{4} + \frac{(2^n - 1)^2}{2} \binom{n-1}{k}. \quad (11)$$

Corollary 4 (low k). *Specializing (10)–(11),*

$$S_{n,1,2} = 4^n \left(n - \frac{7}{3} \right) + 2^{n+1} + n + \frac{1}{3}, \quad S_{n,2,2} = 4^n \left(\frac{n^2}{2} - \frac{17n}{6} + \frac{19}{3} \right) - 2^n (n^2 - n + 10) + \frac{n^2}{2} - \frac{n}{6} + \frac{11}{3}.$$

Remark 5. The value of $S_{n,1,2}$ corrects the formula conjectured in the original question, which omits the “ $+n$ ” term; the two expressions already disagree at $n = 2$, where $T_{2,1} = \{1, 2\}$ gives $S_{2,1,2} = 5$.

Corollary 6 (moments of the uniform distribution on $T_{n,k}$). *Let X be uniform on $T_{n,k}$. Then $\mathbb{E}X = (2^n - 1)/2$, all odd central moments vanish, and*

$$\text{Var}(X) = \frac{[t^k] \widehat{M}_2(n)}{4 \binom{n-1}{k}}, \quad \text{Kurt}(X) = \frac{\mu_4}{\mu_2^2} = \frac{\binom{n-1}{k} [t^k] \widehat{M}_4(n)}{([t^k] \widehat{M}_2(n))^2}.$$

For every fixed k , $\text{Kurt}(X) \rightarrow 1$ as $n \rightarrow \infty$: with few transitions the value is dominated by the top run and the distribution becomes two-lobed. (For $k = 0$ and $k = n - 1$ the kurtosis is exactly 1.) Sample values at $n = 64$: 1.0297 at $k = 1$, 1.2302 at $k = 8$, 1.7869 at $k = 31$.

Remark 7 (on “closed form”). For each fixed p , (4) with Lemma 3 is an exact formula with $O(k+p)$ terms, uniform in n and k . A single hypergeometric-free expression in all three parameters simultaneously seems out of reach: the k -slices irreducibly involve the partial sums $\sum_{i \leq k} 3^i \binom{n}{i}$ (and $\frac{9}{7}$ -weighted analogues at $p = 4$), which do not telescope.

4 Striped generators: independent transitions

The generators that motivated the question do not fix k . A *striped* generator with mean run length $m \geq 2$ draws b_0 uniformly and then repeats the previous bit with probability $1 - \theta$, where $\theta = 1/m$; that is, the s_j of Section 1 are i.i.d. with $\Pr[s_j = -1] = \theta$. Set

$$\beta = \mathbb{E}[s_j] = 1 - 2\theta = 1 - \frac{2}{m}.$$

The analysis of Theorem 1 goes through verbatim with the marker t replaced by expectation: the recursion for $\mathbb{E}[Y^p]$ has the triangular matrix $\binom{p}{j} 2^j e_j$ with $e_j = 1$ for even j and $e_j = \beta$ for

odd j , hence eigenvalues $1, 2\beta, 4, 8\beta, 16, \dots$. For the centered value $c = 2X - (2^n - 1)$ all odd moments vanish, and the even ones are short rational expressions:

$$\mathbb{E}[c^2] = \frac{1 + 2\beta}{3(2\beta - 1)} + \frac{4\beta^2 (2\beta)^{n-1}}{(2\beta - 1)(\beta - 2)} + \frac{4(2 + \beta) 4^{n-1}}{3(2 - \beta)}, \quad (12)$$

$$\begin{aligned} \mathbb{E}[c^4] = & \frac{(6\beta + 1)(8\beta + 7)}{15(2\beta - 1)(8\beta - 1)} - \frac{8\beta^2(3\beta + 4) (2\beta)^{n-1}}{(2\beta - 1)(2 - \beta)(8 - \beta)} - \frac{8(2\beta + 1)(\beta + 2) 4^{n-1}}{3(2\beta - 1)(\beta - 2)} \\ & + \frac{32\beta^2(4\beta + 3) (8\beta)^{n-1}}{(\beta - 2)(8\beta - 1)(2\beta - 1)} + \frac{16(7\beta + 8)(\beta + 6) 16^{n-1}}{15(8 - \beta)(2 - \beta)}, \end{aligned} \quad (13)$$

with $\text{Var}(X) = \frac{1}{4}\mathbb{E}[c^2]$ and $\text{Kurt}(X) = \mathbb{E}[c^4]/\mathbb{E}[c^2]^2$.

Remark 8 (removable singularities). The parameter values $\beta = \frac{1}{2}$ (mean run length $m = 4$) and $\beta = \frac{1}{8}$ are removable singularities of (12)–(13), caused by eigenvalue collisions ($2\beta = 1$, resp. $8\beta = 1$); evaluate by taking limits, or by running the recursion directly. For example, at $m = 4$,

$$\mathbb{E}[c^2] = \frac{5 \cdot 4^n - 6n - 5}{9}.$$

Corollary 9 (limits). *As $n \rightarrow \infty$,*

$$\text{Var}\left(\frac{X}{2^n}\right) \longrightarrow \frac{2 + \beta}{12(2 - \beta)}, \quad \text{Kurt}(X) \longrightarrow \frac{3(2 - \beta)(6 + \beta)(8 + 7\beta)}{5(8 - \beta)(2 + \beta)^2}.$$

At $\beta = 0$ (unstriped, uniform bits) these are $\frac{1}{12}$ and $\frac{9}{5}$, the variance and kurtosis of the uniform distribution; as $\beta \rightarrow 1$ they tend to $\frac{1}{4}$ and 1, those of a symmetric two-point distribution.

m	β	$\lim \text{Var}(X/2^n)$	$\lim \text{Kurt}(X)$	
2	0	$1/12 \approx 0.0833$	$9/5 = 1.8$	(uniform)
3	$1/3$	$7/60 \approx 0.1167$	$1767/1127 \approx 1.5679$	
4	$1/2$	$5/36 \approx 0.1389$	$897/625 = 1.4352$	
8	$3/4$	$11/60 \approx 0.1833$	≈ 1.2234	
16	$7/8$	$23/108 \approx 0.2130$	≈ 1.1130	
32	$15/16$	$47/204 \approx 0.2304$	≈ 1.0568	

5 Verification

All results were checked in exact rational arithmetic: the eigen-expansions (7)–(9) against exhaustive enumeration of all 2^{n-1} transition strings for $p \leq 4$ and $n \leq 12$; the bridge (2) against the raw integer sums $\sum_m m^p$ over $[0, 2^n)$ for $n \leq 10$ and all k ; the closed forms against an independent iteration of (3) at $n = 40$; the explicit formulas (10)–(11) and Corollaries 4–6 against brute force on the stated ranges; and (12)–(13) against probability-weighted enumeration at $\beta \in \{\frac{1}{3}, \frac{3}{5}, \frac{15}{16}, -\frac{1}{3}\}$.