

REPRODUCED REPORT: REM sleep in naps differentially relates to memory consolidation
in typical preschoolers and children with Down syndrome.

Goffredina Spano¹ & Rebecca L. Gomez^{1,3}

¹ University of Arizona

² University College London

Author Note

This is a reproduction of an article published on October 20, 2018.

Correspondence concerning this article should be addressed to Goffredina Spano,
Department of Psychology, University of Arizona, Tucson, AZ 85821. E-mail:
g.spano@ucla.ac.uk

Abstract

Naps are beneficial for learning in typically developing infants, children, and adults. They show greater retention when a delay between training and test contains sleep than when it is a comparable period of wake. However, individuals with Down syndrome have a high rate of disordered sleep than seen in the typical population. Do they experience the same benefits of sleep on learning? The current experiment suggests they do not. While typically developing preschoolers showed more retention after a period filled with sleep, children with Down syndrome had greater retention after a period of wakefulness.

Keywords: naps, sleep, memory, development, Down syndrome

Word count: X

REPRODUCED REPORT: REM sleep in naps differentially relates to memory consolidation in typical preschoolers and children with Down syndrome.

Methods

Participants

Warning: package 'xtable' was built under R version 3.5.3

Groups	N	Mean_age	PercentFemale
DS	25	9.49	52
TD	24	5.03	54

Materials & Procedure

The goal of this study was to assess the retention of new words with various intervals between training and test. Children received all conditions 1-2 weeks apart. The conditions included: 1. after a 5 min delay 2. after a nap (4 hour delay) 3. after 24 hours

Data analysis

The authors assessed the number of trials needed to reach criterion across conditions and groups.

The first analysis conducted was a repeated measures ANOVA for both wake and nap conditions. The second was a 2x2 ANOVA with delay type as the repeated factor and TD or DS as the between. These were conducted for the 4 and 24 hour delay.

The second was a 2x2 ANOVA with delay type as the repeated factor and TD or DS as the between. These were conducted for the 4 and 24 hour delay.

We used R (Version 3.5.2; R Core Team, 2018) and the R-packages *data.table* (Version 1.12.0; Dowle & Srinivasan, 2019), *dplyr* (Version 0.8.0.1; Wickham, François, Henry, & Müller, 2019), *ggplot2* (Version 3.1.0; Wickham, 2016), *papaja* (Version 0.1.0.9842; Aust &

Barth, 2018), *readxl* (Version 1.3.1; Wickham & Bryan, 2019), and *xtable* (Version 1.8.3; Dahl, Scott, Roosen, Magnusson, & Swinton, 2018) for all our analyses.

Results

Grouping	Timing	meanNTC	SEMNTC
DS	Immediate	1.680000	0.2628054
DS	Sleep	1.640000	0.1620699
DS	Wake	2.080000	0.1993322
TD	Immediate	2.041667	0.2789679
TD	Sleep	1.708333	0.1408973
TD	Wake	1.666667	0.2055980

[1] "factor"

Error: Subjects Df Sum Sq Mean Sq F value Pr(>F) Residuals 48 2.206 0.04596

Error: Subjects:Condition Df Sum Sq Mean Sq F value Pr(>F) Condition 1 0.007

0.00686 0.049 0.825 Residuals 48 6.674 0.13904

Error: Subjects Df Sum Sq Mean Sq GroupPerHr 1 0.151 0.151

Error: Subjects:CondPerHr Df Sum Sq Mean Sq CondPerHr 1 0.4689 0.4689

Error: Within Df Sum Sq Mean Sq F value Pr(>F)

GroupPerHr 1 0.164 0.1641 2.329 0.129

CondPerHr 1 2.349 2.3488 33.348 3.10e-08 **GroupPerHr:CondPerHr 1 1.481**

1.4813 21.032 8.18e-06 Residuals 190 13.382 0.0704

— Signif. codes: 0 “’ **0.001** ” 0.01 ” 0.05 “.” 0.1 “” 1

Error: Subjects Df Sum Sq Mean Sq GroupPerHr 1 0.2538 0.2538

Error: Subjects:CondPerHr Df Sum Sq Mean Sq CondPerHr 1 1.582 1.582

59 Error: Within Df Sum Sq Mean Sq F value Pr(>F)
60 GroupPerHr 1 0.594 0.594 8.552 0.00387 ** CondPerHr 1 5.653 5.653 81.333 < 2e-16
61 ***GroupPerHr:CondPerHr 1 5.362 5.362 77.155 9.23e-16*** Residuals 190 13.205
62 0.069
63 — Signif. codes: 0 “’ ***0.001*** ’’ *0.01* ’” 0.05 “.” 0.1 “” 1

64 Discussion

65 This study challenges the assumption that sleep is beneficial for learning across
66 populations. Contrary to the pattern displayed by typically developing children, children
67 with Down Syndrome showed improved performance during the wake condition as compared
68 to the sleep condition.

References

69

70 Aust, F., & Barth, M. (2018). *papaja: Create APA manuscripts with R Markdown*.

71 Retrieved from <https://github.com/crsh/papaja>

72 Dahl, D. B., Scott, D., Roosen, C., Magnusson, A., & Swinton, J. (2018). *Xtable: Export*

73 *tables to latex or html*. Retrieved from <https://CRAN.R-project.org/package=xtable>

74 Dowle, M., & Srinivasan, A. (2019). *Data.table: Extension of 'data.frame'*. Retrieved from

75 <https://CRAN.R-project.org/package=data.table>

76 R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna,

77 Austria: R Foundation for Statistical Computing. Retrieved from

78 <https://www.R-project.org/>

79 Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.

80 Retrieved from <http://ggplot2.org>

81 Wickham, H., & Bryan, J. (2019). *Readxl: Read excel files*. Retrieved from

82 <https://CRAN.R-project.org/package=readxl>

83 Wickham, H., François, R., Henry, L., & Müller, K. (2019). *Dplyr: A grammar of data*

84 *manipulation*. Retrieved from <https://CRAN.R-project.org/package=dplyr>

Table 1

ANOVA table for Experiment 1

Effect	F	df_1	df_2	MSE	p	$\hat{\eta}_G^2$
Condition	0.05	1	48	0.14	.825	.001

Table 2

ANOVA table for 4 hour delay

Effect	F	df_1	df_2	MSE	p	$\hat{\eta}_G^2$
GroupPerHr	8.55	1	190	0.07	.004	.043
CondPerHr	81.33	1	190	0.07	< .001	.300
GroupPerHr \times CondPerHr	77.15	1	190	0.07	< .001	.289

Table 3

ANOVA table for 24 hour delay

Effect	F	df_1	df_2	MSE	p	$\hat{\eta}_G^2$
GroupPerHr	2.33	1	190	0.07	.129	.012
CondPerHr	33.35	1	190	0.07	< .001	.149
GroupPerHr \times CondPerHr	21.03	1	190	0.07	< .001	.100




Figure 1. Methods




Figure 2. Average number of trials to criterion per group per condition.