

REPRODUCED REPORT: REM sleep in naps differentially relates to memory consolidation  
in typical preschoolers and children with Down syndrome.

Goffredina Spano<sup>1</sup> & Rebecca L. Gomez<sup>1,3</sup>

<sup>1</sup> University of Arizona

<sup>2</sup> University College London

#### Author Note

This is a reproduction of an article published on October 20, 2018.

Correspondence concerning this article should be addressed to Goffredina Spano,  
Department of Psychology, University of Arizona, Tucson, AZ 85821. E-mail:  
g.spano@ucla.ac.uk

## Abstract

11

12 Naps are beneficial for learning in typically developing infants, children, and adults. They  
13 show greater retention when a delay between training and test contains sleep then when it is  
14 a comparable period of wake. However, individuals with Down syndrome have a high rate of  
15 disordered sleep than seen in the typical population. Do they experience the same benefits of  
16 sleep on learning?

17

*Keywords:* naps, sleep, memory, development, Down syndrome

18

Word count: X

REPRODUCED REPORT: REM sleep in naps differentially relates to memory consolidation in typical preschoolers and children with Down syndrome.

## Methods

### Participants

**##** Warning: package 'xtable' was built under R version 3.5.3

Groups	N	Mean_age	PercentFemale
DS	25	9.49	52
TD	24	5.03	54

### Materials & Procedure

The goal of this study was to assess the retention of new words with various intervals between training and test. Children received all conditions 1-2 weeks apart. The conditions included: 1. after a 5 min delay 2. after a nap (4 hour delay) 3. after 24 hours

### Data analysis

The authors assessed the number of trials needed to reach criterion across conditions and groups.

The first analysis conducted was a repeated measures ANOVA for both wake and nap conditions. The second was a 2x2 ANOVA with delay type as the repeated factor and TD or DS as the between. These were conducted for the 4 and 24 hour delay.

The second was a 2x2 ANOVA with delay type as the repeated factor and TD or DS as the between. These were conducted for the 4 and 24 hour delay.

We used R (Version 3.5.2; R Core Team, 2018) and the R-packages *data.table* (Version 1.12.0; Dowle & Srinivasan, 2019), *dplyr* (Version 0.8.0.1; Wickham, François, Henry, & Müller, 2019), *ggplot2* (Version 3.1.0; Wickham, 2016), *papaja* (Version 0.1.0.9842; Aust &

Barth, 2018), *readxl* (Version 1.3.1; Wickham & Bryan, 2019), and *xtable* (Version 1.8.3; Dahl, Scott, Roosen, Magnusson, & Swinton, 2018) for all our analyses.

## Results

Grouping	Timing	meanNTC	SEMNTC
DS	Immediate	1.680000	0.2628054
DS	Sleep	1.640000	0.1620699
DS	Wake	2.080000	0.1993322
TD	Immediate	2.041667	0.2789679
TD	Sleep	1.708333	0.1408973
TD	Wake	1.666667	0.2055980

[1] “factor”

Error: Subjects Df Sum Sq Mean Sq F value Pr(>F) Residuals 48 4.412 0.09192

Error: Subjects:Condition Df Sum Sq Mean Sq F value Pr(>F) Condition 1 0.014

0.01372 0.049 0.825 Residuals 48 13.348 0.27808

Error: Within Df Sum Sq Mean Sq F value Pr(>F) Residuals 98 4.568 0.04661

## Discussion

## References

- Aust, F., & Barth, M. (2018). *papaja: Create APA manuscripts with R Markdown*. Retrieved from <https://github.com/crsh/papaja>
- Dahl, D. B., Scott, D., Roosen, C., Magnusson, A., & Swinton, J. (2018). *Xtable: Export tables to latex or html*. Retrieved from <https://CRAN.R-project.org/package=xtable>
- Dowle, M., & Srinivasan, A. (2019). *Data.table: Extension of 'data.frame'*. Retrieved from <https://CRAN.R-project.org/package=data.table>
- R Core Team. (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. Retrieved from <http://ggplot2.org>
- Wickham, H., & Bryan, J. (2019). *Readxl: Read excel files*. Retrieved from <https://CRAN.R-project.org/package=readxl>
- Wickham, H., François, R., Henry, L., & Müller, K. (2019). *Dplyr: A grammar of data manipulation*. Retrieved from <https://CRAN.R-project.org/package=dplyr>




Figure 1. Methods




Figure 2. Average number of trials to criterion per group per condition.