HW 6

Max Horowitz-Gelb

$\mathbf{Q}\mathbf{1}$

Let our state space be [T, C]. Then we can think of one vehicle following another vehicle as a transition with transition matrix,

$$p_{i,j} = \begin{bmatrix} 1/4 & 3/4 \\ 1/5 & 4/5 \end{bmatrix}$$

We can then solve for the stationary distribution of this transition matrix simply since it is a 2x2 matrix.

$$\pi_T = \frac{1/5}{1/5 + 3/4} = 0.2105, \pi_C = \frac{3/4}{1/5 + 3/4} = 0.7895$$

Then since $p_{i,j} \neq 0$ for all i, j, then clearly p is irreducible and aperiodic. Then since our state space is finite and we have a stationary distribution π , then theorem 1.19 says the probability of being a truck converges to $\pi_T = 0.2105$.

$\mathbf{Q2}$

For this we simply solve for the set of functions

$$\pi_1 - (0.86\pi_1 + 0.05\pi_2 + 0.03\pi_3) = 0$$

$$\pi_2 - (0.08\pi_1 + 0.88\pi_2 + 0.05\pi_3) = 0$$

$$\pi_1 + \pi_2 + \pi_3 = 1$$

Then,

$$\pi = \begin{bmatrix} 0.014 & -0.08 & 1 \\ -0.05 & 0.12 & 1 \\ -0.04 & -0.05 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 0.2155477, 0.33215548, 0.45229682 \end{bmatrix}$$

Then since p is finite and $p_{i,j} \neq 1$ for all i,j then p is irreducible and is aperiodic, we may again apply theorem 1.19 to say that in the long run, $\pi_1 = 0.2155477$ fraction of the population will reside in cities, $\pi_2 = 0.33215548$ fraction of people will reside in suburbs and $\pi_3 = 0.45229682$ fraction of people will reside in rural areas.

Q3

a.

Using the same strategy as Q2 we first solve for a stationary distribution to get

$$\pi = [0.36363636, 0.35664336, 0.27972028]$$

Then since p is finite and $p_{i,j} \neq 0$ for all i,j then we again can apply 1.19 to say,

$$\lim_{n \to \infty} p^n(1,2) = 0.35664336$$

b.

Then since p is irreducible and closed, then it is recurrent, and we may apply theorem 1.21 which says,

$$\frac{N_n(2)}{n} \to \frac{1}{E_2 T_2}$$

and as shown in a.,

$$\frac{N_n(2)}{n} \to \pi_2$$

so,

$$\pi_2 = \frac{1}{E_2 T_2}$$

and

$$E_2T_2 - 1 = 1/\pi_2 - 1 = 1.8039$$

c.

$$E[\text{walking distance}] = \sum_n p(n) * (0.3 + 0.1n)$$

$$= \sum_n \pi_n * (0.3 + 0.1n) = 0.4916$$

$\mathbf{Q4}$

i

Our state space is [1,2,2'] where the number indicates the number of light bulbs working and ' indicates we just replaced the light bulbs. Then our transition matrix is

$$\begin{bmatrix} 0.95 & 0 & 0.05 \\ 0.02 & 0.98 & 0 \\ 0.02 & 0.98 & 0 \end{bmatrix}$$

Solving for the stationary distribution we get,

$$\pi = [0.2857, 0.7, 0.0143]$$

Then since p is finite and nonzero for all i, j, then it is irreducible and theorem 1.19 says that the long run fraction of time one light bulb is working is $\pi_1 = 0.2857$.

ii

Then we can use theorem 1.21 to say that

$$E_2T_{2'}-1=1/\pi_{2'}-1=69$$

So the expected number of days between light bulb is 69 days.

Q_5

First since states $\{1...M\}$ are closed, then for any $x \in \{1...M\}$.

$$\sum_{m} p(x,m) = \sum_{m} f((x-m) \mod M) = \sum_{n=0}^{M-1} f(n) = 1$$

Then for any $x \in \{1...M\}$,

$$\sum_{m} p(m, x) = \sum_{m} f((m - x) \mod M) = \sum_{n=0}^{M-1} f(n) = 1$$

therefore the THMC is doubly stochastic implying at has a uniform stationary distribution

$$\pi = [1/M...1/M]$$

Clearly our state space is finite. Then since p(1,2)>0 and p(2,4)>0 then for all $x, p(x, (x+1) \mod M), p(x, (x-1) \mod M), p(x, (x+2) \mod M), p(x, (x-2) \mod M)$ are all positive. So for any integers a, b such that $a+2b=M, p^{a+b}(x,x)\neq 0$. For any $M\geq 4$ the set of $\{a+b\}$ has a gcd of 1. Therefore our THMC is aperiodic and by theorem 1.19

$$\lim_{n\to\infty} p^n(1,1) = \pi_1 = 1/M$$

Then using theorem 1.23

$$\lim_{n\to\infty} \sum_{m=1}^{n} X_m^2 = \sum_{i=1}^{M} \pi_i * i^2 = 1/M * \sum_{i=1}^{M} i^2$$