

CS761 Spring 2015 Homework 1

Assigned Jan. 18, due Jan. 20 before class

Instructions:

- Homeworks are to be done individually.
- Typeset your homework in latex using this file as template (e.g. use pdf_lat_ex). Show your derivations.
- Hand in the compiled pdf (not the latex file) online. Instructions will be provided. We do not accept hand-written homeworks.
- Homework will no longer be accepted once the lecture starts.
- Fill in your name and email below.

Name: Max Horowitz-Gelb
Email: horowitzgelb@wisc.edu

Each question has 5 points if answered correctly. Wrong answers or blank answers 0 points. However, an explicit “I don’t know” answer is worth 1 point.

1. Define inner product on real functions

$$\langle g, h \rangle = \int_0^1 g(x)h(x) dx.$$

Orthogonalize the basis $1, x, x^2$.

$$1, x - \frac{1}{2}x^2 - x + \frac{1}{6}$$

2. Here is a little fact: Let X be a random variable with finite mean μ and finite variance σ^2 . Then $\forall t > 0$ we have

$$P(|X - \mu| \geq t) \leq \frac{\sigma^2}{t^2}.$$

Now let X_1, X_2, \dots be a sequence of independent and identically distributed random variables with mean μ and variance σ^2 . Let $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$. Use the little fact to prove

$$\forall \epsilon > 0, \forall \delta > 0, \exists N > 0, \forall n \geq N, P(|\bar{X}_n - \mu| \leq \epsilon) \geq 1 - \delta.$$

Proof

If X_1, \dots, X_n all have variance σ^2 and mean μ , then \bar{X}_n has variance σ^2/n and mean μ . Therefore by our little fact,

$$P(|\bar{X}_n - \mu| \geq t) \leq \frac{\sigma^2}{nt^2}$$

Now let $\epsilon > 0, \delta > 0$ be given. Then set $N = \lceil \frac{\sigma^2}{\delta \epsilon^2} \rceil$

Then for any $n \geq N$

$$P(|\bar{X}_n - \mu| \geq \epsilon) \leq \frac{\sigma^2}{n\epsilon^2} \leq \frac{\sigma^2}{N\epsilon^2} \leq \delta$$

Then since $P(|\bar{X}_n - \mu| \geq \epsilon)$, then

$$P(|\bar{X}_n - \mu| \leq \epsilon) \geq 1 - \delta$$