

Method C v2 (gamma, tau) grid: structural-deficit map across the parameter plane


Fixed-Point Factory

June 2026

Setup

10×10 grid: $\gamma \in \{0.1, 0.3, 1, 3, 10, 30, 100, 300, 1000, 3000\}$, $\tau \in \{0.001, 0.005, 0.01, 0.05, 0.1, 0.3, 0.5, 0.7, 0.85, 1.0\}$.
100 cells, Method C v2 additive Bayesian-symmetric monotone ansatz fit against the Lin-CDF R4 operator. 2 min total. Chain warm-start across τ within each γ column.


1 Structural-deficit residual $|F|_\infty$



This is the residual after fitting the best 5-parameter additive monotone ansatz. It measures **how much the equilibrium needs cross-signal joint structure beyond the additive form** (= the


true "deficit"). Smooth, monotone in τ , weakly increasing in γ above some threshold.

2 Slope α^*




Monotone in both τ and γ . Same qualitative shape as in the R4 sweep but with no kernel-band bias.

3 P range



4 Cross-sections



Left: slope vs τ at each γ — monotone, asymptote near $\gamma \geq 30$.

Right: residual vs τ at each γ — the structural non-additivity. Grows smoothly with τ ; relatively γ -independent at high γ , smaller at low γ .

5 Reproducibility

projects/REZN/solver_code/highprec_dd_qd/dd_k3_method_c_v2.py (operator), dd_k3_method_c_v2_grid.py (grid driver), dd_k3_method_c_v2_grid_figs.py (figures), dd_k3_mu_analytic.py (analytic 1D lookup).
 Saved coefficients: projects/REZN/solved.fixed.points/dd_k3_overnight/method_c_v2_grid/coeffs/.