

Basic Graph Theory

KNU Math 420

Classnotes

These notes are for a undergrad course in Graph theory based on Douglas West's 'Introduction to Graph Theory', which is referred to as 'the text'. Students should have the text. Test problems will be based on the problems in the notes. Not only are the lists of problems at the end of each section problem from the notes, then problems written out in the notes are usually asking you to fill in details from the text. Some, but not all, of these problems will be answered in class.

Students should be familiar with the set theory and logic seen in Discrete Math.

1 Fundamental Concepts


1.1 What is a graph?

A graph G consists of a non-empty finite set $V = V(G)$ of elements called *vertices* and a relation $E = E(G)$ on V , called the *edge* relation, which relate pairs of (not necessarily distinct) vertices. When edge relation is irreflexive and symmetric, the graph is called *simple*. Usually our graphs are simple. If $\{u, v\} \in E$, we say they u and v are *adjacent*, or that they are *neighbours*, and that they are the *end points of the edge* $\{u, v\}$. If $\{v, v\}$ is an edge, then it is a *loop*. Simple graphs don't have loops. If E is not symmetric, we call G a *digraph*.

For example

$$G = (\{1, 2, 3, 4\}, \{\{1, 2\}, \{2, 3\}, \{3, 4\}, \{4, 1\}, \{1, 3\}\})$$

is a graph. It has four vertices and five edges. As this mess of brackets gets troublesome to write, we drop them when it doesn't confuse things and write an edge $\{x, y\}$ as xy . The graph G can be represented pictorially as



Matrices and Isomorphism

To describe a particular graph we have to describe its vertices and its edges. The names of the vertices are usually not important, and so we usually only talk of a graph up to isomorphism.

Definition 1.1.1. An *isomorphism* is a bijective map $f : V(G) \rightarrow V(H)$ such that

$$uv \in E(G) \iff f(u)f(v) \in E(H).$$

If there is an isomorphism f from G to H , we say G and H are *isomorphic*, and write $G \cong H$.

The relation of being isomorphic is clearly an equivalence relation on the set of all graphs.

As we usually consider graphs to be the same if they are isomorphic. Mostly, graph theory is the study of *graph properties*—properties of a graph that are invariant under isomorphism. The number of vertices, and the size of the smallest cycle in the graph are such properties.

Occasionally, one does talk of *labelled graphs*, distinguishing graphs in the same isomorphism class, as it can help make things easier to count.

Problem 1.1.1. How many graphs are there on 4 vertices? How many non-isomorphic (unlabelled) graphs?

Obviously we will not always be able to describe a graph with a picture. To put a graph into a computer, we often represent it as a matrix.

For a simple graph G on the vertex set $V = \{v_1, \dots, v_n\}$ and having edgeset $E = \{e_1, \dots, e_m\}$ the *adjacency matrix* $A(G)$ is the symmetric $n \times n$ matrix $A = [a_{ij}]$ where

$$a_{ij} = \begin{cases} 1 & \text{if } v_i \sim v_j \\ 0 & \text{otherwise} . \end{cases}$$

The incidence matrix $M(G)$ is the $m \times n$ matrice $M = [m_{ij}]$ such that

$$m_{ij} = \begin{cases} 1 & \text{if } v_j \in e_i \\ 0 & \text{otherwise} . \end{cases}$$

Problem 1.1.2. Where G is the four vertex graph drawn above, give $A(G)$ and $M(G)$.

Representing graphs this way often makes it easier to decide if they are isomorphic, but deciding if two graphs are isomorphic is a computationally hard problem.

Graphs as Models

See Section 1.1 of the text for several typical examples of things that are modelled by graphs. The following are typical questions we might ask about a graph. Try to answer them for the graph P shown in Figure 1. The definitions you need to understand these questions come later, but they are probably what you think they are.

- How many colours do we need, when colouring the vertices of P to be sure that adjacent vertices get different colours.
- Can you draw P without intersecting edges?
- How many edges (vertices) must you remove to *disconnect* P ?





Figure 1: The Petersen Graph P

- For any two pairs of vertices u, u' and v, v' can you find a *path* between u and u' , and another between v and v' , such that the two are disjoint?
- What is the smallest *cycle* in P ?
- How many 5-cycles are there in P ?
- Can we partition the edges of P into disjoint 5-cycles?
- Can we find a *walk* in P that visits every vertex (edge) exactly once?



- Is P isomorphic to this graph?

Some Common Graphs

We generally now consider graphs to be the same if they are isomorphic, and take blatant liberties with the names of their vertices. As such, we define some common graphs, these names apply to any graph that is isomorphic.

- The *complete graph*, or *clique* on n vertices, K_n , has $V(K_n) = \{v_1, v_2, \dots, v_n\}$ and $E(K_n) = \{v_i v_j \mid 1 \leq i < j \leq n\}$.
- The *n -cycle*, C_n , has $V(C_n) = \{v_1, v_2, \dots, v_n\}$ and $E(C_n) = \{v_i v_{i+1} \mid i = 1, \dots, n-1\} \cup \{v_n v_1\}$.
- The *n -path*, P_n , has $V(P_n) = \{v_1, \dots, v_n\}$ and $E(P_n) = \{v_i v_{i+1} \mid i = 1, \dots, n-1\}$. (Many books use P_n to mean the path with n edges. This is actually the more common usage these days. Be careful of this when you are looking at other sources.)
- The *complete bipartite graph*, or *biclique* $K_{m,n}$, has $V(K_{m,n}) = \{a_1, \dots, a_m\} \cup \{b_1, \dots, b_n\}$ and $E(K_{m,n}) = \{a_i b_j \mid i \in [m], j \in [n]\}$.

- The n -cube, Q_n , $V(Q_n)$ is the set of binary strings of length n , vertices u and v are adjacent if they differ in exactly one coordinate.

Subgraphs

Given a graph G , another graph G' is a *subgraph* of G , written $G' \leq G$ if $V(G') \subset V(G)$ and $E(G') \subset E(G)$.

A subgraph G' of G is a *proper* subgraph of G if $E(G')$ is a proper subset of $E(G)$. It is *spanning* if $V(G') = V(G)$, or *induced* if $E(G') = \{uv \in E(G) | u, v \in V(G')\}$. For a subset $T \subset V(G)$ the subgraph $G[T]$ of G *induced* on T is the induced subgraph of G with vertex set T ; we get it from G by removing all vertices not in T .

Problem 1.1.3. How many non-labelled subgraphs are there of K_4 ? How many are proper? ...induced? ...spanning? ...spanning, induced and proper?

A subgraph of G isomorphic to a path (cycle, clique) is called a path (cycle, clique) of G . The *complement* $\overline{G}$ of a (simple) graph G is the graph with the same set of vertices and edgeset

$$E(\overline{G}) = \{uv \mid u \neq v \in V(G)\}.$$

An independent set of G is a clique of $\overline{G}$.

A graph G is *bipartite* if it is a subgraph of a biclique.

Problem 1.1.4. Show that the following properties are equivalent for a graph G .

- G is bipartite.
- G there is a partition of $V(B)$ into sets A and B such that every edge has one endpoint in A and one in B .
- G contains no odd cycles.

(Showing that every non-bipartite graph has an odd cycle is non-trivial.)

Problem 1.1.5. Show that a subgraph G' of G is induced if and only if $\overline{G'}$ is a subgraph of $\overline{G}$.

Solution

Let $G' \subset G$ be induced. Clearly $V(\overline{G'}) \subset V(\overline{G})$, so we have to show that $E(\overline{G'}) \subset E(\overline{G})$. Let $e \in E(\overline{G'})$ then $e \notin E(G')$ so $e \notin E(G)$. So $e \in E(\overline{G})$.

On the other hand, let $\overline{G'} \leq \overline{G}$. We want to show that G' is induced. For $u, v \in V(G')$, with $uv \in E(G)$ we want to show that $uv \in E(G')$. Well $uv \in E(G)$ implies that $uv \notin E(\overline{G})$ so not in $E(\overline{G'})$, so because $u, v \in V(G')$, $uv \in E(G')$.

Decompositions and Special graphs

A graph is *self-complementary* if $G \cong \overline{G}$. Can you think of a self-complementary graph? How many edges must it have? A self complementary subgraph H of K_n yields a decomposition of K_n into two copies of H .

A family of subgraphs of G is a *decomposition* of G if the edgesets of the subgraphs partition the edgeset of G .

Problem 1.1.6. Show that K_7 has a decomposition into copies of K_3 . Does K_6 ?

The graph P from Figure 1 is called the Petersen graph. It comes up a lot. It has a neat construction that leads to its symmetry. Its vertex set can be viewed as the family of two-element subsets of the set $[5] = \{1, 2, 3, 4, 5\}$. Two vertices are adjacent if the subsets are disjoint.

This construction is useful in proving things about the Petersen graph. It has no triangles because this would require three mutually disjoint two element subsets in $[5]$; which is impossible. It has no 4-cycles because any two non-adjacent vertices must, as sets, share one element, and so there is exactly one vertex adjacent to both of them. Thus the smallest cycle in P is a C_5 . Can you find a 5 cycle? Can you show that any edge of P is in a 5-cycle?

The *girth* of a graph G is the length of its shortest cycle. An automorphism of a graph G is an isomorphism $f : G \rightarrow G$. A graph is *vertex transitive* if for every pair u, v of vertices, there is an automorphism f of G such that $f(u) = v$.

Problem 1.1.7. Show that the Petersen graph is vertex transitive.

The *degree* $d(v) = d_G(v)$ of a vertex v of a graph G is the number of edges of G that contain v (loops containing v are counted twice).

Problems from the Text

Section 1.1: 1 - 10, 12, 13, 14, 18, 23, 25, 41

1.2 Paths, cycles, and trails

Let x and y be vertices in a graph G . An *xy-walk* in G is a finite sequence

$$x = x_1 x_2 \dots x_{n-1} x_n = y$$

of vertices such that $x_{i-1} \sim x_i$ for $i = 1, \dots, n$. The *length* of the walk is $n - 1$, (we count edges, not vertices). If the edges $x_i x_{i+1}$ of the walk are distinct, it is a *trail*, and if the vertices are distinct, it is a *path*. It is closed if $x_1 = x_n$.

Theorem 1.2.1. Any *xy-trail* in a graph has an *xy-path* as a subgraph.

Intuitively, we just start with an xy -trail and 'chop' out the repeated bits until it is a path. A lot of times it is very hard to write the intuitive proof rigorously. The text uses an induction on the length of the trail. We do the same proof, using the suggestion in exercise 13b.

Proof. Let $T = x_0x_1 \dots x_d$ be the shortest xy -trail in G ; one exists by assumption.

We claim that T is an xy -path. If not, there are some $0 \leq i < j \leq d$ such that $x_i = x_j$. But then

$$x_0, x_1, \dots, x_{i-1}, x_i, x_{j+1}, \dots, x_d$$

is a shorter xy -trail T , which contradicts our choice of T . $\square$

Problem 1.2.1. Prove the Theorem 1.2.1 using induction.

Problem 1.2.2. Prove the following. Any odd circuit in a graph has an odd cycle as a subgraph.

A graph G is *connected* if for every pair u, v of vertices there is a uv -path in G . A maximal connected subgraph of G is a *component*. A component is *trivial* if it contains only one vertex, the vertex is an *isolated* vertex. The *distance* $d(u, v)$ between two vertices of G is the length of the shortest path between them; it is infinite if they are in different components. A *cut-edge* or *cut-vertex* of a graph is a edge or vertex whose removal increases the number of components of the graph.

Theorem 1.2.2. *An edge is a cut-edge if and only if it is not in any cycles.*

Proof. See text. $\square$

1.2.1 Bipartite Graphs

Recall that a graph G is bipartite if there is a partition $V(G) = A \cup B$ of its vertex set such that all edges are between A and B . We call A and B its bipartition. It can have more than one, but only if it is disconnected.

We asked you to show this earlier.


Theorem 1.2.3 (König (1936)). *A graph is bipartite if and only if it has no odd cycles.*

Proof. See text. $\square$

1.2.2 Eulerian Circuits

It is often told that the first problem of graph theory was the problem of the Bridges of Königsberg. In the fictional city of Königsberg (Problem: prove or disprove my unpopular claim that Königsberg is fictional.) there were seven bridges. The dandies of the city, on slow Sunday, would walk about the town, playing a riddish game. Starting wherever they pleased, they were to walk around, crossing every bridge, without crossing any bridge more than once. This game went on for centuries, and several liars claimed to have done it. Fermat once wrote in the margin of a book that it was simple enough. Until eventually, Euler proved it was impossible, and stabbed Fermat in a dual. (Prove that any of this is true.)

The city of Königsburg consisted of four dots, connected by seven bridges. It looked a lot like this:



A closed trail is called a circuit. A circuit of a graph G is an *Eulerian circuit* if it contains all edges and vertices of G . A graph is *Eulerian* if it contains an Eulerian circuit. A graph is *even* if all vertices have even degree.

The following is known as Euler's Theorem.

Theorem 1.2.4. *A graph is Eulerian if and only if it is even and connected.*

Proof. That an Eulerian graph must be connected is clear. That it must be even is almost as clear— let $C = v_1v_2 \dots v_N$ be an Eulerian circuit. For any vertex v , count the number d of times that v occurs in $v_1v_2 \dots v_{N-1}$. For each occurrence $v = v_i$ is in two edges in C : $v_{i-1}v$ and vv_{i+1} . So v is in $2d$ edges in C . But C contains each edge of G once, so v is in $2d$ edges in G .

Now, in the other direction, assume that G is a connected even graph. Our proof is by induction on the number m of edges. When $m = 0$, the fact that G is connected means that it has only one vertex, and this vertex is a closed trail containing all edge.

Assume then that $m > 0$. As it is even and non-empty it must contain a circuit C — indeed pick a vertex v_0 with an edge, start walking, whenever we enter a vertex other than v_0 , we have visited an odd number of that vertices edges, so one remain to leave it by. We can keep walking then, until we get back to v_0 .

Now, the graph $G' = G - C$ need not be connected, but each component is even, so contains a Eulerian circuit of the component by induction. Moreover

C visits each component, as G is connected. Traversing C , the first time we encounter a vertex of a component X , we pause our traversal of C , traverse the Eulerian circuit of X which brings us back to where we entered X , and then continue our traversal of C .

This yields an Eulerian circuit of G , as needed. $\square$

Problem 1.2.3. Using Euler's Theorem, show that in a connected graph G with $2k$ vertices of odd degree, there is a decomposition of G into k trails, and no fewer.

Problems from the Text

Section 1.2: 1-12, 14, 16, 17, 27, 29, 43

1.3 Vertex Degrees and Counting

The *size* of a graph $G = (V, E)$ is the number $e(G) = |E|$ of edges it has, while the *order* is the number $n(G) = |V|$ of vertices it has. Quite often we use n on its own for the order of the graph we are talking about, and m for its size. (We can't use e , because we usually use that for an edge of G .) Or we will specify n by calling G an n -vertex graph.

For a vertex v of G , the *neighbourhood* $N(v) = N_G(v)$ of v in G is the set of edges in $E(G)$ that contain v . So for a simple graph, the degree of a vertex is $d(v) = |N(v)|$. The *min-degree* $\delta(G)$ of G is the minimum of the degrees of its vertices, and the *max-degree* $\Delta(G)$ is the maximum. If all vertices of G have degree k , then G is *k -regular*.

While many of the 'common graphs' we have seen are regular, it is a very rare property for a graph to be regular.

Problem 1.3.1. What is the size of a k -regular graph of order n ?

A vertex transitive graph is, of course, regular. We showed earlier that the Petersen graph is vertex transitive, so is clearly 3-regular.

This example, from Section 1.3 of the text, shows how nice it can be to have vertex transitivity. A *claw* is the graph $K_{1,3}$.

Example 1.3.1. We show that the Petersen graph P has ten 6-cycles, by showing that there is a 1-to-1 bijection between 6-cycles and claws. As each of the 10 vertices of P is the center of exactly one claw, P has ten claws, so this is enough.

Pick the claw in P with center v , and leaves a, b, c . Each of the leaves has two other edge, but as the leaves are distance 2 apart, none of their edges other than the ones to v , can share an endpoint, as P has no 3-cycles. So when we

remove a claw, the six edges from the claw to the rest of P , all go to different vertices of P . Thus what is left is a 2-regular graph of order 6. As P has no triangles, this can only be a 6-cycle.

Not all regular graphs are as nice as the Petersen graph.

Problem 1.3.2. Find a regular graph that is not vertex transitive.

But regularity can be still be nice. There is a difficult open conjecture. The *deck* of a graph G is the multiset of *unlabelled* vertex deleted subgraphs

$$\{G \setminus v_i \mid v_i \in V(G)\}.$$

The following conjecture posits that no two non-isomorphic graphs with at least 3 vertices can have the same deck.

Conjecture 1.3.2 (Graph Reconstruction Conjecture (Kelly and Ulam 1942)). *A graph of order at least 3 is uniquely determined by its deck.*

Problem 1.3.3. Why do we need three vertices to conjecture this?

This conjecture is almost 100 years old, and we cannot prove it. But we do know that it is true for certain classes of graphs. In the problems you will show the following; it is quite easy— maybe you see already why it is true.

Fact 1.3.3. *The Graph Reconstruction Conjecture is true for regular graphs.*

Extremal graphs

Problems in extremal graph theory ask such things as how big or small a graph can be and have a certain property.

Some simple such questions are:

Problem 1.3.4. What is the maximum number of edges an n -vertex graph can have? What is the minimum?

Maybe this problem was too simple. This one is only marginally harder.

Problem 1.3.5. What is the maximum number of edges in a **disconnected** graph on n vertices?

Still too simple? Maybe, but it introduces some ideas. To answer this question, you found an extremal graph, a disconnected graph on n vertices with _____ edges. This is called the *extremal graph* for the problem. Then you had to prove that there was no graph on n vertices with more edges.

There are many extremal graphs for the following question.

Problem 1.3.6. What is the minimum number of edges in a **connected** n -vertex graph?

The following problem is maybe not an extremal problem, but has the flavour of one, and is a useful tool for many extremal problems.

Theorem 1.3.4. *Every simple graph has a bipartite subgraph containing at least half of its edges.*

Proof. The proof is by induction on the number, n , of vertices of the graph G . The statement is clearly true when $n = 1$, so assume that $n \geq 2$ and that the result holds on all graphs of order less than n . Let v be a vertex of G and let $G' = G \setminus v$. By the induction hypothesis, there is a partition $V(G') = A' \cup B'$ such that at least half of the edges of G' are between A' and B' . We may assume that at least half of the edges of G that contain v are to vertices in B' , and so setting $A = A' \cup \{v\}$ we get a partition $V(G) = A \cup B'$ such that at least half of the edges of G are between A and B' . $\square$

A graph G is H -free if it contains no *induced* copy of H . Mantel proved one of the first results of extremal graph theory. We will later see a generalisation by Turan.

Problem 1.3.7. How many edges can a graph on n vertices have, and have no triangles?

There is a pretty good family of candidates for the extremal graph—graphs with lots of edges and no triangles: the bipartite graph $K_{a, (n-a)}$ has no triangles and has $a(n-a)$ edges. You can use calculus, or an argument as we used in the above theorem, to do the following.

Problem 1.3.8. Show that $a(n-a)$ is maximized when $a = \lfloor n/2 \rfloor$, and so show that there is a bipartite graph of order n with $\lfloor n/2 \rfloor \lceil n/2 \rceil = \lfloor n^2/4 \rfloor$ edges.

So a graph can have $\lfloor n^2/4 \rfloor$ edges and have no triangles. Can it have more?

Definition 1.3.5. Let $T(n)$ be the maximum number of edges in a triangle free n -vertex graph.

The above example showed that $T(n) \geq \lfloor n^2/4 \rfloor$.

Theorem 1.3.6 (Mantel (1907)). *For all $n \geq 1$ we have $T(n) \leq \lfloor n^2/4 \rfloor$.*

Proof. As $T(n)$ is an integer, it is enough to show that $T(n) \leq n^2/4$. The base case is $n = 1$ and 2 for which it is clearly true. Assume the theorem holds for all graphs on at least n vertices, and let $G = (V, E)$ be a triangle free graph on $n+2$ vertices.

Choose an edge $e = \{x, y\}$ from E , and let $G' = (V', E')$ be the subgraph of G induced by $V' = V \setminus \{x, y\}$. Let E_x be the set of edges from x to a vertex of V' , and E_y be the set of edges from y to a vertex of V' .

As $E = E' + E_x + E_y + \{e\}$, and from the fact that G is triangle-free we get that $|E_x| + |E_y| \leq n$. Thus from the induction hypothesis we get that that

$$|E| \leq \frac{n^2}{4} + n + 1 = \frac{(n+2)^2}{4}.$$


□

Graphic sequences

Definition 1.3.7. The *degree sequence* of a graph G is the multiset of the degrees of the vertices of G . The minimum degree, $\delta(G)$, and maximum degree, $\Delta(G)$, of a graph G are the minimum and maximum degrees respectively in its degree sequence.

Problem 1.3.9. Show that every graph contains a path of length at least $\delta(G)$.

Clearly, degree sequence, minimum degree, and maximum degrees are graph properties. So the following two graphs, which have the same number of vertices, and edges are not isomorphic as they have different degree sequences: $(4, 2, 2, 2, 2)$ and $(3, 3, 2, 2, 2)$ respectively, (and in fact different maximum degrees.)



Here is a simple observation about degree sequence:

Theorem 1.3.8. If $(d_1, d_2, \dots, d_n)$ is the degree sequence of a graph, then the degree sum: $\sum_{i=1}^n d_i$ is equal to $2|E(G)|$.

Proof. Each edge of the graph contributes 2 to the degree sum, one for each endpoint. □

Corollary 1.3.9. *In any graph G , there is an even number of vertices of odd degree.*

Problem 1.3.10. What is the degree sequence of the cube Q_n ?

Clearly isomorphic graphs have the same degree sequence. But the converse is clearly not true.

Problem 1.3.11. Find graphs with the same degree sequence that are not isomorphic.

A sequence $D = (d_1, d_2, \dots, d_n)$ non-increasing integers is *graphic* if there is a graph G on the vertices $v_1, \dots, v_n$ such that $d(v_i) = d_i$ for all $i = 1, \dots, n$. Such a graph G is called a *realisation* of D . We saw above that one necessary condition for a sequence to be graphic is that it sums to an even number. But this is not sufficient: $(4, 2, 0)$ is not graphic.

There is a recursive method to decide if a given sequence is graphic.

Lemma 1.3.10. (Havel-Hakimi) *If a sequence $D = (d_1, d_2, \dots, d_n)$ is graphic, then there is a graph with degree sequence D such that the maximum degree vertex v_1 is adjacent to the d_1 vertices of next highest degree.*

Before we prove this. Let's see how it allows us to decide if a sequence is graphic. If D is graphic, then it has a realisation as in the lemma. If we remove the edges from v_1 then the graph we get has degree sequence

$$D' = (0, d_2, d_3 - 1, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_{n-1}, d_n)$$

which we get by changing d_1 to 0 and reducing the next d_n values by one.

On the other hand, if the sequence D' is graphic, then so is D .

This gives us a recursive algorithm for deciding if a degree sequence D is graphic: keep reducing the sequence as above until all the degrees are reduced to 0 or less. This clearly terminates, as the sum of degrees is reduced at each step.

If we end with a sequence of zeros, D was graphic. If we get any negative numbers, it was not graphic.

(Sometimes this is done reordering the sequences at each step, but it is useful to keep them unordered, as it is then easier to reconstruct the graph if the sequence is graphic.)

Problem 1.3.12. Use this algorithm to decide if $(3, 3, 2, 2, 2)$ is graphic.

Now let's prove the lemma.

Proof. Let $D = (d_1, d_2, \dots, d_n)$ be a graphic sequence. Given a realisation G of D , let $T = \{v_2, \dots, v_{d_1+1}\}$, and let $t_G = |N(v_1) \cap T|$. As D is graphic, it

has some realisation. Let G be on that maximises t_G . If $t_G = \Delta$ then we are done, so we may assume, towards contradiction, that $t_G < \Delta$. We find a new realisation G' with $t_{G'} > t_G$, so the lemma follows by contradiction.

As $N(v_1) \neq T$, there is $x \in T$ with $v_1 \not\sim x$ and $y \notin T$ with $v_1 \sim y$. Furthermore, since $x \in T$ and $y \notin T \cup \{v_1\}$, $d(x) \geq d(y)$. So there is some vertex s with $s \sim x$ and $s \not\sim y$.

Removing the edges v_1y and xs from G and adding v_1x and ys we get a graph G' with the same degrees on every vertex, (so another realisation of D), but with $t_{G'} = |N_{G'}(v_1) \cap T| = |(N_G(v_1) \cap T) \cup \{x\}| = t_G + 1$, as needed for the contradiction. $\square$

Problem 1.3.13. Decide, using Havel-Hakimi, whether or not the sequence $(5, 5, 5, 3, 2, 2, 1, 1)$ is graphic.

Our proof above used what we might call a *2-switch*. Let a graph G have vertices $a, b, c, d \in V(G)$ such that

$$a \sim b \not\sim c \sim d \not\sim a.$$

The graph we get from G by removing the edges ab and cd , and adding the edges bc and da , is a 2-switch of G . The process of adding and removing the edges is also called a *2-switch*.

Corollary 1.3.11 (Berge (1973)). *Let G and H be graphs with the same degree sequences. One can get from G to H by a sequence of 2-switches.*

Proof. By the above proof of Havel-Hakimi, we can switch both graphs to the same graph. $\square$

Problems from the Text

1.3: 1-8, 9, 10, 12, 13, 14, 21, 28, 30, 36, 37, 38, 39, 46, 62

We skip Section 1.4 on digraphs, I do it in discrete math, and don't want too much overlap.

2 Trees and Distance

2.1 Basic Properties

A *forest*, or *acyclic graph*, is a graph with no cycles as subgraphs. A *tree* is a connected forest. A *leaf* is a vertex of degree 1. A *spanning tree* of a graph G is a spanning subgraph which is a tree.

A lot of the results we give about trees might seem obvious. This, then is a good place to practice proper proof. For any of these results, I can draw a picture, and wave my hands, and convince you that they are true. But a picture is usually not a proof. And proving what seems clear is often quite difficult.

Problem 2.1.1. Show that a graph is connected if and only if it has a spanning tree.

Hand-waving proof: The spanning tree connects the graph, so a graph with a spanning tree is clearly connected. On the other hand, let G be connected and H be a minimal connected subgraph. We are done unless H has a cycle, but if it has a cycle we can remove some edge from the cycle without disconnecting H , contradicting the choice of H .

Everything I say there is true. And you believe it. But is it enough. Should I justify some of my statements? What does it mean that 'the spanning tree connects the graph'? Why does removing an edge from the cycle not disconnect H ?

Problem 2.1.2. Show that a connected graph is a tree if and only if every edge is a cut-edge.

The following will be useful in inductions. This result requires the assumption, which we always make, that graphs are finite.

Lemma 2.1.1. *Any tree T of order at least 2 has at least two leaves, and removing a leaf from such a tree produces a smaller tree.*

Proof. Let T be a tree with at least two vertices. Let P be a maximal path in T . It is non-trivial because T has at least one edge. The endpoints of P are leaves, because if they have more than one neighbour in P , then we have a cycle, which is impossible in a tree, and if they have any neighbours not in P , then P was not maximal. Thus we have at least two leaves.

Let ℓ be a leaf and $T' = T - \ell$. T' is certainly acyclic, as T was. It is also connected, as ℓ , having degree 1 is not in the path in T between any two vertices u, v in $V(T) \setminus \{\ell\}$. $\square$

As trees are so basic, having several characterisations of them can be useful in proofs.

Theorem 2.1.2. Let $G = (V, E)$ be an n -vertex graph. Then the following are equivalent.

- i. G is connected and acyclic.
- ii. G is connected and has $n - 1$ edges.
- iii. G is acyclic and has $n - 1$ edges.
- iv. Any two vertices of G are connected by a unique path.

Proof. The text breaks the proof down to proving the following five implications.

$$i \Rightarrow \{ii, iii\} \quad ii \Rightarrow \{i, iii\} \quad iii \Rightarrow \{i, ii\} \quad i \Rightarrow iv \quad iv \Rightarrow i$$

Problem 2.1.3. Pick one implication and write a proof for it. (In the text, the proofs of the second and third implications use that the first is already proved; this is allowed.) Don't look at the text until you have complete your proof.

□

Proposition 2.1.3. Let T and T' be spanning trees of a connected graph G and let $e \in E(T) - E(T')$.

- There is an edge $e' \in E(T') - E(T)$ such that $T - e + e'$ is a spanning tree of G .
- There is an edge $e' \in E(T') - E(T)$ such that $T' + e - e'$ is a spanning tree of G .

Proof. i) As T is a tree, so e is a cut-edge, $T - e$ has two components. There is a path in T' between a vertex u in the first and a vertex v in the second, so there is an edge e' of T' in that path going between the components. Thus $T - e + e'$ is connected, and so by the edge count, is a spanning tree.

ii) The graph $T' + e$ has one cycle, and any edge $e' \neq e$ in this cycle can be removed without disconnecting it, so $T' + e - e'$ is connected, and so by the edge count is a spanning tree. □

This result has a cool application given at the end of the Section in the text.

Distance in trees and graphs

Recalling that the distance $d(u, v)$ between vertices u and v in a graph G is the length (number of edges) of the shortest uv -path in G , the *diameter* of G is

$$\text{diam } G = \max_{u, v \in V(G)} d(u, v).$$

The *eccentricity* of a vertex u is

$$\epsilon(u) = \max_{v \in V(G)} d(u, v),$$

and the *radius* of G is

$$\text{rad } G = \min_{u \in V(G)} \epsilon(u).$$

The *center* of G is the set of vertices of minimum eccentricity, or the subgraph of G induced by them.

Problem 2.1.4. Find the diameter and radius of K_n, P_n, C_n, Q_n and P .

Graphs with big radius tend to have few edges while graphs with small radius have many. One particular example of this is the following.

Problem 2.1.5. Show that if $\text{diam}(G) \geq 3$ then $\text{diam}(\overline{G}) \leq 3$.

Theorem 2.1.4 (Jordan (1869)). *The center of any tree is a vertex or an edge.*

Proof. See the book for the strong induction on n . □

Problems from the Text

2.1: 1-16, 18, 19, 32, 33, 38, 40, 51, 54