

# Example 2 - Scipy special functions and PyFVTool

September 1, 2025

## 1 Example 2.

First, take an initial look at this notebook. Then, use “Kernel-> Restart & Run All” to re-evaluate the entire notebook. Running this notebook is a good test for your Python installation.

This example will use Scipy to evaluate the analytic solution for a heat transfer problem, and PyFVTool to solve the same problem by the finite-volume method.

No more details are given here. Look up more theory in the book by Crank (“Mathematics of Diffusion”).

```
[1]: import numpy as np
from numpy import exp
from scipy.special import jn_zeros, j0, j1
import matplotlib.pyplot as plt
import pyfvtool as pf
print('PyFVTool version', pf.__version__)
```

PyFVTool version 0.4.1

### 1.1 Analytic solution.

See the book by Crank (“Mathematics of Diffusion”), page 78, section 5.3.

Equation (5.22) reads

$$\frac{C - C_1}{C_0 - C_1} = 1 - \frac{2}{a} \sum_{n=1}^{\infty} \frac{\exp(-D\alpha_n^2 t) J_0(r\alpha_n)}{\alpha_n J_1(a\alpha_n)}$$

Here we will evaluate and plot this equation.

```
[2]: def crank522(r, t, a, D):
    '''evaluate eqn 5.22 for a given r,t
    a : radius of cylinder
    D : diffusion coefficient

    the following global variables need to be set

    Nterm_crank : number of terms to be evaluated
    '''
```

```

global Nterm_crank

aalp = jn_zeros(0, Nterm_crank)
alpha = aalp/a

XJ0 = exp(-D * alpha**2 * t) * j0(r*alpha)
AJ1 = alpha * j1(aalp)

S = np.sum(XJ0/AJ1)

return 1.0 - (2.0/a) * S

```

```
[3]: # set world parameters
```

```

a_val = 2.9
D_val = 1.9
Nterm_crank = 30

```

```
[4]: # create radial axis for plotting analytic solution
```

```

r_an = np.linspace(0.,a_val,200)


```

```
[5]: # evaluate crank522 at different points in time, and plot
```

```

for t in [0.005, 0.01, 0.02, 0.03, 0.04, 0.06, 0.08, 0.1, 0.15,
          0.2, 0.3, 0.4, 0.6, 0.8]:
    c = np.array([crank522(rr, t, a_val, D_val) for rr in r_an])
    plt.plot(r_an, c)
plt.xlabel('rel. radial position')
plt.ylabel('rel. temperature')
plt.title('analytic solution')
plt.show()

```



## 1.2 Finite-volume solution with PyFVTool.

Define 1D cylindrical grid with a variable called ‘c’, initialized to an initial value of 0.0 everywhere.

```
[6]: Nr = 50
     Lr = a_val
     c_outer = 1.0 # (outer) boundary concentration
     c0 = 0.0
     deltat = 0.001
```

```
[7]: mesh = pf.CylindricalGrid1D(Nr, Lr)
```

```
[8]: c = pf.CellVariable(mesh, c0)
```

By default, the boundary conditions for a variable are of the ‘no flux’ (Neumann) type.

Here, we apply a different boundary condition: the outer (rightmost) boundary will be kept at 1.0 (Dirichlet boundary condition).

```
[9]: # switch the right (=outer) boundary to Dirichlet: fixed concentration
     c.BCs.right.a[:] = 0.0
```

```
c.BCs.right.b[:] = 1.0
c.BCs.right.c[:] = c_outer
```

When changing the BCs, it is necessary to update everything in the cell variable object by calling `apply_BCs()`

```
[10]: c.apply_BCs()
```

```
[11]: D = pf.CellVariable(mesh, D_val) # diffusion coefficient
      alfa = pf.CellVariable(mesh, 1.0) # transientterm coefficient
```


```
[12]: t = 0.0
```

Now, we solve the equation by taking time steps. We plot the solution at several specified time-points, together with the analytic solution.

```
[13]: r_an = np.linspace(0., a_val, 200) # axis for plotting analytic solution

      sample_i = [5, 10, 20, 30, 40, 60, 80, 100, 150, 200, 300, 400, 600, 800]

      for i in range(0, 1001):
          if i in sample_i:
              r, phi = c.plotprofile()
              plt.plot(r, phi)
              c_an = np.array([crank522(rr, t, a_val, D_val) for rr in r_an])
              plt.plot(r_an, c_an, 'k:')
              # calculate the value of D at the faces
              # as the harmonic mean of the values at the centers
              Dave = pf.harmonicMean(D)
              pf.solvePDE(c,
                          [ pf.transientTerm(c, deltat, alfa),
                            -pf.diffusionTerm(Dave)])
              t += deltat
      plt.xlabel('radial position $r$')
      plt.ylabel('rel. temperature $T$')
      plt.title('FVM (PyFVTool) solution. Dotted line = analytic');
```



1.3 End.

[ ]: