

Example 2 - Scipy special functions and Fipy

June 15, 2022

1 Example 2.

First, take an initial look at this notebook. Then, use “Kernel-> Restart & Run All” to re-evaluate the entire notebook. Running this notebook is a good test for your Python installation.

This example will use Scipy to evaluate the analytic solution for a heat transfer problem, and Fipy to solve the same problem by the finite-volume method.

No more details are given here. They will be given in the *Thermodynamique & Phénomènes de transport* course.

```
[1]: # Use the following for JupyterLite
      # import piplite
      # await piplite.install("fipy")
```

```
[2]: import numpy as np
      from numpy import exp
      from scipy.special import jn_zeros, j0, j1
      import matplotlib.pyplot as plt
      import fipy as fp
      print('Fipy version',fp.__version__)
```

Fipy version 3.4.2.1

1.1 Analytic solution.

See the book by Crank (“Mathematics of Diffusion”), page 78, section 5.3.

Equation (5.22) reads

$$\frac{C - C_1}{C_0 - C_1} = 1 - \frac{2}{a} \sum_{n=1}^{\infty} \frac{\exp(-D\alpha_n^2 t) J_0(r\alpha_n)}{\alpha_n J_1(a\alpha_n)}$$

Here we will evaluate and plot this equation.

```
[3]: def crank522(r, t):
      '''evaluate eqn 5.22 for a given r,t

      the following global variables need to be set
      a : radius of cylinder
```

```

D : diffusion coefficient
Nterm : number of terms to be evaluated
'''
global a
global D
global Nterm

aalp = jn_zeros(0,Nterm)
alpha = aalp/a

XJ0 = exp(-D * alpha**2 * t) * j0(r*alpha)
AJ1 = alpha * j1(aalp)

S = np.sum(XJ0/AJ1)

return 1.0 - (2.0/a) * S

```

```

[4]: # set world parameters
a = 1.0
D = 1.0
Nterm = 20

```

```


[5]: # create radial axis
r = np.linspace(0.,1.,200)

```

```

[6]: # evaluate crank522 at different points in time, and plot
for t in [0.005, 0.01, 0.02, 0.03, 0.04, 0.06, 0.08, 0.1, 0.15,
          0.2, 0.3, 0.4, 0.6, 0.8]:
    c = np.array([crank522(rr,t) for rr in r])
    plt.plot(r,c)
plt.xlabel('rel. radial position')
plt.ylabel('rel. temperature')
plt.title('analytic solution')
plt.show()

```



1.2 Finite-volume solution with Fipy.

Define 1D cylindrical grid with a variable called 'c', initialized to an initial value of 0.0 everywhere.

```
[7]: mesh = fp.CylindricalGrid1D(nr = 50, Lr = 1.0)
```

```
[8]: c = fp.CellVariable(mesh = mesh, value = 0.0)
```

Apply boundary condition: outer wall will be kept at 1.0. (Dirichlet boundary condition).

```
[9]: c.constrain(1.0, mesh.facesRight)
```


We should also define our partial differential equation! (With Fipy, this step comes last).

```
[10]: eq = fp.TransientTerm(var=c) == fp.DiffusionTerm(var=c, coeff=1.0)
```

Now, we solve the equation by taking time steps. We plot the solution at several specified time-points.

```
[11]: sample_i = [5,10,20,30,40,60,80,100,150,200,300,400,600,800]
for i in range(0,1001):
    if i in sample_i:
        plt.plot(c.mesh.cellCenters.value[0], c.value)
    eq.solve(dt = 0.001)
```

```
plt.xlabel('rel. radial position')
plt.ylabel('rel. temperature')
plt.title('FVM (Fipy) solution')
plt.show()
```



1.3 End.

[]: