

Chorin's Projection Method with Spectral Collocation for 2D Incompressible Navier Stokes

August 23, 2019

The setup and time discretization with Crank Nicholson remains the same as the finite difference version of this (see other directory in repo). Here, we focus on deriving a Chebyshev pseudo-spectral estimator.

Recall the NSE (with no force function)

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p &= \Delta \mathbf{u} \\ \nabla \cdot \mathbf{u} &= 0 \\ \mathbf{u} &= 0 \quad \text{on} \quad \partial\Omega\end{aligned}$$

Chorin: Step 1 Also recall that the Chorin's projection method first ignores pressure to compute an intermediate velocity field

$$\begin{aligned}\frac{\partial \mathbf{u}^*}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= \Delta \mathbf{u}^* \\ \mathbf{u}^* &= 0 \quad \text{on} \quad \partial\Omega\end{aligned}$$

We discretize time first with Adams-Bashford and implicit Crank-Nicholson

$$\frac{\mathbf{u}^* - \mathbf{u}^n}{\Delta t} + (\mathbf{u}^{\frac{n+1}{2}} \cdot \nabla) \mathbf{u}^{\frac{n+1}{2}} = \Delta \mathbf{u}^{\frac{n+1}{2}}$$

So far things are identical to the finite difference version of Chorin's. But to solve this, we do not discretize spatial dimensions anymore: we use (pseudo)spectral methods:

We want to approximate the solution \mathbf{u}^* as truncated series of Chebyshev polynomials: $\{T_k(x)\}_{k=1}^{\infty}$ where $T_k(x) = \cos(k \cos^{-1} x)$; each polynomial is restricted to $[-1, 1]$.

$$u_N^*(x) = \sum_{k=0}^{\infty} \hat{u}_k^* T_k(x) \approx \sum_{k=0}^N \hat{u}_k^* T_k(x)$$

where $\mathbf{u}^* = (u^*, v^*)$. We just write the derivation for the first equation for simplicity. To get the Chebyshev coefficients, we have to compute the (normalized) inner product:

$$\hat{u}_k = \frac{2}{\pi c_k} \int_{-1}^1 u T_k w dx$$

where $c_k = \begin{cases} 2 & \text{if } k = 0 \\ 1 & \text{if } k \geq 1 \end{cases}$. This is hard to compute, so we need to estimate the integral. Normally, this would require a lot of points but we can get away with carefully selected ones and a Gaussian quadrature. The best points turn out to be the roots of another Chebyshev polynomial with degree one higher; these are called the Gauss-Lobatto points – they end up with denser spread near the boundaries (-1 and 1).

$$x_i = \cos \frac{\pi i}{N}, i = 0, \dots, N \quad (1)$$

resulting in,

$$\hat{u}_k = \frac{2}{\bar{c}_k N} \sum_{i=0}^N \frac{1}{\bar{c}_i} u_i T_k(x_i), k = 0, \dots, N \quad (2)$$

where $\bar{c}_k = \begin{cases} 2 & \text{if } k = 0 \\ 1 & \text{if } 1 \leq k \leq N-1 \\ 2 & \text{if } k = N \end{cases}$. Note that to convert back and forth between spectral coefficients (\hat{u}_k) and the values at the collocation points ($u_N(x_i)$) is a matrix multiplication. To be explicit, let $\mathcal{T} = [\cos k\pi i/N], k, i = 0, \dots, N$ and $\mathcal{T}^{-1} = [2(\cos \pi i/N)/(\bar{c}_k \bar{c}_i N)]$. Then,

$$\begin{aligned} \mathcal{U}^* &= \mathcal{T} \hat{\mathcal{U}}^* \\ \hat{\mathcal{U}}^* &= \mathcal{T} \mathcal{U}^* \end{aligned}$$

where $\mathcal{U}^* = [u^*(x_0), \dots, u^*(x_N)]$ and $\hat{\mathcal{U}}^* = [\hat{u}_0, \dots, \hat{u}_N]$.