

<u>MA (EU) number</u>	<u>(Invented) name</u>	<u>Strength</u>	<u>Pharmaceutical Form</u>	<u>Route of Administration</u>	<u>Immediate Packaging</u>	<u>Content (concentration)</u>	<u>Pack size</u>
EU/1/25/1913/001	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	1 pre-filled syringe
EU/1/25/1913/002	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	1 pre-filled syringe + 1 needle
EU/1/25/1913/003	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	1 pre-filled syringe + 2 needles
EU/1/25/1913/004	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	10 pre-filled syringes
EU/1/25/1913/005	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	10 pre-filled syringes + 10 needles
EU/1/25/1913/006	CAPVAXIVE	-- <sup>1</sup>	Solution for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	10 pre-filled syringes + 20 needles

--<sup>1</sup>

One dose (0.5ml) contains:

Pneumococcal polysaccharide serotype 3 <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 6A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 7F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 8 <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 9N <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 10A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 11A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 12F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 15A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide from deOAc15B (de-O-acetylated serotype 15B) <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 16F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 17F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 19A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 20A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 22F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 23A <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 23B <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 24F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 31 <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 33F <sup>1</sup>	4 mcg
Pneumococcal polysaccharide serotype 35B <sup>1</sup>	4 mcg

<sup>1</sup>Conjugated to CRM<sub>197</sub> carrier protein. CRM<sub>197</sub> is a nontoxic mutant of diphtheria toxin (originating from *Corynebacterium diphtheriae* C7) expressed recombinantly in *Pseudomonas fluorescens*.

One dose (0.5 mL) contains approximately 65 mcg CRM<sub>197</sub> carrier protein.