




Thresholds = mean of the last 10 of 14 reversals (recomputed from the trial CSV); dB = 20 log10(A). Open circles: interval_bias; +: near_ceiling (all retained). Means weight conditions equally.
The Won et al. (2015) literature line of the PDF is not part of these data and is not redrawn.




Shading: 5th reversal to run end; dashed: mean of the final 10 reversals; orange diamonds = reversals used. IB = interval_bias, NC = near_ceiling (stored app flags).



Shading: 5th reversal to run end; dashed: mean of the final 10 reversals; orange diamonds = reversals used. IB = interval_bias, NC = near_ceiling (stored app flags).


STM | threshold changes between repeat runs

A 3 of 12 cells with |difference| > 3 dB; 3 have a third run



B Repeat thresholds and descriptive summaries (dB)

Condition	Ear	n	Run1	Run2	Run3	Mean	SD	R2-R1
0.5 c/o 5 Hz	R	2	-20.2	-18.4	—	-19.30	1.27	+1.8
0.5 c/o 5 Hz	L	2	-10.2	-11.2	—	-10.70	0.71	-1.0
0.5 c/o 10 Hz	R	2	-13.8	-14.6	—	-14.20	0.57	-0.8
0.5 c/o 10 Hz	L	3	-10.2	-15.6	-9.2	-11.67	3.44	-5.4
1 c/o 5 Hz	R	2	-13.2	-11.2	—	-12.20	1.41	+2.0
1 c/o 5 Hz	L	2	-7.0	-9.2	—	-8.10	1.56	-2.2
1 c/o 10 Hz	R	3	-9.6	-13.8	-11.0	-11.47	2.14	-4.2
1 c/o 10 Hz	L	2	-7.0	-4.4	—	-5.70	1.84	+2.6
2 c/o 5 Hz	R	2	-10.2	-8.6	—	-9.40	1.13	+1.6
2 c/o 5 Hz	L	3	-6.8	-10.2	-6.2	-7.73	2.16	-3.4
2 c/o 10 Hz	R	2	-4.0	-5.2	—	-4.60	0.85	-1.2
2 c/o 10 Hz	L	2	-6.4	-4.8	—	-5.60	1.13	+1.6



1370 valid trials, 1037 correct; 0 technical failures. Ceiling-blocked moves: 0 in analysis windows, 1 in an initial segment.