



UNIVERSITATEA BABEȘ-BOLYAI
BABEȘ-BOLYAI TUDOMÁNYEGYETEM
BABEȘ-BOLYAI UNIVERSITÁT
BABEȘ-BOLYAI UNIVERSITY
TRADITIO ET EXCELLENTIA

Mathematical Analysis

1st Year Computer Science

Mihai Nechita *

Last update: October 15, 2025

Contents

Real numbers	2
Sequences	8

*math.ubbcluj.ro/~mihai.nechita. If you find any typos, please let me know on Teams or by email.

※ Real numbers

Let us start with some standard notation: $\emptyset$ is the empty set; $\mathbb{N} = \{1, 2, \dots\}$ the set of natural numbers; $\mathbb{Z} = \{\dots, -1, 0, 1, \dots\} = \{m - n \mid m, n \in \mathbb{N}\}$ the set of integers; $\mathbb{Q} = \{\frac{m}{n} \mid m, n \in \mathbb{Z}, n \neq 0\}$ the set of rational numbers; $\mathbb{R}$ the set of real numbers.

You are very much used with the real numbers. However, it is not trivial to define them in a rigorous way, see for example [1] or [2] for different methods of constructing $\mathbb{R}$ from $\mathbb{Q}$. We will straightforwardly start working with the real numbers – for this reason some of their properties, e.g. Definition 1.5, will simply be given as definitions.

Definition 1.1

Let A be a subset of $\mathbb{R}$, denoted as $A \subseteq \mathbb{R}$. We define $x \in \mathbb{R}$ to be

a *lower bound* for A if $x \leq a, \forall a \in A$; an *upper bound* for A if $x \geq a, \forall a \in A$.

We define

$lb(A) := \{x \in \mathbb{R} \mid x \leq a, \forall a \in A\}$ the set of lower bounds of A ,

$ub(A) := \{x \in \mathbb{R} \mid x \geq a, \forall a \in A\}$ the set of upper bounds of A .

We define $x \in \mathbb{R}$ to be

the *minimum* of A if $x \in lb(A) \cap A$; the *maximum* of A if $x \in ub(A) \cap A$,

denoted by $\min(A)$, respectively $\max(A)$. In other words, we have that

$\min(A) \in A$ and $\min(A) \leq a, \forall a \in A$; $\max(A) \in A$ and $\max(A) \geq a, \forall a \in A$.

Note that there are sets which do not have minimum or maximum, e.g. $(0, 1)$.

Definition 1.2

A set $A \subseteq \mathbb{R}$ is defined to be

- bounded (from) below if $lb(A) \neq \emptyset$;
- bounded (from) above if $ub(A) \neq \emptyset$;
- bounded if it is both bounded below and above;
- unbounded if it is not bounded.

Definition 1.3

We say that $x \in \mathbb{R}$ is the *supremum* of $A \subseteq \mathbb{R}$, $x := \sup(A)$, if and only if:

1. $x \geq a, \forall a \in A$, that is $x \in ub(A)$.
2. if u is an upper bound for A , then $x \leq u$.

The supremum is *the least upper bound*, i.e. $\sup(A) := \min(ub(A))$.

Definition 1.4

We say that $x \in \mathbb{R}$ is the *infimum* of $A \subseteq \mathbb{R}$, $x := \inf(A)$, if and only if:

1. $x \leq a, \forall a \in A$, that is $x \in lb(A)$.
2. if u is a lower bound for A , then $x \geq u$.

The infimum is *the greatest lower bound*, i.e. $\inf(A) := \max(lb(A))$.

Definition 1.5 (Completeness Axiom)

Every set $A \subseteq \mathbb{R}$ that is bounded above has a supremum. Similarly, every set $A \subseteq \mathbb{R}$ that is bounded below has an infimum.

Note that if A has a maximum, then $\sup(A) = \max(A)$. Similarly, if A has a minimum, then $\inf(A) = \min(A)$. Also, if $\sup(A) \in A$, then $\max(A) = \sup(A)$.

Example 1.6

(a) $A = \{\frac{1}{n} \mid n \in \mathbb{N}\}$, $\sup(A) = 1 = \max(A)$, $\inf(A) = 0$, $\nexists \min(A)$.

(b) $A = \{x \in \mathbb{Q} \mid x^2 \leq 2\}$, $\sup(A) = \sqrt{2}$, $\nexists \max(A)$, $\inf(A) = -\sqrt{2}$, $\nexists \min(A)$.

Proposition 1.7

Let $A \subseteq \mathbb{R}$ be a bounded set. For $\sup(A)$ and $\inf(A)$ the following are true:

$$\forall \varepsilon > 0, \exists x \in A \text{ such that } \sup(A) - \varepsilon < x,$$

$$\forall \varepsilon > 0, \exists x \in A \text{ such that } x < \inf(A) + \varepsilon.$$

Proof. By definition, for any $y < \sup(A)$, say $y = \sup(A) - \varepsilon$ with $\varepsilon > 0$, we have that $y \notin ub(A)$. Hence there exists $x \in A$ such that $y < x$. Similar proof for $\inf(A)$. $\square$

Proposition 1.8

Let $A \subseteq B \subseteq \mathbb{R}$ be (nonempty) bounded sets. Then

$$\inf(B) \leq \inf(A) \leq \sup(A) \leq \sup(B)$$

and

$$\sup(A \cup B) = \max\{\sup(A), \sup(B)\},$$

$$\inf(A \cup B) = \min\{\inf(A), \inf(B)\}.$$

Proof. It follows directly from the definitions. □

Definition 1.9

Define the *extended real line* $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$, where ∞ and $-\infty$ are such that

$$\forall x \in \mathbb{R}, -\infty < x < \infty.$$

If a set A is not bounded above, we define $\sup(A) := \infty$.

If a set A is not bounded below, we define $\inf(A) := -\infty$.

[Seminar] The empty set $\emptyset$ is bounded by any number. In $\overline{\mathbb{R}}$, $\sup(\emptyset) = -\infty$, $\inf(\emptyset) = \infty$.

Definition 1.10

A set $V \subseteq \mathbb{R}$ is a *neighborhood (vecinity)* of $x \in \mathbb{R}$ if

$$\exists \varepsilon > 0 \text{ such that } (x - \varepsilon, x + \varepsilon) \subseteq V.$$

A set $V \subseteq \mathbb{R}$ is a *neighborhood* of ∞ if $\exists a \in \mathbb{R}$ such that $(a, \infty) \subseteq V$.

A set $V \subseteq \mathbb{R}$ is a *neighborhood* of $-\infty$ if $\exists a \in \mathbb{R}$ such that $(-\infty, a) \subseteq V$.

We denote all the neighborhoods of x by

$$\mathcal{V}(x) := \{V \subseteq \mathbb{R} \mid V \text{ is a neighborhood of } x\}.$$

Definition 1.11

Let $A \subseteq \mathbb{R}$. The following set is called the *interior* of A

$$\text{int}(A) := \{x \in \mathbb{R} \mid \exists V \in \mathcal{V}(x) \text{ such that } V \subseteq A\},$$

and the following set is called the *closure* of A

$$\text{cl}(A) := \{x \in \mathbb{R} \mid \forall V \in \mathcal{V}(x), V \cap A \neq \emptyset\}.$$

Proposition 1.12

For any $A \subseteq \mathbb{R}$, it holds that $\text{int}(A) \subseteq A \subseteq \text{cl}(A)$.

Proof. To prove that $\text{int}(A) \subseteq A$ we prove that if $x \in \text{int}(A)$, then $x \in A$. Let $x \in \text{int}(A)$, then $\exists \varepsilon > 0$ such that $(x - \varepsilon, x + \varepsilon) \subseteq A$. Since $x \in (x - \varepsilon, x + \varepsilon)$, we have that $x \in A$.

To prove that $A \subseteq \text{cl}(A)$ we show that if $x \in A$, then $x \in \text{cl}(A)$. Let $x \in A$. Then for any $V \in \mathcal{V}(x)$ it holds that $x \in V$, giving that $x \in V \cap A$. Hence $x \in \text{cl}(A)$ since $V \cap A \neq \emptyset$. $\square$

Definition 1.13

If $A = \text{int}(A)$, then A is called *open*. If $A = \text{cl}(A)$, then A is called *closed*.

Remark 1.14

To prove that a set A is open, it is sufficient to prove that $A \subseteq \text{int}(A)$.

To prove that a set A is closed, it is sufficient to prove that $\text{cl}(A) \subseteq A$.

Proposition 1.15

The following statements are true:

- The complement of an open set is closed.
- The complement of a closed set is open.

Proof. Let us prove the first statement, the other one being similar. Consider A an open set, i.e. $A = \text{int}(A)$, and denote by $A^c = \{x \in \mathbb{R} \mid x \notin A\}$ its complement. To prove that A^c is closed, we prove that $\text{cl}(A^c) \subseteq A^c$. Consider $x \in \text{cl}(A^c)$ and let's assume that $x \notin A^c$, i.e. $x \in A$, aiming to obtain a contradiction. Since A is open, there exists $V \in \mathcal{V}(x)$ such that $V \subseteq A$, giving that $V \cap A^c = \emptyset$: contradiction with $x \in \text{cl}(A^c)$. Hence the assumption $x \notin A^c$ is false, and we have that if $x \in \text{cl}(A^c)$, then $x \in A^c$. In other words, $\text{cl}(A^c) \subseteq A^c$. $\square$

Proposition 1.16

Any union of open sets is open. Any intersection of closed sets is closed. Any finite intersection of open sets is open. Any finite union of closed sets is closed.

Proof. (Optional) Left as extra homework. $\square$

Exercises. Seminar 1

1. Find the lower and the upper bounds, then $\sup$, $\inf$, $\max$, $\min$ for each of the following:

- | | |
|----------------------------------|---------------------------------|
| (a) $[-3, 2) \cup \{3\}$. | (c) $(-5, 5) \cap \mathbb{Z}$. |
| (b) $(-1, 1] \cup (2, \infty)$. | (d) $\emptyset$. |

2. Find the $\sup$, $\inf$, $\max$, $\min$ for each of the following sets:

- | | |
|--|--|
| (a) $\{x \in \mathbb{Q} \mid x^2 < 3\}$. | (c) $\{\frac{n}{n+1} \mid n \in \mathbb{N}\}$. |
| (b) $\{x^2 - 4x + 3 \mid x \in \mathbb{R}\}$. | (d) $\{2^{-k} + 3^{-m} \mid k, m \in \mathbb{N}\}$. |

3. Suppose that S is nonempty and bounded above. Show that the set $-S := \{-x \mid x \in S\}$ is bounded below and $\inf(-S) = -\sup(S)$.

4. Let $f : D \rightarrow \mathbb{R}$ and $g : D \rightarrow \mathbb{R}$ be two functions defined on a nonempty set D . Prove that

$$\inf_{x \in D} (f(x) + g(x)) \geq \inf_{x \in D} f(x) + \inf_{x \in D} g(x) \quad \text{and} \quad \sup_{x \in D} (f(x) + g(x)) \leq \sup_{x \in D} f(x) + \sup_{x \in D} g(x).$$

Give examples where the above inequalities are strict.

5. Which of the following sets are neighborhoods of 0?

$$[-1, 1] \cup \{2\}; \quad (-1, 1) \cap \mathbb{Q}; \quad \bigcap_{n=1}^{\infty} \left[-\frac{1}{n}, \frac{1}{n}\right].$$

6. Let $x \in \mathbb{R}$ and $U, V \in \mathcal{V}(x)$. Prove that $U \cap V \in \mathcal{V}(x)$.

7. Find the interior and the closure for each of the following sets:

(a) $(1, 2]$.

(c) $(-1, 1] \cup (2, \infty)$.

(b) $[-3, 2) \cup \{3\}$.

(d) $(-5, 5) \cap \mathbb{Z}$.

Exercises. Self study

1. Let $a, b \in \mathbb{R}$ with $a > 0$. If $S \subset \mathbb{R}$ is nonempty and bounded above, prove that

$$\sup_{x \in S} (ax + b) = a \sup(S) + b.$$

2. Let $a, b \in \mathbb{R}$. Prove that there exist neighborhoods $U \in \mathcal{V}(a)$ and $V \in \mathcal{V}(b)$ such that

$$U \cap V = \emptyset.$$

3. Let $A = (0, 1) \cap \mathbb{Q}$. Show rigorously (using the definitions) that

$$\inf A = 0, \sup A = 1, \text{int} A = \emptyset \text{ and } \text{cl} A = [0, 1].$$

※ Sequences

A set $\{x_n \mid n \in \mathbb{N}\}$ is called a sequence and is denoted by $(x_n)_{n \in \mathbb{N}}$ or simply (x_n) . A sequence (x_n) is bounded above (or below) if the set $\{x_n \mid n \in \mathbb{N}\}$ is bounded above (or below). A sequence (x_n) is increasing if $x_{n+1} \geq x_n, \forall n \in \mathbb{N}$, and decreasing if $x_{n+1} \leq x_n, \forall n \in \mathbb{N}$. A sequence is monotone if it is either increasing or decreasing.

Definition 2.1

A sequence (x_n) has a limit $\ell \in \overline{\mathbb{R}}$, and we write $\lim_{n \rightarrow \infty} x_n = \ell$ or $x_n \rightarrow \ell$, if

$$\forall V \in \mathcal{V}(\ell), \exists N_V \in \mathbb{N} \text{ such that } x_n \in V, \forall n \geq N_V.$$

If $\ell \in \mathbb{R}$, we say that (x_n) converges to ℓ : $\forall \varepsilon > 0, \exists N_\varepsilon \in \mathbb{N}$ such that $|x_n - \ell| < \varepsilon, \forall n \geq N_\varepsilon$. $x_n \rightarrow \infty$ if $\forall a > 0, \exists N_a \in \mathbb{N}$ such that $x_n > a, \forall n \geq N_a$.

$x_n \rightarrow -\infty$ if $\forall a < 0, \exists N_a \in \mathbb{N}$ such that $x_n < a, \forall n \geq N_a$.

Example 2.2

Prove that $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$. Let $\varepsilon > 0$; we need to find an index $N_\varepsilon \in \mathbb{N}$ such that $\frac{1}{n} < \varepsilon, \forall n \geq N_\varepsilon$. Since $\frac{1}{n} \leq \frac{1}{N_\varepsilon}$ for $n \geq N_\varepsilon$, it is enough to have $\frac{1}{N_\varepsilon} < \varepsilon$ or $N_\varepsilon > \frac{1}{\varepsilon}$. We conclude by taking $N_\varepsilon > \frac{1}{\varepsilon}$, e.g. $N_\varepsilon = [1/\varepsilon] + 1$.

Proposition 2.3

A sequence (x_n) converges to $\ell \in \mathbb{R}$ if and only if $\lim_{n \rightarrow \infty} |x_n - \ell| = 0$.

Proposition 2.4

Any convergent sequence is bounded.

Proof. Let $\ell \in \mathbb{R}$ be the limit. For $\varepsilon = 1$, there exists $N_1 \in \mathbb{N}$ such that $|x_n - \ell| < 1, \forall n \geq N_1$. All the terms after index N_1 are in $(\ell - 1, \ell + 1)$. Since there are finitely many terms before index N_1 , we conclude that the sequence is bounded. $\square$

Theorem 2.5 (Weierstrass)

Any monotone and bounded sequence is convergent.

Proof. Assume that the sequence is increasing, for example. Let $S = \{x_n \mid n \in \mathbb{N}\}$ and consider $\sup(S) \in \mathbb{R}$ (we know that S is bounded). From [Proposition 1.7](#) we have that

$$\forall \varepsilon > 0, \exists N_\varepsilon \in \mathbb{N} \text{ such that } \sup(S) - \varepsilon < x_{N_\varepsilon}.$$

As (x_n) is increasing, $\sup(S) - \varepsilon < x_{N_\varepsilon} \leq x_n \forall n \geq N_\varepsilon$. Hence $\sup(S) - x_n < \varepsilon, \forall n \geq N_\varepsilon$. The sequence converges to $\sup(S)$ by [Definition 2.1](#). Similarly, a decreasing and bounded sequence converges to its infimum. $\square$

Proposition 2.6

Any monotone sequence has a limit in $\overline{\mathbb{R}}$.

Proof. If the sequence is bounded and monotone, then it is convergent by the Weierstrass theorem. If the sequence is unbounded and monotone, then its limit will be infinite. $\square$

Theorem 2.7 (Squeeze/Sandwich theorem)

Let (x_n) , (y_n) , (z_n) be sequences for which there is an $n_0 \in \mathbb{N}$ such that

$$x_n \leq y_n \leq z_n, \forall n \geq n_0,$$

and

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} z_n.$$

Then

$$\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} z_n.$$

Proof. Let $\ell := \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} z_n$ and assume first that $\ell \in \mathbb{R}$. Let $\varepsilon > 0$. Then

$$\exists N_1 \in \mathbb{N} \text{ such that } |x_n - \ell| < \varepsilon, \forall n \geq N_1$$

and

$$\exists N_2 \in \mathbb{N} \text{ such that } |z_n - \ell| < \varepsilon, \forall n \geq N_2.$$

Taking $N_\varepsilon := \max\{N_1, N_2\}$, we have that

$$|y_n - \ell| \leq \max\{|x_n - \ell|, |z_n - \ell|\} < \varepsilon, \forall n \geq N_\varepsilon,$$

hence the conclusion. When ℓ is infinite the proof is similar. □

Theorem 2.8 (Cantor's nested intervals)

Let (a_n) be increasing and (b_n) decreasing such that $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n$, $\forall n \in \mathbb{N}$. Consider the closed intervals $I_n := [a_n, b_n]$, with $I_{n+1} \subseteq I_n$. If $\lim_{n \rightarrow \infty} (b_n - a_n) = 0$, then there exists $x \in \mathbb{R}$ such that

$$\bigcap_{n=1}^{\infty} I_n = \{x\}.$$

Proof. Consider the bounded sets $A := \{a_n \mid n \in \mathbb{N}\}$ and $B := \{b_n \mid n \in \mathbb{N}\}$. For any $k \in \mathbb{N}$, we have that

$$a_k \leq \sup(A) \leq b_k$$

and

$$b_k \geq \inf(B) \geq a_k.$$

Hence by the squeeze [Theorem 2.7](#) we have that $\sup(A) = \inf(B)$ and $\bigcap_{n=1}^{\infty} I_n = \{\sup(A)\}$. □

Theorem 2.9 (Bolzano-Weierstrass)

Any bounded sequence has a convergent subsequence.

Proof. Consider the bounded set $A := \{x_n \mid n \in \mathbb{N}\}$. Let $a_1 := \inf(A)$ and $b_1 := \sup(A)$, and define $I_1 := [a_1, b_1]$. Bisect I_1 and notice that at least one of the two halves must contain infinitely many terms from the sequence. Take $I_2 := [a_2, b_2]$ to be the half that does. Continuing this procedure we obtain for each $k \in \mathbb{N}$ an interval $I_k := [a_k, b_k]$ containing (at least) a term $x_{n_k} \in A$, such that $I_{k+1} \subseteq I_k$ and $b_k - a_k \rightarrow 0$.

From Cantor's nested intervals [Theorem 2.8](#) we have that there exists $x \in \mathbb{R}$ such that $\bigcap_{n=1}^{\infty} I_n = \{x\}$, and hence the subsequence (x_{n_k}) converges to x . $\square$

Proposition 2.10 (Ratio test)

Let (x_n) be a sequence with positive terms such that

$$\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = \ell \in \overline{\mathbb{R}}.$$

If $\ell < 1$, then $\lim_{n \rightarrow \infty} x_n = 0$. If $\ell > 1$, then $\lim_{n \rightarrow \infty} x_n = \infty$. If $\ell = 1$, the test is inconclusive.

Proof. Let us consider the case $\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = \ell < 1$. Let $\varepsilon > 0$ such that $q := \ell + \varepsilon < 1$. Then there exists $N \in \mathbb{N}$ such that $\frac{x_{n+1}}{x_n} - \ell < \varepsilon$, $\forall n \geq N$. We have that $x_{n+1} < x_n \cdot q$, $\forall n \geq N$, hence $x_n < q^{n-N} x_N$, giving that $0 < x_n < q^n \frac{x_N}{q^N}$. Since $q < 1$, $q^n \rightarrow 0$, and $x_n \rightarrow 0$ by the squeeze theorem. The proof is similar when $\ell > 1$. $\square$

Lemma 2.11 (Stolz-Cesàro)

Let $(a_n), (b_n)$ be two sequences such that (i) $a_n \rightarrow 0$ and $b_n \rightarrow 0$ with (b_n) decreasing; or (ii) $b_n \rightarrow \infty$ with (b_n) increasing.

$$\text{If } \lim_{n \rightarrow \infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = \ell, \text{ then } \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \ell.$$

Proof. (Optional) See [3, Theorem 2.5.6]. $\square$

Definition 2.12 (Cauchy sequence)

A sequence (x_n) is called *Cauchy* (or *fundamental*) if

$$\forall \varepsilon > 0, \exists N_\varepsilon \in \mathbb{N} \text{ such that } |x_m - x_n| < \varepsilon, \forall m, n \geq N_\varepsilon.$$

Proposition 2.13

Any Cauchy sequence is bounded.

Proof. For $\varepsilon = 1$, there exists $N_1 \in \mathbb{N}$ such that $|x_m - x_n| < 1, \forall m, n \geq N_1$. In particular, $|x_n - x_{N_1}| < 1, \forall n \geq N_1$, hence all the terms after index N_1 are in $(x_{N_1} - 1, x_{N_1} + 1)$. There are finitely many terms before index N_1 , thus we conclude that the sequence is bounded. $\square$

Theorem 2.14

A sequence is convergent if and only if it is Cauchy.

Proof. Let's consider first a convergent sequence (x_n) with $x_n \rightarrow \ell$. For any $\varepsilon > 0$, there exists $N_\varepsilon \in \mathbb{N}$ such that $|x_n - \ell| < \frac{\varepsilon}{2}$, for any $n \geq N_\varepsilon$. Then $|x_m - x_n| \leq |x_m - \ell| + |x_n - \ell| < \varepsilon$, for any $n \geq N_\varepsilon$. Hence the sequence (x_n) is Cauchy.

Assume now that (x_n) is a Cauchy sequence. From the previous proposition we have that (x_n) must be bounded, and thus it has a convergent subsequence (x_{n_k}) , $x_{n_k} \rightarrow x \in \mathbb{R}$. Let $\varepsilon > 0$. There exists thus $K_\varepsilon \in \mathbb{N}$ such that $|x_{n_k} - x| < \varepsilon, \forall k \geq K_\varepsilon$. Also, there exists $N_\varepsilon \in \mathbb{N}$ such that $|x_m - x_n| < \varepsilon, \forall m, n \geq N_\varepsilon$. In particular, $|x_{n_k} - x_n| < \varepsilon, \forall k, n \geq N_\varepsilon$. Hence $|x_n - x| \leq |x_n - x_{n_k}| + |x_{n_k} - x| < 2\varepsilon, \forall n \geq \max\{K_\varepsilon, N_\varepsilon\}$, meaning that $x_n \rightarrow x$. $\square$

Example 2.15

The sequence defined by $x_n = 1 + \frac{1}{2} + \dots + \frac{1}{n}$ is not convergent. Indeed, one can see, for example, that

$$x_{2n} - x_n = \frac{1}{n+1} + \dots + \frac{1}{2n} > \frac{n}{2n},$$

hence $x_{2n} - x_n > \frac{1}{2}$ for any $n \in \mathbb{N}$. Thus (x_n) is not Cauchy, hence it is not convergent.

Definition 2.16

For a sequence (x_n) we define the set of its *limit points* by

$$\text{LIM}(x_n) := \{x \in \overline{\mathbb{R}} \mid \text{there exists a subsequence } (x_{n_k}) \text{ s.t. } x_{n_k} \rightarrow x\},$$

and

$$\liminf_{n \rightarrow \infty} x_n := \inf (\text{LIM}(x_n)),$$

$$\limsup_{n \rightarrow \infty} x_n := \sup (\text{LIM}(x_n)).$$

Example 2.17

For $x_n = \frac{(-1)^n n}{n+1}$, $\text{LIM}(x_n) = \{-1, 1\}$, $\liminf_{n \rightarrow \infty} x_n = -1$, $\limsup_{n \rightarrow \infty} x_n = 1$.

Proposition 2.18

$\lim_{n \rightarrow \infty} x_n = \ell \in \overline{\mathbb{R}}$ if and only if $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = \ell$.

Exercises. Seminar 2

1. Prove using the ε -definition that $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$.
2. Study if the sequence (x_n) is bounded, monotone, and convergent, for each of the following:

(a) $x_n = \sqrt{n+1} - \sqrt{n}$. (b) $x_n = \frac{1}{1 \cdot 2} + \dots + \frac{1}{n(n+1)}$. (c) $x_n = \frac{2^n}{n!}$.

3. Find the limit for each of the following sequences:

(a) $\sqrt{n}(\sqrt{n+1} - \sqrt{n})$. (c) $\sqrt[n]{n}$. (e) $(a_1^n + a_2^n + \dots + a_k^n)^{\frac{1}{n}}$
 (b) $\frac{2^n + (-1)^n}{3^n}$. (d) $\frac{(an+1)^2}{4n^2 - 2n + 1}$, $a \in \mathbb{R}$. with $a_i > 0$.

4. Consider the sequence (e_n) given by $e_n = \left(1 + \frac{1}{n}\right)^n$. Prove that (e_n) is increasing and bounded, hence convergent. Its limit is denoted by e .

5. Find the limit for each of the following sequences:

(a) $\left(\frac{2n+1}{2n-1}\right)^n$. (b) $n(\ln(n+2) - \ln(n+1))$.

6. (a) Let (x_n) be convergent. What can you say about the sequence (a_n) of averages,

$$a_n = \frac{x_1 + x_2 + \dots + x_n}{n}?$$

Can the averages converge, but the sequence not?

- (b) Compare $1 + \frac{1}{2} + \dots + \frac{1}{n}$ with n and $\ln n$ by taking the ratio, respectively.

7. Let (x_n) be a sequence such that $\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = \ell$. Prove that $\lim_{n \rightarrow \infty} \sqrt[n]{x_n} = \ell$.

8. Find the limit for each of the following sequences: (a) $\frac{n}{\sqrt[n]{n!}}$. (b) $\frac{1^p + 2^p + 3^p + \dots + n^p}{n^{p+1}}$, $p \in \mathbb{N}$.

Exercises. Self study

1. Prove using the ε -definition that $\lim_{n \rightarrow \infty} \frac{n+1}{2n+3} = \frac{1}{2}$.

2. Find the limit for each of the following sequences:

(a) $(1 + 2 + \dots + n)^{1/n}$.

(b) $\left(\frac{\ln(n+1)}{\ln n}\right)^n$.

(c) $\frac{n^n}{1 + 2^2 + 3^3 + \dots + n^n}$.

3. For $x_n = \frac{\sin(n)}{n}$ study if (x_n) is bounded, monotone, and convergent. Find its limit.

4. Prove that the sequence (x_n) given by

$$x_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} - \ln n$$

is decreasing and bounded, hence convergent. Its limit is typically denoted by γ and is known as the Euler-Mascheroni constant.

※ References

- [1] W. Rudin, *Principles of Mathematical Analysis 3rd ed*, McGraw-Hill, 1976.
- [2] T. Tao, *Analysis I*, Springer, 2016.
- [3] Ș. Cobzaș, *Analiză matematică – Calcul Diferențial*, Presa Universitară Clujeană, 2019.
- [4] J.E. Marsden, A. Tromba, *Vector Calculus 6th ed*, W.H. Freeman and Company, 2012
- [5] M. Oberguggenberger, A. Ostermann, *Analysis for Computer Scientists*, Springer, 2018
- [6] G. Strang, *Linear Algebra and Learning from Data*, Wellesley Cambridge Press, 2019