

Steklov's Problem for OPUC

Michel Alexis

joint with Alexander Aptekarev and Sergey Denisov

UW-Madison

ORAM, 2021


Orthogonal Polynomials on the Unit Circle (OPUC)

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.


Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.



Orthogonal Polynomials on the Unit Circle (OPUC)


Setting: $\mathbb{T} \subset \mathbb{C}$.



Can write $z = e^{i\theta}$.

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.




$$1, z, z^2, z^3, \dots$$

Can write $z = e^{i\theta}$.

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.




Can write $z = e^{i\theta}$.

$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.



Can write $z = e^{i\theta}$.


$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$



$$w(\theta) \frac{d\theta}{2\pi}$$

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.



Can write $z = e^{i\theta}$.


$$1, z, z^2, z^3, \dots \quad \perp \text{ w.r.t. } \frac{d\theta}{2\pi}$$

↓ Gram-Schmidt

$$\downarrow \\ w(\theta) \frac{d\theta}{2\pi}$$

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.




Can write $z = e^{i\theta}$.

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.




Can write $z = e^{i\theta}$.

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$ is orthonormal polynomial of degree n .

Orthogonal Polynomials on the Unit Circle (OPUC)

Setting: $\mathbb{T} \subset \mathbb{C}$.



Can write $z = e^{i\theta}$.

$$\begin{array}{ccc} 1, z, z^2, z^3, \dots & \perp \text{ w.r.t. } & \frac{d\theta}{2\pi} \\ \downarrow \text{Gram-Schmidt} & & \downarrow \\ \varphi_0, \varphi_1, \varphi_2, \varphi_3, \dots & \perp \text{ w.r.t. } & w(\theta) \frac{d\theta}{2\pi} \end{array}$$

- $\varphi_n(z)$ is orthonormal polynomial of degree n .
- $\{\varphi_n(z)\}$ are called Orthogonal Polynomials on the Unit Circle.

How do OPUC behave?

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

Trend seems to be $|\varphi_n| \rightarrow w^{-1/2}$;

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

Trend seems to be $|\varphi_n| \rightarrow w^{-1/2}$; indeed, $\int |\varphi_n|^2 w = 1$ and φ_n oscillate more and more.

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

Trend seems to be $|\varphi_n| \rightarrow w^{-1/2}$; indeed, $\int |\varphi_n|^2 w = 1$ and φ_n oscillate more and more.


We expect this behavior pointwise as well.

How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

Trend seems to be $|\varphi_n| \rightarrow w^{-1/2}$; indeed, $\int |\varphi_n|^2 w = 1$ and φ_n oscillate more and more.

We expect this behavior pointwise as well.





How do OPUC behave?

- $\frac{1}{|\varphi_n|^2} \rightarrow w$ in the weak-* sense as measures.
- $\log \frac{1}{|\varphi_n|^2} \rightarrow \log w$ in L^1 and weak-* sense if $\int_{\mathbb{T}} \log w > -\infty$.

Trend seems to be $|\varphi_n| \rightarrow w^{-1/2}$; indeed, $\int |\varphi_n|^2 w = 1$ and φ_n oscillate more and more.

We expect this behavior pointwise as well.



$$|\varphi_n| \rightarrow w^{-1/2}$$

Conjecture (Steklov's Conjecture)


If $w \geq \delta > 0$, then $\{\varphi_n\}$ are bounded above, i.e. $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$.

$$|\varphi_n| \rightarrow w^{-1/2}$$

Conjecture (Steklov's Conjecture)

If $w \geq \delta > 0$, then $\{\varphi_n\}$ are bounded above, i.e. $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$.

False! Can create weights w which oscillate rapidly, forcing $\{\varphi_n\}$ to blow-up.




$$|\varphi_n| \rightarrow w^{-1/2}$$

Conjecture (Steklov's Conjecture)

If $w \geq \delta > 0$, then $\{\varphi_n\}$ are bounded above, i.e. $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$.

False! Can create weights w which oscillate rapidly, forcing $\{\varphi_n\}$ to blow-up.



But $\|\varphi_n\|_{L^2(w)} = 1$ by definition.


Maybe we're asking for too much regularity?

$$|\varphi_n| \rightarrow w^{-1/2}$$

Conjecture (Steklov's Conjecture)

If $w \geq \delta > 0$, then $\{\varphi_n\}$ are bounded above, i.e. $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$.

False! Can create weights w which oscillate rapidly, forcing $\{\varphi_n\}$ to blow-up.



But $\|\varphi_n\|_{L^2(w)} = 1$ by definition.

Maybe we're asking for too much regularity?

Problem (Steklov problem)


Does there exist $p > 2$ such that $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$?

$$|\varphi_n| \rightarrow w^{-1/2}$$

Conjecture (Steklov's Conjecture)

If $w \geq \delta > 0$, then $\{\varphi_n\}$ are bounded above, i.e. $\sup_n \|\varphi_n\|_{L^\infty(w)} < \infty$.

False! Can create weights w which oscillate rapidly, forcing $\{\varphi_n\}$ to blow-up.



But $\|\varphi_n\|_{L^2(w)} = 1$ by definition.

Maybe we're asking for too much regularity?

Problem (Steklov problem)

Does there exist $p > 2$ such that $\sup_n \|\varphi_n\|_{L^p(w)} < \infty$?

Remark: If $\int_{\mathbb{T}} \log w > -\infty$, then $|\varphi_n(z)| \sim |\Phi_n(z)|$, where $\Phi_n(z)$ are the monic orthogonal polynomials of degree n .

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq \Lambda$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq \textcolor{red}{X}1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof:

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq \chi 1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq \chi$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n = z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \end{cases}$$

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq X$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]} \Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]} w \Phi_n \quad (\text{orthogonal}) \end{cases}$$

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq X$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n & (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n & (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq K1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq X1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

If we can show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2, then done!

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq K1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

If we can show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2, then done!

- Indeed, $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1 + \epsilon)(1 - \epsilon) \leq 1 - \epsilon^2 < 1$

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq X1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

If we can show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2, then done!

- Indeed, $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq K1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

If we can show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2, then done!

- Indeed, $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1+\epsilon)(1-\epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p$$

Does there exist $p > 2$ such that $\sup_n \|\Phi_n\|_{L^p(w)} < \infty$?

Theorem (Nazarov, 2016)

Suppose $\epsilon \leq w \leq K1$. Then there exists $p > 2$ for which $\sup_n \|\Phi_n\|_{L^p} < \infty$.

Proof: $\mathcal{P}_{[0,n]} \stackrel{\text{def}}{=} \text{Projection onto } \text{Span}\{1, z, z^2, z^3, \dots, z^n\}$.

$$\begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]}\Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]}w\Phi_n \quad (\text{orthogonal}) \end{cases} \Rightarrow \Phi_n = z^n + \mathcal{P}_{[0,n-1]}(1-w)\Phi_n,$$

or rather $(I - \mathcal{P}_{[0,n-1]}(1-w))\Phi_n = z^n$.

If we can show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2, then done!

- Indeed, $\|\mathcal{P}_{[0,n-1]}(1-w)\|_{p,p} \leq (1 + \epsilon)(1 - \epsilon) \leq 1 - \epsilon^2 < 1$
- Now by geometric sum,

$$\|\Phi_n\|_p \leq \|(I - \mathcal{P}_{[0,n-1]}(1-w))^{-1}\|_{p,p} \|z^n\|_p \leq \left\| \sum_{k=0}^{\infty} (\mathcal{P}_{[0,n-1]}(1-w))^k \right\|_{p,p}$$


Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,



Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,






Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,






Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,



Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2




- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,



- Given $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$, one can show $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$ uniformly in n .

Show $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon$ for p close to 2

- Have uniform control of $\|\mathcal{P}_{[0,n-1]}\|_{p,p}$ since $\mathcal{P}_{[0,n-1]}$ is a linear combination of Hilbert transforms,



- Given $\|\mathcal{P}_{[0,n]}\|_{2,2} = 1$, one can show $\|\mathcal{P}_{[0,n]}\|_{p,p} \leq 1 + O(|p - 2|)$ uniformly in n .
- Choose p close enough to 2 so that $\|\mathcal{P}_{[0,n-1]}\|_{p,p} \leq 1 + \epsilon < 1$

□

A_p weights

A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty.$

A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :

A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
- $w \in A_2$ means w cannot be too singular/oscillatory.

A_p weights


- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.

A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.
 - Power weights $|\theta|^\alpha \in A_2$ if $-1 < \alpha < 1$


A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.
 - Power weights $|\theta|^\alpha \in A_2$ if $-1 < \alpha < 1$




A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{ arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.
 - Power weights $|\theta|^\alpha \in A_2$ if $-1 < \alpha < 1$
 - Can define A_p weights similarly for $1 < p < \infty$.



A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.
 - Power weights $|\theta|^\alpha \in A_2$ if $-1 < \alpha < 1$
 - Can define A_p weights similarly for $1 < p < \infty$.




Theorem (Hunt-Muckenhoupt-Wheeden)

If $w \in A_p$, then $\|w^{1/p} \mathcal{H} w^{-1/p}\|_{p,p} = \|\mathcal{H}\|_{L^p(w) \rightarrow L^p(w)} < \infty$.

A_p weights

- We say $w \in A_2$ if $\sup_{I: \text{arc in } \mathbb{T}} \left(\frac{1}{|I|} \int_I w \right) \left(\frac{1}{|I|} \int_I w^{-1} \right) < \infty$.
- Facts/notions about A_2 :
 - $w \in A_2$ means w cannot be too singular/oscillatory.
 - if $w \in A_2$ then $\log w \in \text{BMO}$.
 - Power weights $|\theta|^\alpha \in A_2$ if $-1 < \alpha < 1$
 - Can define A_p weights similarly for $1 < p < \infty$.



Theorem (Hunt-Muckenhoupt-Wheeden)

If $w \in A_p$, then $\|w^{1/p} \mathcal{H} w^{-1/p}\|_{p,p} = \|\mathcal{H}\|_{L^p(w) \rightarrow L^p(w)} < \infty$.

A_p plays nicely with the Hilbert transform \mathcal{H} , so makes sense to adapt previous proof to A_p weights.

The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

$$\bullet \begin{cases} \Phi_n &= z^n + \mathcal{P}_{[0,n-1]} \Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0,n-1]} w \Phi_n \quad (\text{orthogonal}) \end{cases}$$



The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0, n-1]} & \Phi_n \text{ (monic)} \\ 0 &= \mathcal{P}_{[0, n-1]} w & \Phi_n \text{ (orthogonal)} \end{cases}$$



The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0, n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= \mathcal{P}_{[0, n-1]} w \Phi_n \quad (\text{orthogonal}) \end{cases}$$



The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

$$\bullet \begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0, n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'} \mathcal{P}_{[0, n-1]} w^{1/p'} w^{1/p} \Phi_n \quad (\text{orthogonal}) \end{cases}$$



The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

- $$\begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'} w^{1/p} \Phi_n \quad (\text{orthogonal}) \end{cases}$$
- Subtract bottom from top and re-arrange to get

$$(I - Q_{w,p}) w^{1/p} \Phi_n = w^{1/p} z^n,$$

where $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$.



The Steklov problem for A_2 weights

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then $\sup_n \|w^{1/p} \Phi_n\|_{L^p} = \sup_n \|\Phi_n\|_{L^p(w)} < \infty$ for some $p > 2$.

Proof.

- $$\begin{cases} w^{1/p} \Phi_n &= w^{1/p} z^n + w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} w^{1/p} \Phi_n \quad (\text{monic}) \\ 0 &= w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'} w^{1/p} \Phi_n \quad (\text{orthogonal}) \end{cases}$$
- Subtract bottom from top and re-arrange to get

$$(I - Q_{w,p}) w^{1/p} \Phi_n = w^{1/p} z^n,$$

where $Q_{w,p} = w^{1/p} \mathcal{P}_{[0,n-1]} w^{-1/p} - w^{-1/p'} \mathcal{P}_{[0,n-1]} w^{1/p'}$.

- Can invert $I - Q_{w,p}$ for some p near 2 using spectral theory and analytic interpolation. (Skipped because not main goal of talk).



Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.

Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.
- But can also measure regularity from an operator perspective.

Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.
- But can also measure regularity from an operator perspective.
- Define $\mathcal{P}_{[0,n]}^w \stackrel{\text{def}}{=} \text{Proj}_w \text{Span} \{\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n\}$.

Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.
- But can also measure regularity from an operator perspective.
- Define $\mathcal{P}_{[0,n]}^w \stackrel{\text{def}}{=} \text{Proj}_w \text{Span} \{\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n\}$.
 - When $w = 1$ we just get back $\mathcal{P}_{[0,n]}$, bounded on L^p for $1 < p < \infty$.

Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.
- But can also measure regularity from an operator perspective.
- Define $\mathcal{P}_{[0,n]}^w \stackrel{\text{def}}{=} \text{Proj}_w \text{Span} \{\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n\}$.
 - When $w = 1$ we just get back $\mathcal{P}_{[0,n]}$, bounded on L^p for $1 < p < \infty$.

Problem

Does there exist $p \neq 2$ for which $\{\mathcal{P}_{[0,n]}^w\}_{n \geq 0}$ is uniformly bounded as operators on $L^p(w)$?

Another measure of regularity

- Till now, we measured the regularity of $\{\varphi_n\}$ using L^p norms.
- But can also measure regularity from an operator perspective.
- Define $\mathcal{P}_{[0,n]}^w \stackrel{\text{def}}{=} \text{Proj}_w \text{Span} \{\varphi_0, \varphi_1, \varphi_2, \dots, \varphi_n\}$.
 - When $w = 1$ we just get back $\mathcal{P}_{[0,n]}$, bounded on L^p for $1 < p < \infty$.

Problem

Does there exist $p \neq 2$ for which $\{\mathcal{P}_{[0,n]}^w\}_{n \geq 0}$ is uniformly bounded as operators on $L^p(w)$?

A: Yes!

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then there exists $\delta > 0$ such that whenever $|\frac{1}{p} - \frac{1}{2}| < \delta$, then

$$\sup_n \|\mathcal{P}_{[0,n]}^w\|_{L^p(w) \rightarrow L^p(w)} < \infty.$$

Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then there exists $\delta > 0$ such that whenever $|\frac{1}{p} - \frac{1}{2}| < \delta$, then

$$\sup_n \|\mathcal{P}_{[0,n]}^w\|_{L^p(w) \rightarrow L^p(w)} < \infty.$$

Proof.

One can show that when $w \in A_2$, we have $\mathcal{P}_{[0,\infty)}^w$ is bounded on $L^p(w)$. So suffices to show $\{\mathcal{P}_{[n+1,\infty)}^w\}_{n \geq 0}$ are bounded on $L^p(w)$.



Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then there exists $\delta > 0$ such that whenever $|\frac{1}{p} - \frac{1}{2}| < \delta$, then

$$\sup_n \|\mathcal{P}_{[0,n]}^w\|_{L^p(w) \rightarrow L^p(w)} < \infty.$$

Proof.

One can show that when $w \in A_2$, we have $\mathcal{P}_{[0,\infty)}^w$ is bounded on $L^p(w)$. So suffices to show $\{\mathcal{P}_{[n+1,\infty)}^w\}_{n \geq 0}$ are bounded on $L^p(w)$.

$$\begin{cases} 0 \\ \end{cases} = \mathcal{P}_{[0,n]}^w \mathcal{P}_{[n+1,\infty)}^w \quad (\text{orthogonality})$$



Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then there exists $\delta > 0$ such that whenever $|\frac{1}{p} - \frac{1}{2}| < \delta$, then

$$\sup_n \|\mathcal{P}_{[0,n]}^w\|_{L^p(w) \rightarrow L^p(w)} < \infty.$$

Proof.

One can show that when $w \in A_2$, we have $\mathcal{P}_{[0,\infty)}^w$ is bounded on $L^p(w)$. So suffices to show $\{\mathcal{P}_{[n+1,\infty)}^w\}_{n \geq 0}$ are bounded on $L^p(w)$.

$$\begin{cases} \mathcal{P}_{[n+1,\infty)}^w &= (I - \mathcal{P}_{[0,n]})\mathcal{P}_{[0,\infty)}^w + \mathcal{P}_{[0,n]}\mathcal{P}_{[n+1,\infty)}^w & \text{("monic")} \\ 0 &= \mathcal{P}_{[0,n]}w\mathcal{P}_{[n+1,\infty)}^w & \text{(orthogonality)} \end{cases}$$



Theorem (A.-Aptekarev-Denisov, '20)

If $w \in A_2$, then there exists $\delta > 0$ such that whenever $|\frac{1}{p} - \frac{1}{2}| < \delta$, then

$$\sup_n \|\mathcal{P}_{[0,n]}^w\|_{L^p(w) \rightarrow L^p(w)} < \infty.$$

Proof.

One can show that when $w \in A_2$, we have $\mathcal{P}_{[0,\infty)}^w$ is bounded on $L^p(w)$. So suffices to show $\{\mathcal{P}_{[n+1,\infty)}^w\}_{n \geq 0}$ are bounded on $L^p(w)$.

$$\begin{cases} \mathcal{P}_{[n+1,\infty)}^w &= (I - \mathcal{P}_{[0,n]})\mathcal{P}_{[0,\infty)}^w + \mathcal{P}_{[0,n]}\mathcal{P}_{[n+1,\infty)}^w & \text{("monic")} \\ 0 &= \mathcal{P}_{[0,n]}w\mathcal{P}_{[n+1,\infty)}^w & \text{(orthogonality)} \end{cases}$$

Now we can proceed exactly as before!



Thank you for Listening!